

**WATER PRODUCTIVITY FOR *BORO* RICE PRODUCTION: STUDY ON
FLOODPLAIN *BEELS* IN RAJSHAHI, BANGLADESH**

**Md. Istiaque Hossain^{*1,2}, Chamhuri Siwar¹, Mazlin Bin Mokhtar¹, Madan
Mohan Dey³, Abd. Hamid Jaafar⁴ and Md. Mahmudul Alam¹**

¹*Department of Fisheries, University of Rajshahi, Rajshahi-6205, Bangladesh*

²*Universiti Kebangsaan Malaysia (UKM) 43600 UKM BANGI, Selangor,
Malaysia*

³*Aquaculture Economics and Marketing Aquaculture/Fisheries Center University
of Arkansas Pine Bluff, USA*

⁴*Graduate Program Coordinator Faculty of Business, Universiti Kebangsaan
Malaysia*

Citation Reference:

Hossain, M.I., Siwar, C., Mokhtar, M., Dey, M.M., Jaafar, A.H., and Alam, M.M. 2013. Water Productivity for Boro Rice Crop Production: Study on Floodplain Beels in Rajshahi, Bangladesh. *Journal of Bio-Science*, 21: 123-136. Available at <http://www.banglajol.info/index.php/JBS/article/view/22526>

**This is a pre-publication copy.
The published article is copyrighted by the publisher of the journal.**

* Corresponding author: E-mail: bitanrubd@yahoo.com , bitan@ru.ac.bd

WATER PRODUCTIVITY FOR *BORO* RICE PRODUCTION: STUDY ON FLOODPLAIN *BEELS* IN RAJSHAHI, BANGLADESH

Md. Istiaque Hossain^{*1,2}, Chamhuri Siwar¹, Mazlin Bin Mokhtar¹, Madan Mohan Dey³,
Abd. Hamid Jaafar⁴ and Md. Mahmudul Alam¹

¹Department of Fisheries, University of Rajshahi, Rajshahi-6205, Bangladesh

²Universiti Kebangsaan Malaysia (UKM) 43600 UKM BANGI, Selangor, Malaysia

³Aquaculture Economics and Marketing Aquaculture/Fisheries Center University of Arkansas Pine Bluff, USA

⁴Graduate Program Coordinator Faculty of Business, Universiti Kebangsaan Malaysia

Abstract

Context: Water productivity is considered as an important indicator of Agriculture productivity because of the scarcity of freshwater. More yield or output against same or less amount of water has become the global interest.

Objectives: This study measures the productivity of water on the floodplain land in terms of *Boro* rice cultivation for two floodplain *beels* in Rajshahi Bangladesh.

Materials & Methods: For this study, the production and market price data were collected by direct observation based on 30 samples in the year 2006-07.

Results: This study found gross water productivity of rice yield as 0.47 kg m⁻³ in *beel* Mail and 0.43 kg m⁻³ in *beel* Chandpur. In monetary value, water productivity per cubic meter irrigation water were TK 5.65, TK 3.42 and TK 2.64 based on gross return, net return considering cash costs and net return considering full costs in *beel* Mail. In *beel* Chandpur these values were TK 5.19 m⁻³, TK 2.87 m⁻³ and TK 2.14 m⁻³, respectively. The usage of average irrigated water in the *boro* rice farms were estimated 10730.05 m³ and 11236 m³ with an average production of yield 4992.95 kg and 4783.20 kg in *beel* Mail and *beel* Chandpur. Statistical result shows that keeping irrigation water constant, a 1% increase of *boro* rice yield will increase water productivity at 0.916% in *beel* Mail and 0.972% in *beel* Chandpur. The water productivity in *beel* Mail was 4.65% higher than *beel* Chandpur due to the intervention of community based fish culture management.

Conclusion: The findings of this study will help to govern and improve production by proper utilizing floodplain lands.

Keywords: water productivity; *Boro* production; Floodplain; Bangladesh; *beel*.

Introduction

Water productivity (WP) generally refers to productivity against unit amount of water used. So it implies to efficient water use that can be increased by either producing more output per unit of water used or by reducing water losses or by the combination of both. Therefore, the concept of WP in Agricultural production systems has focused on 'producing more food with the same water resources' or 'producing the same amount of food with less water resources'.

Freshwater resources are facing increased competition among a multitude of users (Pimentel *et al.* 2004, Rijsberman 2006). In this water scarcity conditions, the limited available water should be used more efficiently (Bessembinder *et al.* 2005). There are two key challenges in front of Agriculture- food production needs to enhance for rapidly growing world population and this demand needs to be fulfilled under decreasing water resources. FAO's (2003) report on World Agriculture: towards 2015/30 projects mentioned that global food production will need to increase by 60% to close nutrition gaps, cope with the population growth and accommodate changes in diets over the next three decades. Water withdrawals for Agriculture

* Corresponding author: E-mail: bitanrubd@yahoo.com , bitan@ru.ac.bd

are expected to increase by some 14% in that period, representing an annual growth rate of 0.6%, down from 1.9% in the period 1963-1999. Much of the increase will take place on arable irrigated land, forecast to expand globally from some 2 million sq. km to 2.42 million sq. km. In a group of 93 developing countries water use efficiency in irrigation, i.e. the ratio between water consumption by crops and the total amount of water withdrawn, is expected to grow from an average 38% to 42%.

To meet up the food shortage, it is needed to produce more food from limited water by increasing WP (Dugan *et al.* 2006, Guerra *et al.* 1998). Molden and Saktivadivel (1999) define WP as the value or weight obtained per unit of water used or consumed. Sharma (2006) also defines productivity of water as the benefits derived from use of water and is most often given in terms of mass of produce, or its monetary value, per unit of water. According to him, WP is dependent on several factors, including crop genetic material, water management practices and the economic and policy incentives to produce. Integrated Water Management Institute (IWMI) researchers have been of the view that this definition is scale dependent. For a farmer, it means getting more crops per drop of irrigation water. But at a regional, basin or a country level WP means getting more value per unit of water resources used. Increasing WP is then the business of several actors working in harmony at plant, field, irrigation-system and river basin levels (Sharma 2006). Various stakeholder groups define WP differently according to water use. Molden (1997) defines WP as 'crop production' per unit 'amount of water used'. It can be further defined in several ways according to the purpose, scale and field of analysis (Bastiaanssen *et al.* 2003, Molden *et al.* 2001).

Some researchers have focused on measuring crop WP values, calculated as the ratio of yield over evapotranspiration, at crop or field level for different crops. Zawrat and Bastiaanssen (2004) reviewed 84 literatures which are not older than 25 years and found profound variations of crop WP for unit amount of water depletion. The range of crops WP is very wide (wheat, 0.6–1.7 kg m⁻³; rice, 0.6–1.6 kg m⁻³; cotton_{seed}, 0.41–0.95 kg m⁻³; cotton_{lint}, 0.14–0.33 kg m⁻³ and maize, 1.1–2.7 kg m⁻³) and thus offers tremendous opportunities for maintaining or increasing agricultural production by 20–40% less water resources. Different economic WP was also found by authors in various crops. Molden *et al.* (1998) and Molden *et al.* (2001) found economic WP for wheat crop in South East Asia ranging from \$0.07 to \$0.17 per m³ irrigation water. Whereas Merrett (1997) found, economic WP in Jordan ranged from \$0.03 m⁻³ for wheat to \$0.3 m⁻³ for potato. Thus the WP values differ by crops, environment, socioeconomic condition and farming pattern.

Wetlands are important ecosystem that supports a wide range of biodiversity and provides numerous benefits. These wetlands, especially the flood prone ecosystems have the potential to be utilized for WP enhancing measures. Floodplain is a low lying flat land which is adjacent to a stream or river that experiences occasional or periodic flooding (Powell, 2009). It is a dry area susceptible to being inundated by water from any natural resource. Some places, like Bangladesh, farmlands that become flooded (over 1 meter depth) during wet season are known as floodplain. *Beel* is a very common form of floodplain area in the Ganga-Brahmaputra floodplains of the states of West Bengal, and Assam in India and in Bangladesh. It is formed by inundation of low lying lands during flooding, where some water gets trapped even after flood waters recede back from the floodplains. It also may be caused by rain water during monsoon season. There are two types of *beels* viz., the seasonal *beels* which dry-up annually and the perennial *beels* which retain water round the year. *Beels* may be formed from great river when the mainstream of the river changes the direction leaving a remnant. Again this can be formed as a result of silt deposition in the river bed which makes the river to flow with two parallel streams leaving a land in between. The land between a pair of parallel rivers then forms a ditch or depression which is converted to *beel* during flooding (Alam & Hossain 2007).

Bangladesh has one of the largest and richest floodplain lands in the world (Rahman, 1989, Tsai *et al.* 1993). In Bangladesh 80% of the total land area are said to be floodplain. 60% of the country is less than six meters above sea level (Hofer & Messerli 1997). During the rainy season (July -September) huge quantity of water, enter the country on its way to the Bay of Bengal. At the same time about 90% of annual rainfall also occurs.

For this reason, two thirds of Bangladesh is vulnerable to flooding and almost every year a third to a quarter of the country goes under water (Nishat, 1990). More than half of rice land in Bangladesh (10.2 million hectares) is flooded to the depth of more than 50cm during the rainy season (IFAD & World Fish Center 2002). In 1986, the Master Plan Organization (MPO) for National Planning estimated the Net Cultivable Area (NCA) of Bangladesh to be 9,562,402 hectares, out of which 6,300,723 hectares were said to be floodplains susceptible to annual submersion to different depths. Whereas, Welcomme (1979) estimated 9,300,000 hectares of floodplain area in Bangladesh, which includes 2,834,000 hectares of paddy fields.

Therefore, proper utilization of water is very important for sustainable Agriculture and rural economic development. At the same time, new policies and initiatives are required for the adaptation of WP enhancing measures (FAO 2001). With this view in mind, assessment of WP of seasonal floodplain water body is very potential to produce more food from the same water resources. Therefore this study is an attempt to fill up the knowledge gap through measuring the productivity of water of the floodplain land in terms of rice production based on two *beels* in Bangladesh.

Methodology

Study Area

The study was conducted on two *beels* which are situated in Mohanpur upazila of Bangladesh. Mohanpur is a - upazila under Rajshahi district. It is situated between 24° 28' and 24° 38' North latitude and between 88° 34' and 88° 43' east latitude having an area of 162.65 sq km (figure 1).




Fig. 1. Study areas in Mohanpur upazila

Source: http://banglapedia.search.com.bd/HT/N_0049.htm

The *beel* Mail is an open *beel* that is shallow depressions where land is converted to Agriculture while permanent water is limited to deeper areas during dry season (Thompson et al. 1999) connected nearby a local Shiba River, which further connected to Padma river basin. This floodplain *beel* is mainly privately owned lands with total 100 hectare area of which 15 hectare government *Khas* lands having water availability for 5-6 month. After flood draw down the *beel* completely drains out the water during dry season leaving fertile land suitable for rice cultivation. Melandi, Goalpara, Dangapara and Moheskundi villagers are the main beneficiaries of this *beel*.

The *beel* Chandpur is also mostly privately owned lands with total 202.43 hectares of land of which 9.15 hectares of land are Government's khas land. The depth of water is 1.5 to 3.5 meters having 5-6 months of water during the wet season. This *beel* is connected with Barnoi River, which further connected to Padma basin. Farmers mainly grow *Boro* rice at the dry season and some are growing Amon crops in the low level area during the wet season. There are 5 villages surrounding the *beel* area namely Batupara, Horiphala, Chandpur, Chuniapara and Nondonhut.

Data Collection

In the study *beels* farmers started *Boro* rice cultivation in December 2006. The land soaking, land preparation and transplantation of each site were done from 3rd week of December to 2nd week of January 2007. The growth of rice crop was seen from 2nd or 3rd week of January to 3rd or 4th week of April 2007. As a sample of 30 farms were taken at each site for monitoring production, and the yields were valued with prevailing market price. Irrigation and rainfall water were normally used for rice crop cultivation, but the rainfall water was found to be very negligible for *Boro* rice crop cultivation during dry season. So the amount of rainfall water and non-beneficial water consumption (loss of water for irrigation system) in rice farms were not considered for measuring WP in this study.

The monitoring of WP by *boro* rice yield at dry season was divided into two parts, direct observation and interview of landowners of various *boro* rice farms for inputs and outputs of rice production and direct observation of irrigated water amount in rice fields.

Data collection was done separately for large farms, medium farms and small farms, and during four periods namely seedling, land preparation, weeding and harvesting periods. However, 11 visits were done for irrigation water amount monitoring at rice fields during land soaking, vegetation phase, production phase and ripening phase. Production of rice yield and irrigated water application per unit area of rice farms were estimated by interviewing the farm households as well as observing the amount of irrigation water application at the farm level. Per hectare average yield of *boro* rice crop was estimated by the total production *boro* rice yield divided by the total cultivable area in sample rice farms. If the depth of irrigation water per hectare is 1mm, it is then considered that 10³ m water has been used per hectare for irrigation purpose.

Models and Estimations

Water use efficiency (WUE), irrigation efficiency (IE), and water productivity (WP) has been defined according to use, field of research and stakeholder. Crop physiologists defined WUE as carbon assimilated and crop yield per unit of transpiration (Viets, 1962), and then later as the amount of produce (biomass or marketable yield) per unit of evapotranspiration (ET). Taking into account that photosynthesis (and thus dry matter yield) and transpiration are related through the diffusion process of CO₂ and H₂O, the efficiency of crop water use has been defined as,

$$\text{Water efficiency} = \frac{\text{Dry matter growth rate}}{\text{Transpiration rate}} = \frac{Y (\text{kg}^{-1} \text{d}^{-1})}{T (\text{mm d}^{-1})} \quad (1)$$

Irrigation specialists have demonstrated the term WUE to describe how effectively water is delivered to crop and to indicate the amount of water wasted. But this concept provides only a partial view because it does not indicate the benefits produced, nor does it specify that water lost by irrigation is often reused by other uses (Seckler et al. 2003). In daily irrigation practices, WP is a more applicable term in equation 1, as it assimilates the rate of dry matter yield and transpiration over time, i.e. denoted as Y and T respectively.

$$\text{WP}_T = \frac{Y (\text{kg ha}^{-1})}{T (\text{mm})} \rightarrow \frac{Y (\text{kg ha}^{-1})}{T (\text{m}^3 \text{ha}^{-1})} \rightarrow \text{kg m}^{-3} \quad (2)$$

Where, 1 mm is equivalent with 10 m³ ha⁻¹. When applying irrigation at field scale, it is generally difficult to distinguish plant transpiration T (mm) from soil evaporation E (mm). Hence, instead of WP_T, WP_{ET} may be used (Molden, 1997, Molden et al., 2001; Droogers & Bastiaanssen, 2002, Kijne et al., 2003):

$$WP_{ET} = \frac{Y \text{ (kg ha}^{-1}\text{)}}{ET \text{ (m}^3 \text{ ha}^{-1}\text{)}} \rightarrow \text{kg m}^{-3} \quad (3)$$

Where ET is the evapotranspiration of 'crop + soil'. Total dry matter yield Y may also be transformed into marketable yield, i.e. Y_M . If the amount of irrigation and precipitation water is considered as 'water use of the crop' then WP_{I+P} may be used,

$$WP_{I+P} = \frac{Y \text{ (kg ha}^{-1}\text{)}}{[I + P] \text{ (m}^3 \text{ ha}^{-1}\text{)}} \rightarrow \text{kg m}^{-3} \quad (4)$$

Where I stands for amount of seasonal irrigation and P for the seasonal precipitation. Where ER (effective rainfall) is very negligible resulting in very low precipitation like arid regions or for dry season crop, WP_{I+P} may be converted to WP_I (Vazifedoust *et al.*, 2008)

$$WP_I = \frac{Y \text{ (kg ha}^{-1}\text{)}}{I \text{ (m}^3 \text{ ha}^{-1}\text{)}} \rightarrow \text{kg m}^{-3} \quad (5)$$

At the field scale classical WP indicator is crop output and total water supplied (irrigation and effective precipitation) to the field as because it is more appropriate and easy to interpret by management (Senzanje *et al.*, 2005). As the farmer is mainly interested in the economic yield of the crop, WP may be expressed in terms of money as,

$$WP_s = \frac{(\$ \text{ kg}^{-1}) \text{ (kg ha}^{-1}\text{)}}{ET \text{ (m}^3 \text{ ha}^{-1}\text{)}} \rightarrow \$ \text{ m}^{-3} \quad (6)$$

So WP measurement is straight forward in crop per drop analysis, which analyzes output (benefit quantity or value of product), over quantity of input (volume of depleted water). So in dry season, productivity from rice cultivation is expressed by rice yield. The equation 7 measures the WP of crops in physical terms as,

$$WP_y = \frac{Q_y}{WC} \quad (7)$$

Where,

WP_y = water productivity in terms of crops yield

Q_y = yield of crop as physical units

WC = consumption of depleted water, expressed in cubic meters

That is, physical productivity is the quantity (Q_y) of product per unit quantity of depleted water. The equation 8 measures the WP in terms of value as:

$$WP_v = \frac{Q_v}{WC} \quad (8)$$

Where,

WP_v = water productivity in monetary terms of crop

Q_v = gross/ net value of yield output in monetary terms

WC = consumption of depleted water, expressed in cubic meters

Again, it is evident that equation 8 is the more suitable measure in case of multi-product outputs. The value of output, Q_v , is the summation of values of different products resulting from application of unit amount of water.

The WP of rice crop in physical terms as,

$$WP_y \text{ (Kg m}^{-3}\text{)} = \frac{Q_y \text{ (Kg)}}{WC \text{ (m}^3\text{)}} \quad (9)$$

$$WC \text{ (m}^3\text{)} = BWC + NBWC \quad (10)$$

Where,

WP_y = water productivity in physical units in terms of rice yield
 Q_y = yield of rice crop, usually expressed in Kg
 WC = consumption of depleted water, expressed in cubic meters
 BWC = beneficial water consumption in cubic meter
 $NBWC$ = Non-beneficial water consumption in cubic meter

The WP of rice crop in terms of value:

$$WP_y \text{ (TK m}^{-3}\text{)} = \frac{Q_{v1} \text{ (TK)} + Q_{v2} \text{ (TK)}}{WC \text{ (m}^3\text{)}} \quad (11)$$

Where,

WP_y = water productivity in terms of monetary valued
 Q_{v1} = value of net return by rice yield
 Q_{v2} = value of net return of by product
 WC = consumption or depleted water

That is, WP is the value of product per unit quantity of depleted water. The same equation is also applicable for measuring WP of by product of crop. It is evident that equation 11 is the more suitable measure in case of multi-product outputs. The value of output, Q_v , is the summation of values of different products resulting from application of water.

This study used the WP by *boro* rice crop at farm level which was estimated by output of rice yield and it was by product with per cubic meter of irrigation water based on equation 9 to 11.

Assessment of Water Productivity Measurement of Cost of Production

For producing *boro* rice per hectare average costs were calculated from sampled farms of *boro* rice in *beel* Mail and *beel* Chandpur. Per hectare full costs were calculated TK 31168.74 and TK 33237.94 for *beel* Mail and *beel* Chandpur, respectively by adding up all sorts of costs involved in producing rice crop. Whereas, per hectare cash variables costs for *boro* rice were estimated TK 24313.14 and TK 25667.59 which comprised 78% and 77.22% of total costs for *beel* Mail and *beel* Chandpur, respectively (Table 1). Irrigation water cost was found to be the most important cost as 24% and 22% for *beel* Mail and *beel* Chandpur, respectively.

In the winter season, *boro* rice crop is dependent on irrigation because of minimum or negligible rainfall. Without assured irrigation, the production cannot be expected high for this crop. Irrigation cost (calculated based on land size that is 33 decimal land= TK 1000 per season) found to be most important cost for *boro* rice production ranging from 22-24% of total production cost. The costs of irrigation water depend on number of irrigation required for crop cultivation. Among the sample crop farms, per hectare average costs of irrigation were TK 7488.88 in *beel* Mail and TK 7456.04 in *beel* Chandpur (Table 1).

Table 1. Economic analysis of per hectare *boro* rice production in seasonal *beels*.

Items	Mail <i>Beel</i>		Chandpur <i>Beel</i>	
	Value (TK)	% total cost	Value (TK)	% total cost
Cash expenses	24313.14	78.00	25667.59	77.22
Hire Labour cost	6652.84	21.34	7228.35	21.75
Fertilizers	5561.00	17.84	5514.04	16.59
DAP	2705.00	-	2579.70	-

Urea	1737.00	-	1507.24	-
MP	1119.00	-	1354.90	-
Cow dung	0	-	72.20	-
Cost of seeds	1562.93	5.01	1551.84	4.67
Tractor use	2246.75	7.21	3059.06	9.20
Pesticides	800.83	2.57	858.27	2.58
Water cost	7488.80	24.03	7456.04	22.43
Non cash expenses	6855.60	22.00	7570.34	22.78
Lease values of land	5625.03	18.05	5624.41	16.92
Family labour	774.55	2.49	1464.57	4.41
Interest on operating capital	456.02	1.46	481.36	1.45
Full cost (TK)	31168.74	100	33237.94	100

Source: Field survey data

Human labour cost is one of the most important variables input in the production process. Labour cost was found to be second most important production cost of *boro* rice cultivation. Both hired and family supplied labours were used for seedling, land preparation, plantation, fertilizers and pesticide uses, weeding and harvesting, threshing and carrying and drying and storage for production of *boro* rice crop (Table 1). To determine total labour cost unpaid family labour cost was taken into account. Table 2 shows the operation-wise distribution of family and hired labour for per hectare crop production. For dry seasons per hectare crop production total hired labour cost was TK 6652.84 (21.34% of total cost) and family labour cost was TK 774.55 (2.49% of total cost) in *beel* Mail. In *beel Chandpur* for per hectare crop production total hired labour cost was TK 7228.35 (21.75% of total cost) and family labour cost was TK 1464.57 (4.41% of total cost).

Table 2. Operation wise distribution of human labour costs for *boro* rice production.

Operations	Mail <i>beel</i>				Chandpur <i>Beel</i>			
	Family labour		Hire labour		Family labour		Hire labour	
	TK	%	TK	%	TK	%	TK	%
Land preparation	336	43.35	807	12.13	603	41.16	1478	20.45
Transplanting	78	10.06	1301	19.56	138	9.42	1390	19.23
Weeding	62	8	2168	32.59	115	7.85	2107	29.15
Fertilizer used	118	15.23	0	0	242	16.52	0	0
Insecticide used	56	7.23	0	0	89	6.08	0	0
Harvesting, threshing and carrying	58	7.48	1774	26.66	172	11.74	1738	24.05
Drying and storage	67	8.65	523	7.86	106	7.24	420	5.81
Others	0	0	80	1.20	0	0	95	1.31
Total	775	100	6653	100	1465	100	7228	100

Source: Field survey data

Among the raw materials, fertilizer is another important cost item for *boro* rice crop cultivation. In the study area farmers used DAP, urea, TSP, potash (K) and cow-dung as fertilizer (organic and nonorganic). Table 1 shows per hectare average costs of fertilizer usage at *boro* rice crop farms. Among the sample crop farms average per hectare fertilizer costs were TK 5561 and TK 5514.04 which were 22% and 20% of total production cost for *beel* Mail and *beel Chandpur*, respectively. Costs of seed depend on variety or number of plants planted in the rice field by the farmers. In the sample crop farms, per hectare average cost of seeds were TK 1562.93 and TK 1551.84 which occupied 5.01% and 4.67% of total production cost for *beel* Mail and *beel Chandpur*, respectively (Table 1). Farmers used tractor for preparation of land for *boro* rice crop cultivation. This cost depends on work hour of tractor use for land preparation. Among the farms, per hectare

average costs of tractor use were TK 2246.75 and TK 3059.06 which shared 7.21% and 9.2% of total costs of *boro* production for *beel* Mail and *beel* Chandpur respectively (Table 1).

Pesticides were used by the farmers during weeding period. Among the sample farms per hectare average costs of pesticide for *boro* rice crop were TK 800.83 and TK 858.27 which shared 2.57% and 2.58% of total production costs for *beel* Mail and *beel* Chandpur, respectively (Table 1).

The amount of money needed to meet the expenses of hired or purchased inputs such as, human labour, seeds, fertilizers, tractor use, pesticides and leveling etc. for *boro* rice cultivation were treated as operating capital. Interest on operating capital (IOC) was charged @ 10% annually and was estimated for the period which the operating capital was used. IOC was estimated by using the following formula (Miah, 1987).

$$IOC = \frac{\text{Operating capital} \times \text{Rate of interest} \times \text{Time consideration}}{2} \quad (12)$$

IOC for *boro* rice production were estimated at TK 456.02 and TK 481.36 ha⁻¹ and it is constituted 1.46% and 1.45% of the total cost for *beel* Mail and *beel* Chandpur respectively (Table 1).

The value of land has been estimated according to average lease value (local price) of seasonal floodplain land of the farmers. Land is usually charged for 1 season or four and half months. During study period, per hectare *boro* rice land was charged TK 5625.03 and 5624.41 for 4.5 months that shared 18.05% and 16.92% of all cost in *beel* Mail and *beel* Chandpur respectively (Table 1). Cost of land (rent) was considered as opportunity cost as some of the sample farms were privately owned and did not require land value.

Measurement of Returns of *boro* Production

Per hectare average yields of *boro* rice were 4992.95±157.95 kg and 4783.20±117.15 kg in *beel* Mail and *beel* Chandpur, respectively. Average per hectare gross returns from sample *boro* rice farms were calculated TK 59873.31±1844.82 and TK 57642.10±1421.09 for *beel* Mail and *beel* Chandpur respectively by value of rice yields and its by-products. The price rates were measured according to prevailing market price (Table 3 and Figure 2).

Table 3. Per hectare production of *boro* rice in seasonal rice field.

Particulars	<i>Beel</i> Mail		<i>Beel</i> Chandpur	
	Production	SD	Production	SD
Average gross returns (TK)	59873.31	1844.82	57642.10	1421.09
Rice yield by value (TK)	54922.41	1736.92	52615.20	1287.64
By product of rice (TK)	4950.90	118.77	5026.90	284.11
Gross average cash cost (TK)	24313.14	1687.50	25667.59	1061.4
Gross average full cost (TK)	31168.74	1611.52	33237.94	1931.99
Net returns basis on cash cost (TK)	35560.17	2317.93	31974.54	790.99
Net returns basis on full cost (TK)	28704.56	1144.68	24404.19	1115.67

Source: Field survey data

Measurement of Net Benefits of *boro* Production

The per hectare returns over cash variables cost that is net profit of rice crop at *beel* Mail and *beel* Chandpur were TK 35560.17±2317.93 and TK 31974.54±790.99, respectively. Whereas it was TK 28704.56±1144.68 and TK 24404.19±1115.67 for *beel* Mail and *beel* Chandpur respectively on the basis of full cost. The net return of rice crop was found higher in the *beel* Mail than *beel* Chandpur (Table 3 and Figure 2).




Fig. 2. Per hectare costs and returns of *boro* rice crop in sites of *beel*.

Measurement of Usage of Water

The amount of irrigation water required for crop cultivation was observed during land soaking, vegetation phase, production phase and ripening phase. If the depth of irrigation water per hectare is 1mm, it is then considered that 10m³ of water has been used in per hectare area for irrigation purpose. The estimated irrigation water volume applied to the *boro* rice farms are summarized in Table 4.

Table 4. Average irrigated water (mm) for *boro* rice cultivation at different stages.

Growth (week)	Cultivation period (week)	Water depth (mm)		Different phases of crop cultivation
		Mail <i>beel</i>	Chandpur <i>beel</i>	
		50	100	Soaking, land preparation, transplanting
1	Jan 3rd	71.00	72	Vegetation phase
2	Jan 4th	92.33	88	
3	Feb 1st	115.00	109	
4	Feb 2nd	131.00	128	
5	Feb 3rd	132.33	133	Reproductive phase
6	Feb 4th	110.33	114	
7	Mar 1st	63.67	61	
8	Mar 2nd	72.33	76	
9	Mar 3rd	127.67	133.6	Ripening phase
10	Mar 4th	107.33	109	
11	Apr 1st	0	0	No irrigation applied
12	Apr 2nd	0	0	
13	Apr 3rd	0	0	
14	Apr 4th	0	0	Harvest
	Total	1073	1123.6	

Source: Field survey data

After seven and eight weeks of transplanting, farmers commonly apply less water to encourage tillering of rice plants. At *beel* Mail, water depth was 50 mm during land soaking and land preparation, and 1023 mm for crop growth. In *beel* Chandpur water depth was 100 mm during land soaking and land preparation, and 1023.6 mm for crop growth (Table 4).

Measurement of Water Productivity

After flood drawdown, rice crop cultivation on dry land of *beels* is mostly dependent on irrigation due to minimum or no rainfall during this period. The abstraction of water used at rice farms were directly related to deep tube-well (underground water) or both underground and surface water of floodplain *beels*.

Almost all of the cases the irrigated water was collected from Deep tube-well water (DTW). Amount of water consumption for rice crop was measured by millimeter (mm) of irrigated water. Table 5 shows the average water consumption at per hectare area for *boro* crop cultivation. The volumes of water consumption at farm level of *beel* Mail and *beel* Chandpur were 10730.5 m³ ha⁻¹ and 11236 m³ ha⁻¹ respectively.

Table 6 shows the WP by irrigated *boro* rice in winter season (dry *beel*) at the rice farms of *beel* Mail and *beel* Chandpur. Physical WP value ranged from 0.41 to 0.50 Kg m⁻³ among the selected rice farms in two floodplain *beels*. The range of physical WP was 0.44-0.50 Kg m⁻³ (average 0.47±0.018Kg m⁻³) at *beel* Mail, with an average yield of 4992.95±157.92 kg and WC ha⁻¹ of 10730.5 m³ and at *beel* Chandpur range of physical WP was 0.41-0.47 Kg m⁻³(average 0.43±0.014Kg m⁻³) with an average yield of 4783.20±117.15 kg and WC ha⁻¹ of 11236 m³.

However, in *beel* Mail, WP values in monetary term were estimated TK 5.65±0.212 m⁻³, TK 3.42±0.243 m⁻³ and TK 2.64±0.109 m⁻³ for rice crop and its by-product based on gross returns, net return based on cash costs and net return based on full costs respectively. WP values in *beel* Chandpur were TK 5.19±0.163 m⁻³, TK 2.87±0.085 m⁻³ and TK 2.14±0.098 m⁻³ for rice crop and its byproduct based on gross returns, net return based on cash costs and net return based on full costs respectively (Table 6).

Table 5. Per hectare water consumption of *boro* rice.

Name of <i>beels</i>	Average water consumption per hectare		
	Water depth(mm)	SD	Volume of water (m ³)
<i>Beel</i> Mail	1073.05	166.58	10730.5
<i>Beel</i> Chandpur	1123.6	81.07	11236

Source: Field survey data

Table 6. Water productivity by *boro* rice crop in two *beels*.

Water productivity	<i>Beel</i> Mail		<i>Beel</i> Chandpur	
	WP (m ⁻³)	SD	WP (m ⁻³)	SD
WP range by rice yield (Kg)	0.44-0.50	-	0.41-0.47	-
WP by rice yield (Kg)	0.47	0.018	0.43	0.014
WP by gross return (TK)	5.65	0.212	5.19	0.163
WP by net return based on cash cost(TK)	3.42	0.243	2.87	0.085
WP by net return based on full cost (TK)	2.64	0.109	2.14	0.098

Source: Field survey data

Correlation among the Variables

Pearson correlation analysis shows that WP by *boro* rice yield in *beel* Mail has a positive significant correlation with rice yield ($r=0.916$, $p<0.01$) and negative significant correlation with water volume ($r=-0.583$, $p<0.01$) (Table 7).

Table 7. Correlation and significant test of water productivity with water volume and returns in *Beel* Mail.

Particulars	WV	RY	GR	NRBCC	NRBFC	WPBRY	WPBGR	WPBNCC
RY	-0.242							
GR	-0.252	1.00**						
NRBFC	-0.401*	0.697**	0.693**					
NRBFC	0.107	0.509**	0.501**	0.202				
WPBRY	-0.583**	0.916**	0.919**	0.743**	0.352			
WPBGR	-0.618**	0.912**	0.917**	0.732**	0.362*	0.989**		
WPBNCC	-0.576**	0.671**	0.670**	0.980**	0.156	0.788**	0.785**	
WPBNFC	-0.273	0.583**	0.579**	0.340	0.927**	0.559**	0.582**	0.362*

* Correlation is significant at the 0.05 level (2-tailed), ** Correlation is significant at the 0.01 level (2-tailed).

WV= Water volume, RY= Rice yield, GR= Gross return, NRBCC= Net return basis of cash cost, NRBFC= Net return basis of full cost, WPBRY= WP by rice yield (Kg m⁻³), WPBGR= WP by gross return (TK m⁻³), WPBNCC= Net WP basis of net cash cost (TK m⁻³), WPBNFC= Net WP basis of net full cost

Moreover, at *beel Chandpur* WP has positive significant correlation with rice yield ($r=0.972$, $p<0.001$), whereas negative significant correlation with volume of irrigated water ($r=-0.933$, $p<0.001$) (Table 8).

Again WP by *boro* rice of *beel* Mail has a positive significant correlation with gross return ($r=0.917$, $p<0.01$) and net return based on cash cost ($r=0.980$, $p<0.01$). However, water volume shows negative significant correlation with gross return ($r=-0.576$, $p<0.01$) and net return based on cash cost ($r=-0.576$, $p<0.01$) (Table 7). Again at *beel Chandpur*, WP has a positive significant correlation with gross return ($r=0.996$, $p<0.01$) and net return based on cash cost ($r=0.950$, $p<0.01$). However, water volume shows negative significant correlation with gross return ($r=-0.964$, $p<0.01$) and net return based on cash cost ($r=-0.719$, $p<0.01$) (Table 8). Here the sign of correlation of “r” is positive means return will increase with increase of inputs and vice versa.

Table 8. Correlation and significant test of water productivity with water volume and returns in *Beel* Chandpur.

Particulars	WV	RY	GR	NRBCC	NRBFC	WPBRY	WPBGR	WPBNCC
RY	-0.917**							
GR	-0.937**	0.983**						
NRBFC	-0.548**	0.638**	0.675**					
NRBFC	0.188	-0.165	-0.148	0.150				
WPBRY	-0.933**	0.972**	0.965**	0.653**	-0.182			
WPBGR	-0.964**	0.977**	0.996**	0.655**	-0.158	0.969**		
WPBNCC	-0.719**	0.771**	0.805**	0.975**	0.073	0.787**	0.796**	
WPBNFC	0.041	-0.035	-0.014	0.224	0.989**	-0.049	-0.019	0.174

** Correlation is significant at the 0.01 level (2-tailed).

Discussions

This study found gross WP of rice yield at 0.47 ± 0.018 kg m⁻³ in *beel* Mail and 0.43 ± 0.014 kg m⁻³ in *beel Chandpur* during dry season, which is also similar to the findings of Cai, Ximing and Rosegrant (2003). They showed that world WP of rice yield in 1995 was 0.15 to 0.60 Kg m⁻³ which will increase from 1995 to 2025. The findings of this study are a little higher than the WP value in Pakistan but lower than the findings in Philippines and China. The WP values for gross inflow of irrigation were 0.27 kg m⁻³ ranging from 0.20 kg m⁻³ to 0.31 kg m⁻³ in Punjab, Pakistan. The irrigation WP of traditional rice cultivation in Tuanlin China was 1.95

kg m⁻³ (Dong *et al.* 2001) and in Philippines for wet seeded rice was 1.4 to 1.6 kg m⁻³ (Toung and Bouman 2002). This variation may be due to variations of rice crop cultivated, land, and usage of fertilizer, environment and cultivating technology.

Again in dry season WP from per cubic meter irrigation water in monetary term of *beel* Mail were TK 5.65±0.212, TK 3.42±0.243 and TK 2.64±0.109 based on gross return, net return considering cash costs and net return considering full costs. However, in *beel Chandpur* these values were TK 5.19±0.163 m⁻³, TK 2.87±0.085 m⁻³ and TK 2.14±0.098 m⁻³. It was also evident from the study that higher WP was obtained in *beel* Mail area than *beel Chandpur* area in dry season due to higher gross yield of *boro* rice against less amount of irrigation water used. The usage of average irrigation water in the *boro* rice farms were estimated 10730.05±166.58 m⁻³ and 11236±81.07 m⁻³ with an average production of yield 4992.95± 157.92 kg and 4783.20±117.15 kg in *beel* Mail and *beel Chandpur*, respectively. Statistical correlation indicates that 1% increase of rice yield of *boro* rice crop, keeping irrigation water constant at farm levels in *beel* Mail will increase 0.916% of WP, whereas 1% volume of irrigation water increase, keeping rice yield of rice crop constant, would result in decrease of WP by 0.583%. Again at *beel Chandpur*, 1% increase of rice yield of *boro* rice crop, keeping irrigation water constant, will increase 0.972% of WP, and 1% volume of irrigation water increase, keeping rice yield of rice production constant, would result in decrease WP by 0.933%.

WP of *boro* rice yield per cubic meter of water use was 4.65% higher in *beel* Mail than *beel Chandpur*. Production cost considering material input costs were also found less in *beel* Mail specially tractor and labeling cost during land preparation resulting in higher net WP. Per hectare land preparation cost was TK 4769 which shared less than 40% of the production cost in *beel* Mail whereas the cost was TK 6668 which shared above 60% of the production cost for *beel Chandpur*. Per hectare cost for land preparation was more than 28% higher in *beel Chandpur* compared to *beel* Mail.

Conclusions

It was found that WP values were higher in *beel* Mail than the *beel Chandpur* due to intervention of community based fish culture. In dry season land preparation cost was less in *beel* Mail area because water was reserved for fish culture till *boro* rice cultivation so that land was swallow and no weed.

Acknowledgements

The authors would like to acknowledge the financial support for this research from the World Fish Center (Community Based Fish Culture in Irrigation System and Seasonal Floodplain Project (CP35) funded by CGIAR Challenge Program Water & Food, April 2005 -31 Dec 2010) and Malaysian Commonwealth Scholarship (MCS), Ministry of Higher Education, Malaysia (Ref. KPT (BS) 035/11/002, 15.2.2006-31.5.2009).

References

- Alam MS, Hossain MS. (2007). "Beel". *Banglapedia*. Asiatic Society of Bangladesh.
http://banglapedia.search.com.bd/HT/B_0382.htm. (23rd nov. 2007)
- Bastiaanssen WGM, Van Dam JC, Droogers P. (2003). Introduction. In Dam, J.C. van, and R.S. Malik (Eds.), 2003. Water productivity of irrigated crops in Sirsa district, India. Integration of remote sensing, crop and soil models and geographical information systems. *WATPRO final report*, including CD-ROM. 6, 11-19.
- Besseminder JJE, Leffelaar PA, Dhindwal AS, Ponsioen TC. (2005). Which crop and which drop, and the scope for improvement of water productivity. *Agric. Water Manage.* 73 (2), 113–130.
- Cai, X. and Rosegrant, M. (2003). World water productivity: Current situation and future options. In: Kijne, J.W., Barker, R. and Molden, D. (Eds.) *Water Productivity in Agriculture: Limits and Opportunities for Improvement*. CABI, Wallingford, 163-178.
- Dong B, Loeve R, Li YH, Chen CD, Molden D. (2001). Water productivity in Zhanghe irrigation system: Issues of scale. *Proceedings of an international workshop on water saving irrigation for rice*. Wuhan, China.

- Droogers P, Bastiaanssen WGM. (2002). Irrigation performance using hydrological and remote sensing modelling. *J. of Irrigation and Drainage Engineering* 128, 11-18.
- Dugan PJ, Dey MM, Sugunan VV. (2006). Fisheries and Water productivity in tropical river basins: enhancing food security and livelihoods by managing water for fish. *Agricultural Water Productivity*. 80, 262-275.
- FAO. (2001). *The economics of conservation agriculture*. Rome, Italy: FAO, Land and water development division, Food and Agriculture organization of the United Nations.
- FAO. (2003). Raising water productivity. Issues in world agriculture 2001-2004, Spotlight 2001-2004. <http://www.fao.org/AG/magazine/0303sp2.htm>, (4th July 2007) pp-?
- Guerra LC, Bhuiyan SI, Tuong TP, Barker R. (1998). Producing more rice with less water from irrigated systems. *SWIM Paper 5*. Sri Lanka, Colombo: International Water Management Institute.
- Hofer T, Messerli B. (1997). Floods in Bangladesh; Process Understanding and Development Strategies. Institute of Geography, Univ. of Beren, Switzerland. <http://www.blackwell-synergy.com/doi/pdf/10.1046/j.1365-2400.2003.00339.x>, (5th May 2007).
- IFAD, WorldFish Center (2002). Increasing and Sustaining the Productivity of Fish and Rice in the Flood-prone Ecosystems in South and Southeast Asia.
- Kijne J, Barker R, Molden D. (Eds.) (2003). Water productivity in agriculture: limits and opportunities for improvement. Comprehensive assessment of Water Management in Agriculture, Series No. 1, *CABI press*, Wallingford, UK, 352 pp.
- Merrett S. (1997). Introduction to the economics of water resources: An international perspective. London: University College London Press.
- Miah MTH. (1987). Appraisal of deep shallow tube well irrigation project in the Tangail District in Bangladesh, *MSC Dissertation*. University of New England, Armidale, Australia.
- Molden D, Sakthivadivel R. (1999). Water Accounting to Assess Use and Productivity of Water. *International Journal of Water Resources Development*, 15, 55-71.
- Molden D. (1997). Accounting for water use and productivity. *SWIM Paper 1*. Colombo, Sri Lanka: International Irrigation Management Institute.
- Molden D, Murray-Rust H, Sakthivadivel R, Makin I. (2001). A water productivity framework for understanding and action. *Workshop on Water productivity*. Wadduwa, Sri Lanka, November 12 and 13, 2001.
- Molden D, Sakthivadivel R, Perry CJ, de Fraiture, C. Kloezen, WH. (1998). Indicators for comparing performance of irrigated agricultural systems. *Research report 20*. Colombo, Sri Lanka: International Water Management Institute.
- MPO. (1986). *National Water Plan. Sector Analysis. Master Plan Organization (MPO) Technical Report No. 1*, Ministry of Water Resources, Dhaka.
- Nishat A. (1990). Background Paper on Water Resources and Flood Control. Department of Water Resources Engineering, Bangladesh University of Engineering and Technology, Dhaka.
- Pimentel D, Berger B, Filiberto D, Newton M, Wolfe B, Karabinakis E, Clark, S, Poon, E, Abbett, E, Nandagopal S. (2004). Water resources: agricultural and environmental issues. *Bioscience*. 54 (10), 909-918.
- Powell W, Gabe. (2009). Identifying Land Use/Land Cover (LULC) Using [National Agriculture Imagery Program](http://www.fws.gov/naiip/) (NAIP) Data as a Hydrologic Model Input for Local Flood Plain Management. Applied Research Project, Texas State University. <http://ecommons.txstate.edu/arp/296>, (6th May 2007).
- Rahman AKA. (1989). *Freshwater of Bangladesh*. Zoological Society of Bangladesh, Dhaka, 364-366 pp.
- Rijsberman FR. (2006). Water scarcity: fact or fiction? *Agricultural Water Management*. 80, 5-22.

- Seckler D, Molden D, Sakthivadivel R. (2003). *The concept of efficiency in water-resources management and policy*. In: Kijne et al. eds. 2003. *Water Productivity in Agriculture: Limits and Opportunities for Improvement*. CABI Publishing, Wallingford, UK, 37-51.
- Senzanje A, Chimunhu T, Zirebwa J. (2005). Assessment of water productivity trends for parastatal agricultural operations—case of Middle Sabi Estate, Zimbabwe. *Physics and Chemistry of the Earth*. 30, 767–771.
- Sharma BR. (2006). Crop Water Requirements and Water Productivity: Concepts and Practices, presented on October 30, 2006; 1200-1300 hrs, in College of Agricultural Engineering, Punjab Agricultural University, Ludhiana.
- Thompson PM, Sultana P, Islam MN, Kabir MM, Hossain MM, Kabir MS. (1999). An assessment of co-management arrangements developed by the Community Based Fisheries Management Project in Bangladesh, Paper Presented at the *International Workshop on Fisheries Co-management*, 23–28 August 1999, Penang, Malaysia. Toung and Bouman 2002.
- Toung TP, Bouman BMA. (2002). Rice production in water scarce environments. In *Water productivity in agriculture: limits and opportunities for improvement*, Ed. J. W. Kijne. Wallingford, UK: CABI.
- Tsai CF, Ali MY, Ahsan K, Alam SS. (1993). 'Household fish consumption in relation to fishing practices and fish production in flood plains of Bangladesh'. *Indian Fisheries Journal*. 40(3), 181-188.
- Vazifedoust M, Van Dam JC, Feddes RA, Feizi M. (2008). Increasing water productivity of irrigated crops under limited water supply at field scale. *Agricultural water management*. 95, 89 – 102.
- Viets Jr, FG. 1962. Fertilizers and the efficient use of water. *Advances in Agronomy*. 14, 223–264.
- Welcomme RL. (1979). *Fisheries Ecology of Floodplain Rivers*. Longman, London.
- Zwart SJ, Bastiaanssen, WGM. (2004). Review of measured crop water productivity values for irrigated wheat, rice, cotton and maize. *Agricultural Water Management*. 69(2), 115-133.