

IMPACTS OF LOW, MODERATE, AND HIGH SEVERITY FIRE ON
HERPETOFAUNA AND THEIR HABITAT IN A SOUTHERN
USA MIXED PINE/HARDWOOD FOREST

DISSERTATION

Presented to the Graduate Council of
Texas State University-San Marcos
in Partial Fulfillment of
the Requirements

for the Degree

Doctor of PHILOSOPHY

by

Donald J. Brown, B.S., M.S.

San Marcos, Texas
August 2013

IMPACTS OF LOW, MODERATE, AND HIGH SEVERITY FIRE ON
HERPETOFAUNA AND THEIR HABITAT IN A SOUTHERN
USA MIXED PINE/HARDWOOD FOREST

Committee Members Approved:

Michael R. J. Forstner, Chair

John T. Baccus

D. Bruce Means

Weston H. Nowlin

Todd M. Swannack

Approved:

J. Michael Willoughby
Dean of the Graduate College

COPYRIGHT

by

Donald James Brown

2013

FAIR USE AND AUTHOR'S PERMISSION STATEMENT

Fair Use

This work is protected by the Copyright Laws of the United States (Public Law 94-553, section 107). Consistent with fair use as defined in the Copyright Laws, brief quotations from this material are allowed with proper acknowledgment. Use of this material for financial gain without the author's express permission is not allowed.

Duplication Permission

As the copyright holder of this work I, Donald James Brown, authorize duplication of this work, in whole or in part, for educational or scholarly purposes only.

DEDICATION

This dissertation is dedicated to the species that have little or no monetary, medicinal, or recreational value, and thus no voice with which to fight for their inherent worth and right to exist. It is up to us to fight for them.



ACKNOWLEDGEMENTS

I am grateful to my advisor Michael R. J. Forstner for giving me the opportunity to be involved in this important research, and for fostering my development as a wildlife ecologist. Thank you to my committee members John T. Baccus, D. Bruce Means, Weston H. Nowlin, and Todd M. Swannack for their knowledge and assistance with the completion of this dissertation. I am grateful to the Capital Area Council of the Boy Scouts of America for allowing the Forstner Lab to use and manipulate the Griffith League Ranch for ecological research. I am indebted to all of the individuals who contributed to this dissertation and our Houston toad conservation research program in the Lost Pines ecoregion. I especially thank Ivana Mali, Melissa Jones, Devin Preston, Emrah Ozel, Bei DeVolld, Dionne Episcopo, and Adam Duarte, who contributed substantial effort to these projects and co-authored the resulting manuscripts. This research was supported by the Texas Parks and Wildlife Department and U. S. Fish and Wildlife Service through a Traditional Section 6 grant, the National Science Foundation through a GK-12 fellowship, Texas State University through research and instructional assistantships, and research scholarships through Texas State University and the Texas Chapter of the Wildlife Society.

This manuscript was submitted on June 18, 2013.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS.....	vi
LIST OF TABLES.....	xi
LIST OF FIGURES	xiii
ABSTRACT.....	xvi
CHAPTER	
I. INTRODUCTION.....	1
Literature Cited	6
II. POTENTIAL POSITIVE EFFECTS OF FIRE ON JUVENILE AMPHIBIANS IN A SOUTHERN USA MIXED PINE/HARDWOOD FOREST	11
Abstract	12
Introduction.....	13
Study Site	15
Methods.....	17
Sampling design.....	18
Statistical analyses	22
Results.....	24
Discussion	28
Acknowledgements.....	32
References.....	32

III. IMPACTS OF A HIGH SEVERITY WILDFIRE ON ABUNDANCE, MOVEMENT, AND DIVERSITY OF HERPETOFAUNA IN THE LOST PINES ECOREGION OF TEXAS	41
Abstract	42
Introduction.....	43
Materials and Methods.....	45
Study site.....	45
Fire severity	48
Herpetofaunal sampling.....	48
Statistical analyses	50
Results.....	53
Fire severity	53
Herpetofaunal impacts	55
Lizard abundance and detection.....	57
Hurter’s Spadefoot Toad detection	59
Herpetofaunal species composition	62
Species richness of calling anurans.....	62
Discussion	64
Acknowledgements.....	68
Literature Cited	68
IV. COMPARISON OF LOW, MODERATE, AND HIGH SEVERITY FIRE IMPACTS TO AQUATIC AND TERRESTRIAL ECOSYSTEM COMPONENTS OF A SOUTHERN USA MIXED PINE/HARDWOOD FOREST	77
Abstract	78
Introduction.....	80

Methods.....	85
Study area.....	85
Fires.....	86
Prescribed burns.....	86
Wildfires	86
Data collection	89
Water quality.....	89
Aquatic arthropods.....	91
Terrestrial vegetation	92
Ground-dwelling arthropods	93
Data analyses	94
Water quality.....	94
Aquatic arthropods.....	96
Terrestrial vegetation	97
Ground-dwelling arthropods.....	99
Results.....	101
Water quality.....	101
Aquatic arthropods.....	104
Terrestrial vegetation	104
Fire severity	104
Forest structure characteristics and understory community composition.....	108
Ground-dwelling arthropods.....	111
Discussion	114
Acknowledgements.....	121
References.....	122

V. FIRE ANTS IN HOUSTON TOAD HABITAT: ANNUAL ACTIVITY AND RESPONSES TO CANOPY COVER AND FIRE.....	136
Abstract	137

Introduction.....	138
Study Site	141
Methods.....	142
Statistical analyses	143
Results.....	145
Discussion	148
Acknowledgements.....	151
References.....	151

VI. WILDFIRE IMPACTS ON RED IMPORTED FIRE ANT CAPTURES IN THE LOST PINES ECOREGION OF TEXAS.....	158
Abstract.....	159
Introduction.....	160
Study Site	161
Methods.....	162
Statistical analyses	163
Results.....	165
Discussion	167
Acknowledgements.....	169
References.....	169
VII. SUMMARY AND CONCLUSIONS.....	173
Literature Cited	178

LIST OF TABLES

Table	Page
2.1. Number of pre-burn (trapped 17 July 2010 to 24 July 2010) and post-burn (trapped 5 September 2010 to 12 September 2010) captures in three treatments: control, wildfire, and prescribed fire, for juvenile amphibians on the Griffith League Ranch (GLR), Bastrop County, Texas, USA	25
2.2. Results from a mixed effects model analysis with treatment (i.e., control, wildfire, or prescribed fire) and status (i.e., pre-burn or post-burn) as fixed effects and array as a random effect, used to determine if number of captures differed among treatments and sampling periods for three amphibian genera on the Griffith League Ranch (GLR), Bastrop County, Texas, USA	26
2.3. Results from a generalized least squares analysis with treatment (i.e., control, wildfire, or prescribed fire) and status (i.e., pre-burn or post-burn) as factors, used to determine if body condition differed between treatments and sampling periods for 3 amphibian genera on the Griffith League Ranch (GLR), Bastrop County, Texas, USA	28
3.1. Number of unique pre-burn (trapped 16 July 2011 to 23 July 2011), post-burn 1 st (trapped 25 September 2011 to 2 October 2011), and post-burn 2 nd (trapped 19 February 2012 to 30 April 2012) herpetofaunal captures on the Griffith League Ranch (GLR), Bastrop County, Texas, USA, using 24 drift fence arrays	55
3.2. Model selection results from a capture-recapture analysis used to determine if apparent survival (<i>S</i>) and detection probability (<i>P</i>) varied by sex and between non-burned and burned habitat for adult Hurter's Spadefoot Toads (<i>Scaphiopus hurterii</i>) on Griffith League Ranch (GLR), Bastrop County, Texas, USA, during spring 2012.....	60

4.1. Results of generalized least squares analyses used to assess if a low severity winter prescribed burn, a moderate severity summer prescribed burn, and a high severity summer wildfire impacted total captures of ground-dwelling arthropods on the Griffith League Ranch (GLR), Bastrop County, Texas, USA, with sampling season and cumulative precipitation during the sampling period included as covariates	112
6.1. Results from pond-specific tests of a burn status (pre-wildfire/post-wildfire) effect on red imported fire ant (RIFA; <i>Solenopsis invicta</i>) captures-per-unit-effort (CPUE) around pond edges in the Lost Pines ecoregion, Bastrop County, Texas, USA	166

LIST OF FIGURES

Figure	Page
1.1. Number of ha burned through wildfires in the U.S. between 1983 and 2011 (National Interagency Fire Center 2013). Based on a least-squares linear regression, area burned increased by 87,901 ha annually over this time period	3
2.1. Genera included in our investigation on short-term impacts of fire on juvenile amphibians on the Griffith League Ranch (GLR), Bastrop County, Texas, USA in summer 2010: <i>Bufo</i> (A), <i>Scaphiopus</i> (B), and <i>Gastrophryne</i> (C)	16
2.2. Aerial image of the Griffith League Ranch (GLR), Bastrop County, Texas, USA. Overlain on the image are the boundaries of the prescribed burn (7 August 2010) and wildfires (21 August 2010), and the arrays used to determine if number of amphibians captured, amphibian body condition, and number of predatory invertebrates captured differed among treatments and sampling periods	21
2.3. Number of pre-burn (trapped 17 July 2010 to 24 July 2010) and post-burn (trapped 5 September 2010 to 12 September 2010) captures at arrays in three treatments: control ($n = 15$), prescribed fire ($n = 4$), and wildfire ($n = 7$), for three amphibian genera: <i>Bufo</i> (A), <i>Scaphiopus</i> (B), and <i>Gastrophryne</i> (C), and predatory invertebrates (D) on the Griffith League Ranch (GLR), Bastrop County, Texas, USA	27
3.1. Aerial image of the Griffith League Ranch (GLR), Bastrop County, Texas, and its location with respect to a 13,406 ha wildfire that occurred in the Lost Pines ecoregion in September 2011, with a breach of the fire break resulting in an additional 125 ha burned in October 2011	47
3.2. Terrestrial habitat around a drift fence array on the Griffith League Ranch (GLR), Bastrop County, Texas, USA, before (A) and after (B) a high severity wildfire on 4 September 2011	54
3.3. Estimated detection probabilities for Six-lined Race Runners (A) and Southern Prairie Lizards (A), before (Pre) and after (Post) high severity wildfire in control (C) and burn (B) habitats on the Griffith League Ranch (GLR), Bastrop County, Texas, USA	59

3.4. Adult male (M) and female (F) Hurter's Spadefoot Toad (<i>Scaphiopus hurterii</i>) apparent survival (S; A) and detection probability (P; B) estimates in non-burned (C) and burned (B) habitat during spring 2012 on Griffith League Ranch (GLR), Bastrop County, Texas, USA.....	61
3.5. Species accumulation curves for calling anurans at ponds located within non-burned (wildfire; $n = 9$) and burned (control; $n = 10$) habitat on the Griffith League Ranch (GLR), Bastrop County, Texas, USA.....	63
4.1. Aerial image of the Griffith League Ranch (GLR), Bastrop County, Texas, USA, and its location with respect to a 13,406 ha wildfire that occurred in the Lost Pines ecoregion in September 2011, with a breach of the fire break resulting in an additional 125 ha burned in October 2011	88
4.2. Results from Principal Response Curve (PRC) analyses used to assess the impacts of low (A), moderate (B), and high (C) severity fires on pond water quality on the Griffith League Ranch (GLR), Bastrop County, Texas, USA.....	103
4.3. Examples of vegetation plots following low (A), moderate (B), and high (C) severity fires (left panels), and ca. 1 year following the fires (right panel), on the Griffith League Ranch (GLR), Bastrop County, Texas, USA.....	105
4.4. Visual descriptive statistics displaying fire severity to substrate on the Griffith League Ranch (GLR), Bastrop County, Texas, USA for low, moderate, and high severity fires	106
4.5. Overstory (A) and pole-sized (B) tree mortality on the Griffith League Ranch (GLR), Bastrop County, Texas, USA following low, moderate, and high severity fires	107
4.6. Results from Redundancy Analyses (RDA) used to assess the impacts of low severity winter fires (A), moderate severity summer fires (B), and high severity summer fires (C) on forest structure characteristics (left biplot) and species composition of understory vegetation (right biplot) on the Griffith League Ranch (GLR), Bastrop County, Texas, USA.....	110
4.7. Results from Redundancy Analyses (RDA) used to assess the impacts of a low severity winter fire (A), moderate severity summer fires (B), and high severity summer fires (C) on ground-dwelling arthropods in the Lost Pines ecoregion of Texas, USA	113
5.1. Red imported fire ant (RIFA; <i>Solenopsis invicta</i>) captures at pond edges in the Lost Pines ecoregion of Texas	146

5.2. Annual activity pattern of red imported fire ants (RIFA) in the Lost Pines ecoregion of Texas based on captures-per-unit-effort (CPUE)	147
5.3. Relationship between mean percent canopy cover ($\pm$ SE) and mean red imported fire ant (RIFA) captures ($\pm$ SE) at 10 ponds in the Lost Pines ecoregion of Texas	148
6.1. Ordinary least squares (OLS) regression assessing the relationship between differences in mean % canopy cover between the pre-wildfire (September 2009 to October 2010) and post-wildfire (September 2011 to August 2012) sampling periods, and red imported fire ant (RIFA; <i>Solenopsis invicta</i>) captures-per-unit-effort (CPUE) between May and early August (i.e., the peak RIFA activity months common to both the pre-wildfire and post-wildfire sampling periods) around pond edges in the Lost Pines ecoregion, Bastrop County, Texas, USA	167

ABSTRACT

IMPACTS OF LOW, MODERATE, AND HIGH SEVERITY FIRE ON HERPETOFAUNA AND THEIR HABITAT IN A SOUTHERN USA MIXED PINE/HARDWOOD FOREST

by

Donald James Brown, B.S., M.S.

Texas State University-San Marcos

August 2013

SUPERVISING PROFESSOR: MICHAEL R. J. FORSTNER

The primary goals of this dissertation were: 1) to increase our knowledge of fire impacts on amphibians and reptiles, collectively referred to as herpetofauna, the least studied major terrestrial vertebrate groups in relation to fire research; 2) to improve our understanding of fire severity as a factor influencing the response of ecosystem components to fire; and 3) to increase our understanding of the temporal and spatial

ecology, and impacts of fire, on the invasive red imported fire ant (RIFA; *Solenopsis invicta*), a species which has been implicated in the decline of several herpetofaunal species, and for which almost no information exists concerning fire impacts. I accomplished this through the completion of 6 field-based studies in the Lost Pines ecoregion of Texas. I used low and moderate severity prescribed fire to manipulate the habitat, and incorporated unplanned high severity wildfires into my study designs.

The herpetofaunal investigations indicated that direct mortality from fire was not significant, even for high severity wildfires. Further, survivorship of juvenile amphibians may have increased following a moderate severity summer prescribed fire, and the potential increase in survivorship could have been related to a reduction in arthropod predation. The high severity wildfire research indicated the post-wildfire landscape provided suitable habitat for herpetofauna ca. 6 months after the wildfire, which is directly opposed to the general assumption of wildlife managers in the area that the fire killed the majority of herpetofauna and destroyed their habitat. In addition, movement-rates of Hurter's spadefoot toads (*Scaphiopus hurterii*) appeared to increase following the wildfire, and to my knowledge this was the first study that investigated the impacts of fire on amphibian movement-rates.

The study investigating the influence of fire severity on responses of ecosystem components to fire indicated that fire severity was an important factor, and the influence was related to magnitude, but not direction, of effects. For some components (e.g., pond nutrient levels) the magnitude effect was clear, whereas for others (e.g., species composition of understory vegetation) it was dichotomous in that no effect was apparent

for low severity fire and a strong effect was apparent for high severity fire. An additional important finding was mortality of loblolly pine (*Pinus taeda*) trees was nearly 100% in the high severity wildfire zone, and subsequently loblolly pine tree recruitment was low. Thus, restoration of the Lost Pines will require significant active management through reseedling of loblolly pines, with the alternative being a shift to hardwood-dominated forest patches in the high severity wildfire zone.

The RIFA investigations indicated that peak annual RIFA activity coincides with the period when juvenile Houston toads are found at high densities around breeding ponds, and thus are particularly vulnerable to population-level impacts of predation. Further, there was a strong inverse relationship between RIFA captures and overstory canopy cover, and high severity fire appeared to positively affect RIFA through reduction in canopy cover. This is a concerning result given the substantial tree mortality across the high severity wildfire zone, and thus the potential for increases in RIFA abundance and distribution in the Lost Pines ecoregion.

CHAPTER I

INTRODUCTION

Format follows the guidelines for the Journal of Wildlife Management

I. Introduction

Historically fire was an important natural disturbance shaping the structure and composition of pine-dominated forests in the southern U.S. (Gilliam and Platt 1999, Knebel and Wentworth 2007, Hanberry et al. 2012). Longstanding fire suppression policies have resulted in structural and compositional changes, notably accumulation of heavy fuel loads and reduction in vegetation species diversity (Agee 1996, Shang et al. 2007, Crotteau et al. 2013). Fire suppression, coupled with climatic trends towards warmer and drier conditions, have resulted in an increase in high severity wildfires (Figure 1.1), particularly in the southern and western U.S. (Davis 2001, Miller et al. 2009), with this trend projected to continue into the next century (Moritz et al. 2012). The term fire severity refers to amount of organic matter lost through burning, as opposed to the related term fire intensity, which refers to the heat energy of fire (Keeley 2009).




Figure 1.1. Number of ha burned through wildfires in the U.S. between 1983 and 2011 (National Interagency Fire Center 2013). Based on a least-squares linear regression, area burned increased by 87,901 ha annually over this time period.

Prescribed burning is currently one of the primary tools used for restoration of fire suppressed forests (Allen et al. 2002, Glasgow and Matlack 2007, Mitchell et al. 2009). Although site-specific goals of burning vary, overarching goals typically include reduction of heavy fuel loads (Fernandes and Botelho 2003, Shang et al. 2007, Webster and Halpern 2010), and mimicking ecosystem impacts of historically natural wildfires within a controlled setting (Moore et al. 1999, Vose 2000, Allen et al. 2002). Much research has been devoted to effects of burn season and frequency, particularly on vegetation (e.g., Cain et al. 1998, Sparks et al. 1998, Taylor 2000, Webster and Halpern

2010). Much less is known about effects of burn severity, given that prescribed burns are typically low to moderate severity fires. In contrast, high severity wildfires are unplanned, and thus wildfire research is by necessity a response to non-designed treatments, and opportunities for research are more limited. However, the need to improve our understanding of the influence of fire severity on ecosystem responses to fire is becoming an increasingly pressing need as the annual area burned in high severity fires in the U.S. continues to rise.

There are several ways to approach ecosystem-level questions: 1) natural history surveys (i.e., no direct ecosystem manipulation, but potentially incorporating disturbance history); 2) energy budget or mass balance studies; 3) experimental manipulation of ecosystem components; 4) comparisons of previous work across ecosystems, preferably through a meta-analysis approach; and 5) process-based simulation modeling, which benefits greatly from use of data derived from other approaches (Likens 1992). For this dissertation research I used an experimental manipulation approach (i.e., prescribed burning), coupled with natural disturbances (i.e., wildfires), to assess the influence of fire severity on ecosystem components in a southern USA mixed pine/hardwood forest ecosystem. All of the studies utilized a before-after/control-impact (BACI) framework, which is the optimal scenario for experimental field research because it considers background spatial and temporal variability (Stewart-Oaten and Bence 2001).

The studies in this dissertation were conducted on the 1,948 ha Griffith League Ranch (GLR), located in the 34,400 ha Lost Pines ecoregion in Bastrop County, Texas. The Lost Pines was extensively logged in the 1800s and early 1900s (Moore 1977). Since

the early to mid-1900s broad-scale fire suppression has been implemented throughout the ecoregion, resulting in the accumulation of heavy fuel loads, similar to forested regions throughout much of the Southern U.S (Cooper 1960, Hartnett and Krofta 1989, Covington and Moore 1994, Gilliam and Platt 1999). Management and conservation initiatives in the Lost Pines are largely focused on promoting healthy non-game wildlife populations, and low severity prescribed fire is currently a heavily promoted habitat management tool for herpetofaunal species of conservation concern (particularly the federally endangered Houston toad [*Bufo (Anaxyrus) houstonensis*]; KES Consulting et al. 2007), despite the lack of quantitative information on direct or indirect impacts to herpetofaunal populations and their habitat in the Lost Pines. Thus, a major component of this research addressed fire impacts on herpetofauna.

This dissertation includes 6 studies that assess fire impacts on components of a southern mixed pine/hardwood forest ecosystem, with emphasis on fire severity as a factor affecting outcomes. Chapters II and III focus on immediate and short-term impacts of fire on herpetofauna. The focus of Chapter II is on the impacts of moderate severity fires on juvenile amphibian survivorship and body condition, whereas Chapter III assesses the impacts of a high severity fire on abundance, movement, and diversity of the herpetofaunal community. Chapter IV compares the impacts of low, moderate, and high severity fire impacts to several aquatic and terrestrial ecosystem components, including water quality, species composition of aquatic arthropods, forest structure characteristics, species composition of understory vegetation, and species composition of ground-dwelling arthropods. Chapters V and VI investigate the impacts of low and moderate

severity (i.e., Chapter V), and high severity (i.e., Chapter VI), fire on red imported fire ant (*Solenopsis invicta*) captures around forest ponds. Fire impacts on this invasive species are of interest given it is a known predator of juvenile Houston toads (Freed and Neitman 1988), and there is evidence that they also prey upon adult Houston toads (M. C. Jones, Texas State University, personal communication).

Taken together, these studies provide valuable information concerning the influence of fire severity on responses of forested ecosystems to fire for a diverse array of ecosystem components. In addition, they provide useful information for land and wildlife managers engaged in conservation and recovery initiatives in the Lost Pines ecoregion. Finally, these studies provide the necessary baseline data to investigate long-term impacts of fire on the Lost Pines ecoregion through both empirical data collection and process-based modeling approaches.

LITERATURE CITED

- Agee, J. K. 1996. Achieving conservation biology objectives with fire in the Pacific Northwest. *Weed Technology* 10:417–421.
- Allen, C. D., M. Savage, D. A. Falk, K. F. Suckling, T. W. Swetnam, T. Schulke, P. B. Stacey, P. Morgan, M. Hoffman, and J. T. Klingel. 2002. Ecological restoration of southwestern ponderosa pine ecosystems: a broad perspective. *Ecological Applications* 12:1418–1433.

- Cain, M. D., T. B. Wigley, and D. J. Reed. 1998. Prescribed fire effects on structure in uneven-aged stands of loblolly and shortleaf pines. *Wildlife Society Bulletin* 26:209–218.
- Cooper, C. F. 1960. Changes in vegetation, structure, and growth of southwestern pine forests since white settlement. *Ecological Monographs* 30:130–164.
- Covington, W. W., and M. M. Moore. 1994. Southwestern ponderosa forest structure changes since Euro-American settlement. *Journal of Forestry* 92:39–47.
- Crotteau, J. S., J. M. Varner, and M. W. Ritchie. 2013. Post-fire regeneration across a fire severity gradient in the southern Cascades. *Forest Ecology and Management* 287:103–112.
- Davis, C. 2001. The west in flames: the intergovernmental politics of wildfire suppression and prevention. *Publius* 31:97–110.
- Fernandes, P. M., and H. S. Botelho. 2003. A review of prescribed burning effectiveness in fire hazard reduction. *International Journal of Wildland Fire* 12:117–128.
- Freed, P. S., and K. Neitman. 1988. Notes on predation on the endangered Houston toad, *Bufo houstonensis*. *Texas Journal of Science* 40:454–456.
- Gilliam, F. S., and W. J. Platt. 1999. Effects of long-term fire exclusion on tree species composition and stand structure in an old-growth *Pinus palustris* (longleaf pine) forest. *Plant Ecology* 140:15–26.
- Glasgow, L. S., and G. R. Matlack. 2007. Prescribed burning and understory composition in a temperate deciduous forest, Ohio, USA. *Forest Ecology and Management* 238:54–64.

- Hanberry, B. B., D. C. Dey, and H. S. He. 2012. Regime shifts and weakened environmental gradients in open oak and pine ecosystems. *PLoS ONE* 7:e41337.
- Hartnett, D. C., and D. M. Krofta. 1989. Fifty-five years of post-fire succession in a southern mixed hardwood forest. *Bulletin of the Torrey Botanical Club* 116:107–113.
- Keeley, J. E. 2009. Fire intensity, fire severity and burn severity: a brief review and suggested usage. *International Journal of Wildland Fire* 18:116–126.
- KES Consulting, Loomis Austin, and M. R. J. Forstner. 2007. Lost pines habitat conservation plan for Bastrop County, Texas. <<http://www.co.bastrop.tx.us/site/content/lostpineshabitat>>. Accessed 1 Dec 2010.
- Knebel, L., and T. R. Wentworth. 2007. Influence of fire and southern pine beetle on pine-dominated forests in the Linville Gorge Wilderness, North Carolina. *Castanea* 72:214–225.
- Likens, G. E. 1992. The ecosystem approach: its use and abuse. Ecology Institute, Oldendorf/Luhe, Germany.
- Miller, J. D., H. D. Safford, M. Crimmins, and A. E. Thode. 2009. Quantitative evidence for increasing forest fire severity in the Sierra Nevada and Southern Cascade Mountains, California and Nevada, USA. *Ecosystems* 12:16–32.
- Mitchell, S. R., M. E. Harmon, and K. E. B. O'Connell. 2009. Forest fuel reduction alters fire severity and long-term carbon storage in three Pacific Northwest ecosystems. *Ecological Applications* 19:643–655.

- Moore, B. 1977. Bastrop County 1691–1900. Revised edition. Nortex Press, Wichita Falls, Texas, USA.
- Moore, M. M., W. W. Covington, and P. Z. Fule. 1999. Reference conditions and ecological restoration: a southwestern ponderosa pine perspective. *Ecological Applications* 9:1266–1277.
- Moritz, M. A., M.-A. Parisien, E. Batllori, M. A. Krawchuk, J. Van Dorn, D. J. Ganz, and K. Hayhoe. 2012. Climate change and disruptions to global fire activity. *Ecosphere* 3:Article 49.
- National Interagency Fire Center. 2013. Total wildland fire acres (1960-2009). <http://www.nifc.gov/fireInfo/fireInfo_stats_totalFires.html>. Accessed 3 Feb 2013.
- Shang, Z., H. S. He, D. E. Lytle, S. R. Shifley, and T. R. Crow. 2007. Modeling the long-term effects of fire suppression on central hardwood forests in Missouri Ozarks, using LANDIS. *Forest Ecology and Management* 242:776–790.
- Sparks, J. C., R. E. Masters, D. M. Engle, M. W. Palmer, and G. A. Bukenhofer. 1998. Effects of late growing-season and late dormant-season prescribed fire on herbaceous vegetation in restored pine-grassland communities. *Journal of Vegetation Science* 9:133–142.
- Stewart-Oaten, A., and J. R. Bence. 2001. Temporal and spatial variation in environmental impact assessment. *Ecological Monographs* 71:305–339.

- Taylor, A. H. 2000. Fire regimes and forest changes in mid and upper Montane forests of the Southern Cascades, Lassen Volcanic National Park, California, U.S.A. *Journal of Biogeography* 27:87–104.
- Vose, J. M. 2000. Perspectives on using prescribed fire to achieve desired ecosystem conditions. Pages 12–17 *in* W. K. Moser, and C. E. Moser, editors. Proceedings, 21st, Tall Timbers fire ecology conference: fire and forest ecology: innovative silviculture and vegetation management. Tall Timbers Research Station, Tallahassee, Florida, USA.
- Webster, K. M., and C. B. Halpern. 2010. Long-term vegetation responses to reintroduction and repeated use of fire in mixed-conifer forests of the Sierra Nevada. *Ecosphere* 1:Article 9.

CHAPTER II

POTENTIAL POSITIVE EFFECTS OF FIRE ON JUVENILE AMPHIBIANS IN A SOUTHERN USA MIXED PINE/HARDWOOD FOREST

Format follows the guidelines for the Journal of Fish and Wildlife Management

II. Potential Positive Effects of Fire on Juvenile Amphibians in a Southern USA

Mixed Pine/Hardwood Forest

Abstract

Prescribed fire is a common tool used to conserve and manage the integrity of forest ecosystems. We investigated short-term juvenile amphibian capture and body condition changes subsequent to fire (i.e., one prescribed burn and two wildfires) in a southern USA pine forest. We surveyed amphibians and predatory invertebrates before and after fires during summer 2010. We tested for treatment (i.e., control, wildfire, or prescribed burn) and status (i.e., pre-burn or post-burn) differences in: (1) genus-level captures of amphibians, (2) amphibian health (inferred through a body condition index), and (3) predatory invertebrate captures. *Bufo* and *Scaphiopus* captures increased in the prescribed burn treatment; whereas, no differences in *Gastrophryne* captures were observed. We did not detect a burn status effect on amphibian body condition. Predatory invertebrate captures were higher post-burn in control and wildfire treatments. Neither a moderate severity prescribed burn nor moderate severity wildfires negatively impacted short-term juvenile amphibian captures. Further, we speculate that *Bufo* and *Scaphiopus* survivorship may have been higher after the prescribed burn.

Keywords: amphibians; fire; forest; Houston toad; predators.

Introduction

Prescribed fire has increased as a primary tool for conservation and recovery of fire-maintained ecosystems (Agee 1996; Sparks et al. 1998; Kloor 2000; Pyne 2010). Prescribed fire is often used to decrease heavy fuel loads in historically fire-suppressed forests, in the prevention of catastrophic wildfires, and to stimulate the recovery of understory vegetation (Sweeney and Biswell 1961; Lovaas 1976; Cain et al. 1998; Ryu et al. 2006). Restoring understory vegetation in fire-suppressed forests may promote greater animal diversity and abundance (Moseley et al. 2003; Smucker et al. 2005; Benson et al. 2007). Fire also stimulates aquatic productivity through increased nutrient loads in streams and ponds (Gresswell 1999; Scrimgeour et al. 2001) and has been linked to mitigation of spruce beetle (*Dendroctonus rufipennis*) outbreaks (Bebi et al. 2003).

Scientific investigations of fire effects on wildlife have progressed slowly from game to non-game species over the last century. Amphibians are possibly the least-studied major vertebrate group in relation to fire research (Russell et al. 1999; Pilliod et al. 2003). The importance of amphibians to terrestrial and aquatic food webs (Burton and Likens 1975; Semlitsch et al. 1996; Walls and Williams 2001; Davic and Welsh 2004) accentuates their potential use as vertebrate bioindicators (Welsh and Ollivier 1998; Kerby et al. 2010) and inherent ecological value. There is a need to achieve a better understanding of changes in amphibian populations due to fire.

Historically, amphibians were assumed vulnerable to direct mortality from fire, largely due to limited mobility and direct mortality observations (Babbitt and Babbitt 1951). However, the majority of published research found no evidence that fire significantly affects short-term abundance of amphibian species (Ford et al. 1999;

Ruthven et al. 2008). Grafe et al. (2002) suggested surface-aestivating anurans (i.e., frogs and toads) respond to auditory cues produced as a result of approaching fire by seeking burn-resistant refugia. Longer-term responses to fire-induced habitat changes are species and context specific (Moseley et al. 2003; Cummer and Painter 2007; Cano 2009; Rochester et al. 2010).

Prescribed burning during terrestrial post-metamorphic growth phases could potentially affect juvenile amphibian survivorship and health by reducing the quantity and quality of refugia by lessening litter, duff, and coarse woody debris (Tinker and Knight 2000; Matthews et al. 2010). Fire could impact juveniles both negatively and positively through invertebrate mortality. Moretti et al. (2006) and Vasconcelos et al. (2009) reported negative impacts on arthropod communities following fire, while Taber et al. (2008) and Greenberg et al. (2010) found little or no differences. Fire could negatively affect juvenile amphibians through a decline in their food base, but positively affect them through predator mortality (Toledo 2005). Predation probably plays a major role in survivorship of most amphibians at all life stages (Wells 2007). In addition to removal of predatory invertebrates, fire may also reduce predation by mesocarnivores. Jones et al. (2004) found raccoons (*Procyon lotor*) had higher preference for unburned compared to burned longleaf pine (*Pinus palustris*) stands, and Sunquist (1967) found raccoons reduced foraging activity in a Minnesota savannah following a prescribed burn.

The purposes of our study were to determine whether fire affected juvenile amphibian captures per unit effort (CPUE), and thus potentially juvenile amphibian survivorship, whether fire affected health (i.e., body condition) of individuals, and whether fire affected CPUE of potentially predatory invertebrates. To our knowledge this

is the first quantitative study specifically addressing the short-term effects of fire explicitly on terrestrial juvenile amphibians.

Study Site

We conducted this study on the 1,948-ha Griffith League Ranch (GLR) in Bastrop County, Texas, USA. The GLR is located in the Lost Pines ecoregion, a 34,400-ha remnant patch of pine-dominated forest isolated from the East Texas Piney Woods ecoregion during the Pleistocene (Bryant 1977; Al-Rabab'ah and Williams 2004). The GLR is primarily a forested ranch with an overstory dominated by loblolly pine (*Pinus taeda*), post oak (*Quercus stellata*), blackjack oak (*Quercus marilandica*), and eastern red cedar (*Juniperus virginiana*), and an understory dominated by yaupon holly (*Ilex vomitoria*), American beautyberry (*Callicarpa americana*), and farkleberry (*Vaccinium arboreum*). The GLR contains three permanent ponds (i.e., ponds have not dried in at least 12 years), 10 semi-permanent ponds (i.e., ponds typically dry several times per decade), and dozens of ephemeral pools that hold water for days to months annually depending on rainfall. Prior to 2009, the study area had been fire suppressed for at least 60 years.

We documented 12 amphibian species on the GLR; 5 were included in this investigation (Figure 2.1): the coastal plain toad (*Bufo* [*Incilius*] *nebulifer*), the endangered Houston toad (*Bufo* [*Anaxyrus*] *houstonensis*), Hurter's spadefoot toad (*Scaphiopus hurterii*), the eastern narrowmouth toad (*Gastrophryne carolinensis*), and the Great Plains narrowmouth toad (*Gastrophryne olivacea*). These species typically breed in spring and early summer in this region, beginning in February for the Houston

toad, and late March or early April for the remaining species (Forstner and Swannack 2004; Saenz et al. 2006; Brown et al. *In press*).




Figure 2.1. Genera included in our investigation on short-term impacts of fire on juvenile amphibians on the Griffith League Ranch (GLR), Bastrop County, Texas, USA in summer 2010: *Bufo* (A), *Scaphiopus* (B), and *Gastrophryne* (C). Individuals shown are the Houston toad *B. houstonensis*, Hurter's spadefoot toad *S. hurterii*, and the Great Plains narrowmouth toad *G. olivacea*, respectively.

Metamorphosis usually occurs 15 to 50 days after eggs hatch, with a shorter time to metamorphosis for *Scaphiopus* and longer time to metamorphosis for *Bufo* (Wright and Wright 1949; Hillis et al. 1984). Pond-breeding anurans typically aggregate near pond

edges for several weeks to months after metamorphosis and then disperse into the surrounding terrestrial landscape (Arnold and Wassersug 1978; Greuter 2004). Studies have shown directional preferences in initial movement away from ponds (deMaynadier and Hunter 1999; Vasconcelos and Calhoun 2004), after which point movement appears to be essentially random outside of the microhabitat scale (Semlitsch 2008). Juvenile movements and foraging activities occur throughout summer months (June to September), with daily activity patterns likely heavily influenced by weather conditions (Roe and Grayson 2008; Child et al. 2009).

Methods

The primary goal of the GLR prescribed fire management policy is to reduce the depth of litter and duff layers accumulated over the past one-half century. This reduces the probability of a catastrophic wildfire and mimics the type of prescribed burns currently conducted throughout the Lost Pines ecoregion. We conducted a moderate severity prescribed burn on a 262-ha burn unit on 7 August 2010. Two moderate severity wildfires occurred on the GLR on 21 August 2010, burning 153 ha and 36 ha, respectively. Burn breaks were installed during the wildfires due to the intensity of the burns and the risk of the fires spreading beyond the boundaries of the GLR.

We assessed initial burn effects on habitat using vegetation plots (20 m by 50 m) established in 2008 by random placement within forested habitat. We surveyed vegetation plots between 10 and 29 days following the burns using National Park Service (2003) fire monitoring guidelines. Four vegetation plots were burned during the prescribed burn and one vegetation plot was burned during the wildfire. We assessed

burn severity to substrate and vegetation within each plot using four 15 m transect lines, each consisting of four points spaced 5 m apart. We assigned points a burn severity ranking from 1 (heavily burned) to 4 (scorched) using a qualitative visual assessment (National Park Service 2003). In addition, we estimated char height and recorded status (i.e., alive or dead) for all overstory trees (i.e., diameter at breast height > 15 cm) within vegetation plots.

Sampling design

We used 18 Y-shaped and eight linear arrays for trapping amphibians and invertebrates. Y-shaped arrays consisted of three 15 m arms of flashing with a 19 L center bucket and a 19 L bucket at each arm terminus. Seven linear arrays consisted of a 15 m arm with a 19 L bucket at each end, and a double-throated funnel trap in the center of the array on each side of the flashing. One linear array consisted of a 121 m arm with a 19 L bucket at each end and a 19 L bucket near the center of the trap. We equipped pitfall traps with flotation devices to mitigate mortality during bucket flooding, and both pitfall and funnel traps with wet sponges to provide a moist environment for amphibians. We also equipped pitfall traps with predator exclusion devices (Ferguson and Forstner 2006).

The sampling design consisted of six sets of arrays with each set containing three Y-shaped arrays spread along drainages and between ponds, and one linear array adjacent to, and parallel with, a pond. The two remaining linear arrays were used to monitor amphibian movement between forest and grassland habitat, and activity at an additional pond, respectively. We did not create this sampling design explicitly for this experiment, but rather as part of long-term investigations on amphibian activity, movement, treatment

response (e.g., prescribed fire), and habitat use (Swannack et al. 2009). To determine if the sampling design was suitable for detecting juvenile amphibians (arrays were located up to 618 m from breeding ponds) we regressed numbers of juvenile captures of three genera (*Bufo*, *Gastrophryne*, *Scaphiopus*) against array distance from breeding pond. These analyses showed that captures for all 3 genera spanned the distribution of array distance from source ponds; captures were positively associated with distances from breeding pond, and slopes were similar between sampling periods. Therefore, we included all 26 arrays in this study.

We trapped amphibians and invertebrates for seven days between 17 July 2010 and 24 July 2010 (i.e., 21 to 27 days prior to the prescribed burn), and seven days between 5 September 2010 and 12 September 2010 (i.e., 29 to 35 days after the prescribed burn, and 15 to 21 days after the wildfires). The September sampling period was unusually wet due to a heavy rain event caused by Tropical Storm Hermine. We checked traps and processed amphibians daily. We toe-clipped amphibians using an individual numbering system (Martof 1953), recorded snout-vent length (SVL) and head width to the nearest 0.1 mm using digital calipers (Control Company, Friendswood, Texas), and weight to the nearest 0.1 g using spring scales (Pesola, Baar, Switzerland). We collected invertebrates from all pitfall traps on the last day of each sampling period (i.e., we allowed pitfall traps to accumulate invertebrates for 7 days) and euthanized them by freezing. Handling permits were provided by Texas Parks and Wildlife Department (SPR-0102-191) and U.S. Fish and Wildlife Service (TE 039544-0). Trapping and handling methods were approved by the Texas State University-San Marcos Institutional Animal Care and Use Committee (Protocol No. 0810_0208_11).

A 1-m fire break was cleared around arrays in the prescribed burn unit, and we ensured that the burn carried to those breaks and spread throughout the adjacent habitat. Thus, the prescribed fire treatment included all four arrays within the treatment area. The wildfires occurred adjacent to two of the array sets. The distance from wildfire perimeters to these arrays ranged from 0 m to 280 m, with a mean distance of 94 m. Terrestrial juvenile anuran movement patterns are not well-studied (Wells 2007), and thus it is difficult to define the appropriate cutoff distance for wildfire influence on array captures during our study period. Smith and Green (2006) reported that most recaptured juvenile and adult Fowler's toads (*Bufo* [*Anaxyrus*] *fowleri*; $n = 1326$) were within 100 m of the location of initial capture over the course of two month sampling periods. Semlitsch (2008) suggested that long distance dispersal of most amphibian species is probably broken up into discrete events over multiple years due to low stamina in juveniles and a positive relationship between body size and locomotor capacity.

Because our preliminary regression analyses did not indicate that we were capturing juveniles actively dispersing from breeding ponds in the first sampling period (i.e., slopes were similar between sampling periods), we assumed that most individuals remained in the vicinity of their initial capture for the duration of the study. We considered captures at arrays >100 m from wildfire perimeters unlikely to be significantly affected by the wildfires, and we included these arrays as controls. The wildfire treatment included seven arrays, and the mean distance from wildfire perimeters to wildfire treatment arrays was 52 m (Figure 2.2). The prescribed fire treatment contained four arrays, with distance to unburned habitat ranging from 96 m to 209 m. The control

treatment included 15 arrays, with distance to burned habitat ranging from 190 m to 1,295 m.




Figure 2.2. Aerial image of the Griffith League Ranch (GLR), Bastrop County, Texas, USA. Overlain on the image are the boundaries of the prescribed burn (7 August 2010) and wildfires (21 August 2010), and the arrays used to determine if number of amphibians captured, amphibian body condition, and number of predatory invertebrates captured differed among treatments and sampling periods. The prescribed fire unit was 262 ha, and the wildfires burned 153 ha and 36 ha, respectively. The control, wildfire, and prescribed burn treatments contained 15, 7, and 4 arrays, respectively.

Statistical analyses

To increase sample sizes for amphibian capture comparisons we grouped captures by genus using a traditional classification system (Dixon 2000): *Bufo* (*B. nebulifer* and *B. houstonensis*), *Scaphiopus* (*S. hurterii*), and *Gastrophryne* (*G. carolinensis* and *G. olivacea*). We did not analyze the other genera present in our study area, *Acris*, *Ambystoma*, *Hyla*, and *Rana* (*Lithobates*), due to low total captures (i.e., <15 individuals per genus) of these taxa. Based on SVL values reported in Wright and Wright (1949), we captured 55 adults and removed these individuals from the data set.

For the three genera investigated, we used the total numbers of unique juveniles captured at each array within each sampling period as the response variable. We analyzed these data using mixed effects models, which allow for unbalanced designs (Zuur et al. 2009). We designated treatment (i.e., control, wildfire, or prescribed fire) and status (i.e., pre-burn or post-burn) as fixed effects and array as a random effect. We assessed assumptions of normality and homoscedasticity using residual plots. When we found significant differences, we determined which means were different using contrast comparisons (Maindonald and Braun 2003). We performed these analyses with the program R (R Version 2.10.1, www.r-project.org) using the nlme (Pinheiro et al. 2009) and contrast (Kuhn et al. 2010) packages.

We used body condition indices (BCI) to determine if health of individuals was affected by fire (Reading and Clarke 1995). For each genus within each sampling period, we regressed body length (SVL) on weight and used residuals as our response variable (Schulte-Hostedde et al. 2005). Positive and negative residuals indicate above and below average body condition, respectively. We removed 33 individuals from the data set

because SVL or weight was not recorded, as well as recaptures within sampling periods. We linearized data for all three genera using the $\log_{10}$ transformation (Fowler et al. 1998). We analyzed these data using generalized least squares (gls) models, with treatment and status as predictors (Zuur et al. 2009). When we found significant differences, we determined which means were different using contrast comparisons (Maindonald and Braun 2003). We performed these analyses with the program R (R Version 2.10.1, www.r-project.org) using the nlme (Pinheiro et al. 2009) and contrast (Kuhn et al. 2010) packages.

Several larger invertebrates in our study system are likely to exert substantial predation pressure on terrestrial juvenile amphibians, including members of the class Arachnida (orders Scorpiones [i.e., scorpions] and Araneae [i.e., spiders]), and beetles in the family Carabidae (Toledo 2005). Although ants are also known to prey upon juvenile amphibians, we did not include ant captures because they can crawl into and out of pitfall traps, and thus, we could not accurately detect their captures with this sampling design. However, individuals are probably most vulnerable to ant predation when they first leave the water (Freed and Neitman 1988; Toledo 2005). Thus, for the predatory invertebrate analysis we used the total number of scorpions (*Centruroides vittatus*; Taber and Fleenor 2003), ground-dwelling spiders (primarily wolf spiders), and carabid beetles captured at each array within each sampling period as the response variable. We analyzed these data using identical statistical methods as for amphibian captures.

Results

The live fuel moisture during the prescribed burn was ca. 130%, with temperature, relative humidity, and wind speed estimates of 32.4 °C, 56.5%, and 1.6 kph, respectively, taken during the burn. Fire passed through 27 of the 64 substrate burn severity points during the prescribed burn, 13 of which contained vegetation. The mean severity ranking for substrate and vegetation was 3.5 and 3.7, respectively. Of the 164 overstory trees, 73 were charred, with a mean char height of 0.4 m. Four of the 141 overstory trees that were alive prior to the burn were killed. Temperature, relative humidity, and wind speed estimates taken during the wildfire were 37.6 °C, 32.4%, and 3.2 kph, respectively, with wind gusts up to 12.9 kph. Fire passed through all sixteen substrate burn severity points during the wildfires, 10 of which contained vegetation. The mean severity ranking for substrate and vegetation was 2.5 and 2.2, respectively. Of the 37 overstory trees, 33 were charred, with a mean char height of 2.8 m. Ten of the 30 overstory trees that were alive prior to the burn were killed.

We captured 210 *Bufo*, *Scaphiopus*, and *Gastrophryne* pre-burn, and 217 *Bufo*, *Scaphiopus*, and *Gastrophryne* post-burn. Individual species results are shown in Table 2.1. We recaptured two individuals in both sampling periods, and recaptured nine individuals within sampling periods. We found a significant treatment and treatment-status interaction effect for *Bufo*, and a treatment-status interaction effect for *Scaphiopus* (Table 2.2). Contrast comparisons showed that pre-burn and post-burn captures differed in the prescribed fire treatment for *Bufo* ($t_{44} = -3.03$, $P = 0.004$) and *Scaphiopus* ($t_{44} = -3.87$, $P < 0.001$), with no differences detected for the control or wildfire treatments (Figure 2.3).

Table 2.1. Number of pre-burn (trapped 17 July 2010 to 24 July 2010) and post-burn (trapped 5 September 2010 to 12 September 2010) captures in three treatments: control, wildfire, and prescribed fire, for juvenile amphibians on the Griffith League Ranch (GLR), Bastrop County, Texas, USA.

Species	Control ^a		Wildfire ^b		Prescribed fire ^c	
	<u>Pre-burn</u>	<u>Post-burn</u>	<u>Pre-burn</u>	<u>Post-burn</u>	<u>Pre-burn</u>	<u>Post-burn</u>
<i>Bufo houstonensis</i>	3	0	1	0	1	0
<i>Bufo nebulifer</i>	57	51	7	20	3	52
<i>Bufo</i> spp. ^d	8	0	5	0	9	0
<i>Gastrophryne carolinensis</i>	14	4	0	2	5	3
<i>Gastrophryne olivacea</i>	27	17	12	9	4	3
<i>Scaphiopus huerterii</i>	38	31	4	8	2	17

^aControl treatment captures included 15 arrays.

^bWildfire treatment captures included 7 arrays.

^cPrescribed fire treatment captures included 4 arrays.

^dYoung individuals lacked interspecific morphological trait differences.

Table 2.2. Results from a mixed effects model analysis with treatment (i.e., control, wildfire, or prescribed fire) and status (i.e., pre-burn or post-burn) as fixed effects and array as a random effect, used to determine if number of captures differed among treatments and sampling periods for three amphibian genera on the Griffith League Ranch (GLR), Bastrop County, Texas, USA.

Genus	Source of Variation	df	<i>F</i>	<i>P</i>
<i>Bufo</i>	Treatment	2,23	5.03	0.015
	Status	1,23	0.61	0.444
	Treatment*Status	2,23	4.97	0.016
<i>Scaphiopus</i>	Treatment	2,23	0.85	0.439
	Status	1,23	1.48	0.236
	Treatment*Status	2,23	7.51	0.003
<i>Gastrophryne</i>	Treatment	2,23	0.12	0.889
	Status	1,23	2.70	0.114
	Treatment*Status	2,23	0.42	0.662




Figure 2.3. Number of pre-burn (trapped 17 July 2010 to 24 July 2010) and post-burn (trapped 5 September 2010 to 12 September 2010) captures at arrays in three treatments: control ($n = 15$), prescribed fire ($n = 4$), and wildfire ($n = 7$), for three amphibian genera: *Bufo* (A), *Scaphiopus* (B), and *Gastrophryne* (C), and predatory invertebrates (D) on the Griffith League Ranch (GLR), Bastrop County, Texas, USA. Boxes enclose the range, diamonds show the mean, and horizontal bars delineate the median number of captures at arrays within each treatment and sampling period.

For body condition comparisons we found a treatment effect for *Bufo*, and no significant differences for *Scaphiopus* or *Gastrophryne* (Table 2.3). For *Bufo*, the contrast comparisons showed body condition in the prescribed burn treatment differed from both the control ($t_{212} = -2.98$, $P = 0.003$) and wildfire ($t_{212} = -3.59$, $P < 0.001$) treatments. In both pre-burn and post-burn samples, *Bufo* in the control and wildfire treatments had a greater than average mean BCI score, and *Bufo* in the prescribed burn treatment had a lower than average mean BCI score.

Table 2.3. Results from a generalized least squares analysis with treatment (i.e., control, wildfire, or prescribed fire) and status (i.e., pre-burn or post-burn) as factors, used to determine if body condition differed between treatments and sampling periods for 3 amphibian genera on the Griffith League Ranch (GLR), Bastrop County, Texas, USA.

Genus	Source of Variation	df	<i>F</i>	<i>P</i>
<i>Bufo</i>	Treatment	2,212	7.71	<0.001
	Status	1,212	0.40	0.528
	Treatment*Status	2,212	1.64	0.196
<i>Scaphiopus</i>	Treatment	2,86	0.01	0.994
	Status	1,86	0.00	0.996
	Treatment*Status	2,86	0.03	0.967
<i>Gastrophryne</i>	Treatment	2,80	0.48	0.621
	Status	1,80	0.00	0.998
	Treatment*Status	2,80	0.66	0.517

We captured 123 predatory invertebrates pre-burn, and 182 predatory invertebrates post-burn. We found a status effect for predatory invertebrate captures ($F_{1,23} = 6.91$, $P = 0.015$). Treatment ($F_{2,23} = 0.18$, $P = 0.836$) and treatment-status interaction ($F_{2,23} = 0.86$, $P = 0.438$) effects were not significant.

Discussion

We found that neither the prescribed fire nor the wildfires negatively impacted juvenile captures or body condition for the three amphibian genera investigated. Indeed, *Bufo* and *Scaphiopus* captures were higher post-burn in the prescribed fire treatment. The fact that we captured substantially more *Bufo* and *Scaphiopus* post-burn in the prescribed

fire unit warrants the following potential explanations: 1) the statistical increase in individuals was purely a function of weather-influenced activity; 2) new individuals entered the prescribed fire unit after the first sampling period, either through tadpole metamorphosis or dispersal into the unit; 3) capture rates were higher in the prescribed fire unit after the burn (presumably as a function of increased mobility and hence pitfall encounters by these species); or 4) survivorship was higher in the prescribed fire unit.

Weather is often strongly tied to amphibian population trends and activity patterns (Hillis et al. 1984; Pechmann and Wilbur 1994; Wells 2007) and undoubtedly influenced the number of captures within each sampling period. Here Tropical Storm Hermine resulted in substantial rains (8.1 cm) across the study area during the September sampling period. However, if weather were the only explanatory factor, we would expect to see the fire treatment results mirrored in the control treatment results. This appeared to be the case for *Gastrophryne* but not for *Bufo* or *Scaphiopus* with respect to the prescribed fire treatment. Thus, as discussed below, it appears additional factors influenced numbers of post-burn captures for at least the two latter species.

Another potential explanation is that the numbers of individuals increased in the prescribed fire unit after the first sampling period either through recruitment (i.e., tadpole metamorphosis) or dispersal into the unit. If recruitment were a major factor, we would expect SVL to be similar in the second sampling period (i.e., a mix of new small juveniles plus larger juveniles present during the first sampling period). Based on mean (± 1 SD) SVL between sampling periods in the prescribed fire unit (*Bufo* pre-burn: 20.26 mm $\pm$ 8.16, *Bufo* post-burn: 38.11 mm $\pm$ 6.40; *Scaphiopus* pre-burn: 20.30 mm $\pm$ 1.41,

Scaphiopus post-burn: $29.91 \text{ mm} \pm 4.67$), it is unlikely that recruitment can explain the results.

Although we cannot rule out the possible influence of dispersal into the prescribed fire unit, the nearest potential source pond outside of the sampling area was 900 m away. Further, only 1 of the 11 recaptures was found at a trap different from where it was first captured. Two individuals were captured in both sampling periods, and both of them were recaptured at the same trap. Finally, the mean BCI score in the prescribed fire treatment was significantly lower than control and wildfire treatments, but similar between sampling periods (*Bufo* pre-burn: -0.03 ± 0.07 , *Bufo* post-burn: -0.01 ± 0.03 ; *Scaphiopus* pre-burn: 0.01 ± 0.06 , *Scaphiopus* post-burn: 0.00 ± 0.04). These data, coupled with the regression analyses indicate that long distance dispersal was not prevalent during our study period.

The reduction in vegetative cover and litter depth in the prescribed fire treatment could have increased juvenile amphibian movement rates, and thus capture rates. Chelgren et al. (2011) found that detection probability for terrestrial salamanders was higher in burned than unburned plots. However, their sampling design consisted of randomized searching under substrates (i.e., active sampling) and thus did not address movement rates. The prescribed burn left much of the live vegetation unconsumed, and although the upper litter layer was consumed, there were no obvious changes that would have greatly increased mobility potential.

We hypothesize the most likely explanation for increased *Bufo* and *Scaphiopus* captures following the prescribed burn was increased survivorship coupled with increased activity levels as a consequence of Tropical Storm Hermine. Further, based on the BCI

results, if this hypothesis is correct, it was more likely due to reduced predation pressure following the burn, relative to the other treatments, rather than an increase in food resources. Although we did not find statistical support for a treatment difference in predatory invertebrate captures, the significant post-burn increase was driven by increased captures in the control and wildfire treatments (see Figure 2). Further, mesopredators may have dispersed out of the prescribed-burn unit during or following the burn. Jones et al. (2004) found raccoons selected for non-burned longleaf pine (*Pinus palustris*) forest, presumably because fire inhibited production of soft mast. However, we did not monitor vertebrate predators, and thus, have no hard evidence to support or refute this scenario.

The results of our study agree with observations by Means and Campbell (1982) and Grafe et al. (2002) that amphibians do not appear to be particularly vulnerable to direct mortality from fire. In contrast to *Bufo* and *Scaphiopus*, there were no noticeable treatment differences for *Gastrophryne*. However, species and genera-specific results are common in the amphibian fire literature (cf. Moseley et al. 2003; Hossack and Corn 2007). With these trend-based data, it is not possible to discern whether fire had no short-term influence on *Gastrophryne* captures or a balance between negative (e.g., direct mortality) and positive (e.g., above hypotheses) capture influences.

In conclusion, the most important finding of our study was that fire did not appear to negatively impact short-term terrestrial juvenile amphibian survivorship or health. Further, we found intriguing evidence that in some cases fire might benefit amphibians not only through longer-term habitat changes (Means and Moler 1979; Hossack and Corn 2007), but also through short-term changes such as reduction in predator pressure,

increased mobility potential, and increased food resources. However, we note that our interpretations were based on the assumption that seven day sampling periods were sufficient to accurately reflect differences among arrays for captures and body condition of amphibians. Further, we were unable to conduct multiple prescribed burns during the study period, and thus our ability to infer causation is limited. We encourage future research examining effects of fire on amphibian predators and predator-prey interactions. In addition to increasing causative understanding, this knowledge would help to elucidate the influence of burn season on amphibians and their community-level interactions, which influence both short-term and long-term outcomes of habitat restoration efforts.

Acknowledgements

This study was supported by the Texas Parks and Wildlife Department (TPWD) and United States Fish and Wildlife Service (USFWS) through a traditional Section 6 grant. M. Ray, J. Lattanzio, E. Feichtinger, D. Episcopo, K. Jamerson, C. Wiggins, and S. Dolino assisted with invertebrate collecting. B. Weckerly provided helpful statistical guidance. Three anonymous reviewers provided suggestions that improved this manuscript. The Capitol Area Council of the Boy Scouts of America provided access to the GLR, and we are appreciative for their continuing support of our research.

References

- Agee JK. 1996. Achieving conservation biology objectives with fire in the Pacific Northwest. *Weed Technology* 10:417–421.

- Al-Rabab'ah MA, Williams CG. 2004. An ancient bottleneck in the lost pines of central Texas. *Molecular Ecology* 13:1075–1084.
- Arnold SJ, Wassersug RJ. 1978. Differential predation on metamorphic anurans by garter snakes (*Thamnophis*): social behavior as a possible defense. *Ecology* 59:1014–1022.
- Babbitt LH, Babbitt CH. 1951. A herpetological study of burned-over areas in Dade County, Florida. *Copeia* 1951:79.
- Bebi P, Kulakowski D, Veblen TT. 2003. Interactions between fire and spruce beetles in a subalpine rocky mountain forest landscape. *Ecology* 84:362–371.
- Benson TJ, Dinsmore JJ, Hohman WL. 2007. Responses of plants and arthropods to burning and disking of riparian habitats. *Journal of Wildlife Management* 71:1949–1957.
- Brown DJ, Swannack TM, Forstner MRJ. *In press*. Herpetofaunal survey of the Griffith League Ranch in the Lost Pines ecoregion of Texas. *Texas Journal of Science*.
- Bryant VM Jr. 1977. A 16,000 year pollen record of vegetational change in central Texas. *Palynology* 1:143–156.
- Burton TM, Likens GE. 1975. Salamander populations and biomass in the Hubbard Brook Experimental Forest, New Hampshire. *Copeia* 1975:541–546.
- Cain MD, Wigley TB, Reed DJ. 1998. Prescribed fire effects on structure in uneven-aged stands of loblolly and shortleaf pines. *Wildlife Society Bulletin* 26:209–218.
- Cano PD, Leynaud GC. 2009. Effects of fire and cattle grazing on amphibians and lizards in northeastern Argentina (Humid Chaco). *European Journal of Wildlife Research* 56:411–421.
- Chelgren ND, Adams MJ, Bailey LL, Bury RB. 2011. Using multilevel spatial models to understand salamander site occupancy patterns after wildfire. *Ecology* 92:408–421.

- Child T, Phillips BL, Shine R. 2009. Does desiccation risk drive the distribution of juvenile cane toads (*Bufo marinus*) in tropical Australia? *Journal of Tropical Ecology* 25:193–200.
- Cummer MR, Painter CW. 2007. Three case studies of the effect of wildfire on the Jemez mountains salamander (*Plethodon neomexicanus*): microhabitat temperatures, size distributions, and a historical locality perspective. *Southwestern Naturalist* 52:26–37.
- Davic RD, Welsh HH Jr. 2004. On the ecological roles of salamanders. *Annual Review of Ecology, Evolution, and Systematics* 35:405–434.
- deMaynadier PG, Hunter ML Jr. 1999. Forest canopy closure and juvenile emigration by pool-breeding amphibians in Maine. *Journal of Wildlife Management* 63:441–450.
- Dixon JR. 2000. *Amphibians and reptiles of Texas*. 2nd edition. College Station: Texas A&M University Press.
- Ferguson AW, Forstner MRJ. 2006. A device for excluding predators from pitfall traps. *Herpetological Review* 37:316–317.
- Ford WM, Menzel MA, McGill DW, Laerm J, McCay TS. 1999. Effects of a community restoration fire on small mammals and herpetofauna in the southern Appalachians. *Forest Ecology and Management* 114:233–243.
- Forstner MRJ, Swannack TM. 2004. *The Houston toad in context*. Austin: U.S. Fish and Wildlife Service. Technical Report.
- Fowler JF, Cohen L, Jarvis P. 1998. *Practical statistics for field biology*. 2nd edition. Englands: John Wiley & Sons Ltd.
- Freed PS, Neitman K. 1988. Notes on predation on the endangered Houston toad, *Bufo houstonensis*. *Texas Journal of Science* 40:454–456.

- Grafe TU, Dobler S, Linsenmair KE. 2002. Frogs flee from the sound of fire. *Proceedings of the Royal Society of London B: Biological Sciences* 269:999–1003.
- Greenberg CH, Forrest TG, Waldrop T. 2010. Short-term response of ground-dwelling arthropods to prescribed fire and mechanical fuel reduction in a southern Appalachian upland hardwood forest. *Forest Science* 56:112–121.
- Gresswell RE. 1999. Fire and aquatic ecosystems in forested biomes of North America. *Transactions of the American Fisheries Society* 128:193–221.
- Greuter KL. 2004. Early juvenile ecology of the endangered Houston toad, *Bufo houstonensis* (Anura: Bufonidae). Master's thesis. Texas State University-San Marcos.
- Hillis DM, Hillis AM, Martin RF. 1984. Reproductive ecology and hybridization of the endangered Houston toad (*Bufo houstonensis*). *Journal of Herpetology* 18:56–72.
- Hossack BR, Corn PS. 2007. Responses of pond-breeding amphibians to wildfire: short-term patterns in occupancy and colonization. *Ecological Applications* 17:1403–1410.
- Jones DD, Conner LM, Storey TH, Warren RJ. 2004. Prescribed fire and raccoon use of longleaf pine forests: implications for managing nest predation? *Wildlife Society Bulletin* 32:1255–1259.
- Kerby JL, Richards-Hrdlicka KL, Storfer A, Skelly DK. 2010. An examination of amphibian sensitivity to environmental contaminants: are amphibians poor canaries? *Ecology Letters* 13:60–67.
- Kloor K. 2000. Returning America's forests to their 'natural' roots. *Science* 287:573–575.
- Kuhn M, Weston S, Wing J, Forester J. 2010. contrast: a collection of contrast methods. R package version 0.13. Available: <http://cran.r-project.org/web/packages/contrast/contrast.pdf> (August 2011).

- Lovaas AL. 1976. Introduction of prescribed burning to Wind Cave National Park. *Wildlife Society Bulletin* 4:69–73.
- Maindonald J, Braun J. 2003. *Data analysis and graphics using R: an example-based approach*. New York: Cambridge University Press.
- Martof BS. 1953. Territoriality in the green frog, *Rana clamitans*. *Ecology* 34:165–174.
- Matthews CE, Moorman CE, Greenberg CH, Waldrop TA. 2010. Response of reptiles and amphibians to repeated fuel reduction treatments. *Journal of Wildlife Management* 74:1301–1310.
- Means DB, Moler PE. 1979. The pine barrens treefrog: fire, seepage bogs, and management implications. Pages 77–83 in Odum RR, Landers L, editors. Atlanta: Georgia Game and Fish Division. Technical Bulletin WL4.
- Means DB, Campbell HW. 1982. Effects of prescribed burning on amphibians and reptiles. Pages 89–97 in Wood GW, editor. *Prescribed fire and wildlife in southern forests*. Georgetown, South Carolina: Belle W. Baruch Forest Science Institute. Available: <http://www.clemson.edu/extfor/Prescribed%20Fire%20and%20Wildlife%20in%20Southern%20Forests.pdf> (August 2011).
- Moretti M, Duelli P, Obrist MK. 2006. Biodiversity and resilience of arthropod communities after fire disturbance in temperate forests. *Oecologia* 149:312–327.
- Moseley KR, Castleberry SB, Schweitzer SH. 2003. Effects of prescribed fire on herpetofauna in bottomland hardwood forests. *Southeastern Naturalist* 2:475–486.
- National Park Service. 2003. *Fire monitoring handbook*. Boise, Idaho: Fire Management Program Center. Available: http://www.nps.gov/fire/download/fir_eco_FEMHandbook2003.pdf (August 2011).

- Pechmann JHK, Wilbur HM. 1994. Putting declining amphibian populations in perspective: natural fluctuations and human impacts. *Herpetologica* 50:65–84.
- Pilliod DS, Bury RB, Hyde EJ, Pearl CA, Corn PS. 2003. Fire and amphibians in North America. *Forest Ecology and Management* 178:163–181.
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. 2009. nlme: linear and non-linear mixed effects models. R package version 3.1–96. Available: <http://cran.r-project.org/web/packages/nlme/nlme.pdf> (August 2011).
- Pyne SJ. 2010. America's fires: a historical context for policy and practice. North Carolina: Forest History Society.
- Reading CJ, Clarke RT. 1995. The effects of density, rainfall and environmental temperature on body condition and fecundity in the common toad, *Bufo bufo*. *Oecologia* 102:453–459.
- Rochester CJ, Brehme CS, Clark DR, Stokes DC, Hathaway SA, Fisher RN. 2010. Reptile and amphibian responses to large-scale wildfires in southern California. *Journal of Herpetology* 44:333–351.
- Roe AW, Grayson KL. 2008. Terrestrial movements and habitat use of juvenile and emigrating adult eastern red-spotted newts, *Notophthalmus viridescens*. *Journal of Herpetology* 42:22–30.
- Russell KR, Van Lear DH, Guynn DC Jr. 1999. Prescribed fire effects on herpetofauna: review and management implications. *Wildlife Society Bulletin* 27:374–384.
- Ruthven DC III, Kzmaier RT, Janis MW. 2008. Short-term response of herpetofauna to various burning regimes in the south Texas plains. *Southwestern Naturalist* 53:480–487.

- Ryu S-R, Chen J, Zheng D, Bresee MK, and Crow TR. 2006. Simulating the effects of prescribed burning on fuel loading and timber production (EcoFL) in managed northern Wisconsin forests. *Ecological Modelling* 196:395–406.
- Saenz D, Fitzgerald LA, Baum KA, Conner RN. 2006. Abiotic correlates of anuran calling phenology: the importance of rain, temperature, and season. *Herpetological Monographs* 20:64–82.
- Schulte-Hostedde AI, Zinner B, Millar JS, Hickling GJ. 2005. Restitution of mass-size residuals: validating body condition indices. *Ecology* 86:155–163.
- Scrimgeour GJ, Tonn WM, Paszowski CA, Goater C. 2001. Benthic macroinvertebrate biomass and wildfires: evidence for enrichment of boreal subarctic lakes. *Freshwater Biology* 46:367–378.
- Semlitsch RD. 2008. Differentiating migration and dispersal processes for pond-breeding amphibians. *Journal of Wildlife Management* 72:260–267.
- Semlitsch RD, Scott DE, Pechmann JHK, Gibbons JW. 1996. Structure and dynamics of an amphibian community, evidence from a 16-year study of a natural pond. Pages 217–248 in Cody ML, Smallwood JA, editors. *Long-term studies of vertebrate communities*. New York: Academic Press.
- Smith MA, Green DM. 2006. Sex, isolation and fidelity: unbiased long-distance dispersal in a terrestrial amphibian. *Ecography* 29:649–658.
- Smucker KM, Hutto RL, Steele BM. 2005. Changes in bird abundance after wildfire: importance of fire severity and time since fire. *Ecological Applications* 15:1535–1549.

- Sparks JC, Masters RE, Engle DM, Palmer MW, Bukenhofer GA. 1998. Effects of late growing-season and late dormant-season prescribed fire on herbaceous vegetation in restored pine-grassland communities. *Journal of Vegetation Science* 9:133–142.
- Sunquist, ME. 1967. Effects of fire on raccoon behavior. *Journal of Mammalogy* 48:673–674.
- Swannack TM, Grant WE, Forstner MRJ. 2009. Projecting population trends of endangered amphibian species in the face of uncertainty: a pattern-oriented approach. *Ecological Modelling* 220:148–159.
- Sweeney JR, Biswell HH. 1961. Quantitative studies of the removal of litter and duff by fire under controlled conditions. *Ecology* 42:572–575.
- Taber SW, Fleenor SB. 2003. *Insects of the Texas lost pines*. College Station: Texas A&M University Press.
- Taber SW, Fleenor SB, Privette JL. 2008. Arthropod abundance following a prescribed burn in the lost pines forest of central Texas. *Southwestern Entomologist* 33:65–77.
- Tinker DB, Knight DH. 2000. Coarse woody debris following fire and logging in Wyoming lodgepole pine forests. *Ecosystems* 3:472–483.
- Toledo LF. 2005. Predation of juvenile and adult amphibians by invertebrates: current knowledge and perspectives. *Herpetological Review* 36:395–400.
- Vasconcelos D, Calhoun AJK. 2004. Movement patterns of adult and juvenile *Rana sylvatica* (LeConte) and *Ambystoma maculatum* (Shaw) in three restored pools in Maine. *Journal of Herpetology* 38:551–561.
- Vasconcelos HL, Pacheco R, Silva RC, Vasconcelos PB, Lopes CT, Costa AN, Bruna EM. 2009. Dynamics of the leaf-litter arthropod fauna following fire in a neotropical woodland savanna. *PLoS ONE* 4:e7762.

- Walls SC, Williams MG. 2001. The effect of community composition on persistence of prey with their predators in an assemblage of pond-breeding amphibians. *Oecologia* 128: 134–141.
- Wells KD. 2007. The ecology & behavior of amphibians. Illinois: University of Chicago Press.
- Welsh HH Jr., Ollivier LM. 1998. Stream amphibians as indicators of ecosystem stress: a case study from California's redwoods. *Ecological Applications* 8:1118–1132.
- Wright AH, Wright AA. 1949. Handbook of frogs and toads of the United States and Canada. 3rd edition. New York: Comstock Publishing Company.
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. 2009. Mixed effects models and extensions in ecology with R. New York: Springer Science+Business Media.

CHAPTER III

IMPACTS OF A HIGH SEVERITY WILDFIRE ON ABUNDANCE, MOVEMENT, AND DIVERSITY OF HERPETOFAUNA IN THE LOST PINES ECOREGION OF TEXAS

Format follows the guidelines for Herpetological Conservation and Biology

III. Impacts of a High Severity Wildfire on Abundance, Movement, and Diversity of Herpetofauna in the Lost Pines Ecoregion of Texas

Abstract

In September and October 2011, a high severity wildfire burned 39% of the 34,400-ha Lost Pines ecoregion in Bastrop County, Texas, USA. We assessed impacts of the wildfire on abundance, movement, and diversity of herpetofauna using drift fence array trap data collected prior to and after the wildfire, and anuran call survey data collected after the fire, on the 1,948-ha Griffith League Ranch. Based on *N*-mixture model analyses, abundance and movement of Six-lined Race Runners (*Cnemidophorus* [*Aspidoscelis*] *sexlineatus*) and Southern Prairie Lizards (*Sceloporus consobrinus*) were not significantly impacted by the wildfire shortly after its occurrence. A capture-recapture analysis indicated that movement rates were higher in the wildfire zone for Hurter's Spadefoot Toads (*Scaphiopus hurterii*) the following spring. Trap data indicated that the herpetofaunal species composition was not impacted by the wildfire shortly after the fire or subsequently, during the following spring. However, the anuran call survey data indicated that anuran species richness was higher in the wildfire zone. Collectively, it seems the wildfire had minimal negative impacts on abundance and diversity of herpetofauna in the short-term, a positive result for conservation in this ecoregion. In addition, our study indicated that investigations focused on fire impacts to ground-dwelling wildlife should consider detection probability when drawing inferences concerning abundances, particularly when differences in ground structure are apparent.

Key Words.—abundance; detection; diversity; fire; forest; herpetofauna; movement

Introduction

Pine-dominated forests throughout much of the U.S. are fire-maintained systems (Hartnett and Krofta 1989; Agee 1996; Schulte and Mladenoff 2005). In the absence of fire, flooding, and other disturbances, these forests progress towards a climax state dominated by hardwood trees (Gilliam and Platt 1999; Knebel and Wentworth 2007). Fire suppression over the last century altered the structure and composition of historically pine-dominated forests throughout the U.S. (Taylor 2000; Stephens and Ruth 2005; Nowacki and Abrams 2008). One issue associated with fire suppression is a consequent increase in fuel loads, which creates an environment conducive to high-severity fires (Davis 2001; Allen et al. 2002; Collins et al. 2010). Use of prescribed fire for reducing fuel loads and managing forest structure increased dramatically over the last half century. However, much of the U.S. forests remain severely fire-suppressed, and the frequency and severity of wildfires continues to increase in many regions (Houghton et al. 2000; Shang et al. 2007; Littell et al. 2009; Miller et al. 2009).

The Lost Pines ecoregion in Bastrop County, Texas, USA, is a 34,400-ha remnant patch of pine-dominated forest thought to have become finally isolated from the East Texas Piney Woods ecoregion between 10,000 and 14,000 y ago (Bryant 1977; Al-Rabah'ah and Williams 2004). The Lost Pines was extensively logged in the 1800s and early 1900s (Moore 1977). Fire suppression was implemented throughout the ecoregion since the early to mid-1900s, causing heavy fuel loads to accumulate. On 4 September 2011 a high-severity wildfire (i.e., the Bastrop County Complex Fire) began in the Lost

Pines. The fire was unstoppable due to extreme drought conditions in central Texas coupled with wind gusts in excess of 58 kph resulting from the passage of Tropical Storm Lee. The fire after 18 d was 95% contained, and the total burned area encompassed 13,406 ha. The wildfire breached a fire break on 11 October 2011, burning another 125-ha.

Management and conservation initiatives in the Lost Pines largely focus on promoting healthy non-game wildlife populations, particularly the federally endangered Houston Toad (*Bufo* [*Anaxyrus*] *houstonensis*). Most previous research indicates that direct mortality during fires is minimal (e.g., Cunningham et al. 2002; Fenner and Bull 2007; Radke et al. 2008; Ruthven et al. 2008), but the impacts of fire on herpetofauna remains severely understudied (Russell et al. 1999; Pilliod et al. 2003). However, few previous studies assessed high-severity forest fires, and thus there is little information available to gauge direct mortality of such wildfires to herpetofauna. The immediate concern from governmental agencies and landowners was the possibility of substantial direct mortality because of the catastrophic nature of the Bastrop County Complex Fire (Lost Pines Recovery Team 2011). In addition to direct mortality, another topic currently of interest to both managers and researchers is how herpetofaunal species detection probabilities are affected by fire. This is relevant given that fire can significantly modify ground structure, potentially influencing activity and visibility of species (Hossack and Corn 2007; Chelgren et al. 2011).

We have recently studied the use of prescribed fire for managing Houston Toad populations and other herpetofauna in the Lost Pines (Brown et al. 2011). As part of this research, we surveyed herpetofauna on our primary study property in July 2011, 6 weeks

prior to a wildfire. Here, we use these pre-burn data to investigate the possibility of significant direct herpetofaunal mortality during the Bastrop County Complex Fire. We also assessed movement of herpetofaunal species shortly after the fire, as well as the following spring, and determined whether species composition differed between burned and non-burned habitat in all three sampling periods. Finally, we use anuran call survey data collected in spring 2012 to determine whether species richness of calling anurans differed between burned and non-burned habitat. Collectively, these analyses provided useful insights into the short-term impacts of high-severity wildfires in a forest ecosystem, and the influence of detection on monitoring herpetofauna.

Materials and Methods

Study site.—We conducted this study on the 1,948-ha Griffith League Ranch (GLR), which primarily was forested with an overstory dominated by Loblolly Pine (*Pinus taeda*), Eastern Red Cedar (*Juniperus virginiana*), and Post Oak (*Quercus stellata*), and an understory dominated by Yaupon Holly (*Ilex vomitoria*), American Beautyberry (*Callicarpa Americana*), and Farkleberry (*Vaccinium arboreum*). The GLR contains three permanent ponds (i.e., ponds had not dried in at least 12 y), 10 semi-permanent ponds (i.e., ponds that typically dried several times per decade), and dozens of ephemeral pools that held water for days to months annually, depending on rainfall. Fire had been suppressed for at least 60 years prior to 2009 on the GLR, but since 2009 both prescribed burns and wildfires occurred on the study area. We conducted three low-severity prescribed burns between November 2009 and August 2010, burning ca. 378 ha. Two medium- to high-severity wildfires occurred on the GLR on 21 August 2010, which

together burned 189 ha. The high-severity wildfire on 4 September 2011 burned 987 ha, followed by a second wildfire on 4 October 2011, which burned an additional 80.5 ha (Fig. 3.1). Because of their intensity, and thus their potential to dramatically impact the forest ecosystem, burn breaks were installed during all fires to minimize their spread.




Figure 3.1. Aerial image of the Griffith League Ranch (GLR), Bastrop County, Texas, and its location with respect to a 13,406 ha wildfire that occurred in the Lost Pines ecoregion in September 2011, with a breach of the fire break resulting in an additional 125 ha burned in October 2011. Overlain on the image are the wildfires, drift fence arrays and ponds used to study the abundance, movement, and diversity of herpetofauna following the wildfires. The September 2011 wildfire burned 987 ha (50.7%) of the GLR, and the October 2011 wildfire burned 80.5 ha (4.1%) of the GLR.

Fire severity.—We assessed fire severity for the Bastrop County Complex Fire using vegetation plots (20 m by 50 m) randomly placed within forested habitat in 2008, and surveyed vegetation plots between 32-66 d after the wildfire using National Park Service (2003) fire-monitoring guidelines. Of the 31 vegetation plots on the GLR, 15 burned during the wildfire. We assessed burn severity to substrate within each plot using four 15 m transect lines, each consisting of four points spaced 5 m apart. We assigned points a burn severity ranking from one (heavily burned) to five (unburned). In addition, we estimated char height and recorded status (alive or dead) for all overstory trees (i.e., diameter at breast height > 15 cm) within vegetation plots.

Herpetofaunal sampling.—We used 18 Y-shaped and six linear drift fence arrays to trap herpetofauna. The linear drift fence arrays were located adjacent to, and parallel with, ponds to maximize amphibian captures, whereas the Y-shaped arrays were located up to 600 m from the nearest pond. The aboveground height of the drift fence array flashing was at least 18 cm, with the flashing buried ca. 10 cm belowground. We buried flashing ca. Y-shaped arrays consisted of three 15 m arms with a 19 L center bucket and a 19 L bucket at each arm terminus. Linear arrays consisted of a 15 m arm with a 19 L bucket at each end, and a double-throated funnel trap in the center of the array on each side of the flashing. We equipped pitfall traps with flotation devices to mitigate mortality during bucket flooding, and both pitfall and funnel traps had wet sponges to provide a moist environment. We also equipped pitfall traps with predator exclusion devices (Ferguson and Forstner 2006). Most traps within the burned area were destroyed by the fire, so we rebuilt them in their exact pre-burn locations. The burn breaks installed on the ranch during the September 2011 wildfire fortuitously resulted in a balanced sampling

design for the first post-wildfire sampling period, with 12 traps (nine Y-shaped arrays and three linear arrays) located in burned and control areas, respectively. The October 2011 wildfire burned the habitat surrounding two additional arrays, resulting in 14 traps (11 Y-shaped arrays and three linear arrays) located in burned areas, and 10 traps (seven Y-shaped arrays and three linear arrays) located in control areas, for the second post-wildfire sampling period.

We trapped herpetofauna for 7 d between 16-23 July 2011 (hereinafter pre-burn), 7 d between 25 September-2 October 2011 (hereinafter post-burn 1st), and 72 d between 19 February-30 April 2012 (hereinafter post-burn 2nd). We checked traps and processed herpetofaunal captures daily, estimated snout-vent length and tail length to the nearest 0.1 mm using digital calipers (Control Company, Friendswood, Texas, USA) or dial calipers (Wiha, Monticello, Minnesota, USA), and estimated weight to the nearest 0.1 g using spring scales (Pesola, Baar, Switzerland). We marked amphibians and lizards using toe clips (Ferner 2007), with amphibians marked individually and lizards cohort-marked.

We obtained sufficient captures of two lizard species, the Southern Prairie Lizard (*Sceloporus consobrinus*) and the Six-lined Race Runner (*Cnemidophorus* [*Aspidoscelis*] *sexlineatus*), to assess immediate wildfire impacts on abundance and detection probability. We obtained sufficient Hurter's Spadefoot Toad (*Scaphiopus hurterii*) captures and recaptures during the post-burn 2nd sampling period to determine if detection probabilities, and thus movement rates, differed between the control and wildfire habitats using a capture-recapture analysis. Finally, we determined if herpetofaunal species composition differed between the control and wildfire habitats during all three sampling periods.

In addition to sampling herpetofauna using traps, we also completed 24 anuran call surveys between 31 January-16 May 2012 following the protocol of Jackson et al. (2006). On each survey night we surveyed all ponds holding water ($n = 15-19$). We began surveys at dusk and surveyed each pond once per survey night for 5 min. We recorded the number of individuals heard calling for all detected species, unless the number of individuals calling was too large to count accurately (typically > 10 individuals). We used these data to determine if species richness of calling anurans (i.e., anurans engaged in breeding activity) differed between the unburned and burned locations. Thus, we assumed that a sufficient number of call surveys were completed to ensure that if a species was present at a pond it would have been detected at least once. We believe this was likely given previous work on call survey detection probabilities on the GLR (i.e., Jackson et al. 2006), and our knowledge of anuran calling activity on the GLR based on 11 years of call survey monitoring prior to this study.

Statistical analyses.—We used a N -mixture modeling approach to estimate herpetofaunal abundance and detection probability (P) prior to and shortly after the wildfire (Dail and Madsen 2011). This approach uses both spatial and temporal replication of count data to jointly estimate abundance and P (Royle 2004). Thus, it accounts for observed numbers being a product of both ecological and observational processes. We used the open population N -mixture model (i.e., pcountOpen) developed by Dail and Madsen (2011), using the software package unmarked (version 0.9-8) in program R (version 2.14.2; Fiske and Chandler 2011). We assumed our statistical populations were closed within survey periods, but open between survey periods. We used the ‘constant’ population dynamics model in pcountOpen (i.e., apparent survival and

recruitment were not explicitly linked in the model). In addition, we included habitat (control or wildfire) and time (pre-burn or post-burn) as covariates in estimations of P . We used Wald tests to assess the significance of the two covariates and their interaction ($\alpha = 0.05$). We determined the most appropriate distributions for our data sets (i.e., Poisson, zero-inflated Poisson, or negative binomial) by comparing the goodness-of-fit of each distribution using the ‘parboot’ function (Fiske and Chandler 2012. Overview of Unmarked: An R package for the Analysis of Data from Unmarked Animals. Available from <http://cran.r-project.org/web/packages/unmarked/vignettes/unmarked.pdf> [Accessed 1 September 2012]). We used a zero-inflated Poisson distribution for the Six-lined Race Runner analysis ($P = 0.29$), and a negative binomial distribution for the Southern Prairie Lizard analysis ($P = 0.31$).

Our data sets were not robust enough to achieve convergence when including habitat, time, and their interaction as covariates in estimations of apparent survival. Thus, to assess the short-term effect of the wildfire on abundance, we used the trap and time-specific estimated abundances as our count data in an additional analysis. We determined if habitat, time, and their interaction were significant predictors of Six-lined Race Runner and Southern Prairie Lizard abundance using generalized linear models with Poisson distributions (Zuur et al. 2009). We assessed assumptions of normality and homoscedasticity using residual plots. These analyses were conducted using program R (version 2.14.2).

We conducted a capture-recapture analysis to determine if apparent survival and P (and thus movement rates) of Hurter’s Spadefoot Toads differed between burned and non-burned habitat during the post-burn 2nd sampling period. Hurter’s Spadefoot Toads are

explosive breeders (Wells 1977), and thus not all individuals emerged (i.e., entered the statistical population) simultaneously. Thus, we analyzed these capture-recapture data using a Cormack-Jolly-Seber model structure, which assumes the population is open (Cormack 1964, Jolly 1965, Seber 1965), and which allowed us to estimate both apparent survival and P . We specified four groups in the analysis: adult males, adult females, unburned habitat, and burned habitat. Through preliminary analyses we determined data were too sparse to accommodate time effects on either apparent survival or P . Therefore, time was not considered as a possible constraint in subsequent analyses. Consequently, the analysis assumed apparent survival and P to be constant across the duration of the post-burn 2nd sampling period. The constraints we considered assessed whether apparent survival and P did not vary between each sex nor across burned and unburned habitats (denoted *constant*), varied between each sex (*sex*), varied across burned and unburned habitats (*habitat*), or varied between each sex as well as across habitats (*sex x habitat*). Model selection was based on AIC, corrected for small sample size (AIC_C), and AIC_C weights (i.e., the most parsimonious model had the smallest AIC_C and the largest AIC_C weight; Burnham and Anderson 1998). We conducted model averaging to estimate parameters in the event that models were competing (≤ 2 AIC_C unit deviance in the highest ranking models). We conducted this analysis using program MARK (White and Burnham 1999).

To assess if herpetofaunal species composition differed between burned and non-burned habitat, we used an ecological distance approach (Clarke 1993). We performed Analysis of Similarity (ANOSIM) tests, using the Kulczynski ecological distance equation (Kindt and Coe 2005), for the pre-burn, post-burn 1st, and post-burn 2nd

sampling periods. We included habitat as a predictor in the analyses to determine if sites were more similar within than between burned and unburned areas. For this statistical test, the computed test statistic (R) ranges from -1 to 1, with values near 0 indicating differences in species composition between burned and unburned habitats were small. We used a permutation test ($n = 1,000$) to assess the significance of R ($\alpha = 0.05$). For the post-burn 2nd sampling period, we did not include terrestrial juvenile anurans that metamorphosed during the sampling period in the analysis. We assumed individuals did not disperse between burned and unburned habitats within each sampling period, which was supported by our recapture data. We conducted these analyses using the software package BiodiversityR (version 1.6) in program R (version 2.14.2).

To determine if species richness of calling anurans differed between unburned and burned areas we used a species accumulation curve approach. This approach was appropriate because species richness estimates are affected by sample size (Kindt and Coe 2005), and the number of surveyed ponds in each habitat type was not identical ($n = 9$ [burned]; $n = 10$ [non-burned]). We calculated the average pooled species richness ($\pm$ SD) at each pond, within each habitat, for each sample size (i.e., 1 to 10 ponds surveyed), and assessed differences graphically using species accumulation curves. We conducted this analysis using the software package BiodiversityR (version 1.6) in program R (version 2.14.2).

Results

Fire severity.—Fire passed through 231 of the 240 substrate burn severity points during the September 2011 wildfire. The mean burn severity ranking for each plot ranged from

1–4.3, and the mean burn severity ranking among plots was 1.7 (Fig. 3.2). Of the 478 overstory trees that were alive in plots before the wildfire, 235 were killed during the fire (49.2% mortality). Seventy of these trees were not detected following the burn and were likely completely consumed by fire (i.e., 63.8% of overstory trees were either unaccounted for or dead). Mean estimated char height for all overstory trees was 7.3 m.



Figure 3.2. Terrestrial habitat around a drift fence array on the Griffith League Ranch (GLR), Bastrop County, Texas, USA, before (A) and after (B) a high severity wildfire on 4 September 2011. Nearly all of the understory vegetation, litter, duff, and coarse woody debris were consumed in most of the burned area, with substantial overstory tree mortality.

Herpetofaunal impacts.—We captured 13, 9, and 18 herpetofaunal species during the pre-burn, post-burn 1st, and post-burn 2nd sampling periods, respectively. We captured 205 unique individuals during the pre-burn sampling period, 131 unique individuals during the post-burn 1st sampling period, and 7,153 unique individuals during the post-burn 2nd sampling period, of which 5,455 were recently metamorphosed Hurter’s Spadefoot Toads (Table 3.1).

Table 3.1. Number of unique pre-burn (trapped 16 July 2011 to 23 July 2011), post-burn 1st (trapped 25 September 2011 to 2 October 2011), and post-burn 2nd (trapped 19 February 2012 to 30 April 2012) herpetofaunal captures on the Griffith League Ranch (GLR), Bastrop County, Texas, USA, using 24 drift fence arrays. The post-burn 1st sample included 12 arrays located in non-burned (Control) habitat, and 12 arrays located in burned (Wildfire) habitat. The post-burn 2nd sample included 10 arrays located in non-burned habitat, and 14 arrays located in burned habitat. The relative abundance of Hurter’s Spadefoot Toads (*Scaphopus hurterii*) in the wildfire habitat (post-burn 2nd) is biased low due to erosion issues following heavy rainfall events, which caused most of the wildfire habitat pitfall traps to completely fill with sand. Captured juvenile amphibians that entered the terrestrial landscape during the post-burn 2nd sampling period are shown in parentheses.

Species	Control			Wildfire ^a		
	Pre-burn	Post-burn 1 st	Post-burn 2 nd	Pre-burn	Post-burn 1 st	Post-burn 2 nd
<u>Amphibians</u>						
<i>Acris crepitans</i> (Blanchard’s Cricket Frog)	1	0	0	0	0	0
<i>Ambystoma tigrinum</i> (Tiger Salamander)	0	0	0	0	0	59
<i>Bufo [Incilius] nebulifer</i> (Coastal Plain Toad)	7	0	12 (1)	3	2	25 (4)
<i>Gastrophryne carolinensis</i> (Eastern Narrow-mouthed Toad)	1	0	2	2	2	4
<i>Gastrophryne olivacea</i> (Western Narrow- mouthed Toad)	5	4	14	3	10	11

Table 3.1 continued.

<i>Hyla cinerea</i> (Green Tree Frog)	0	0	0	1	0	0
<i>Hyla versicolor</i> (Gray Tree Frog)	0	0	0	0	0	4
<i>Pseudacris streckeri</i> (Strecker's Chorus Frog)	0	0	0	0	0	0 (13)
<i>Rana</i> [<i>Lithobates</i>] <i>catesbeiana</i> (American Bullfrog)	0	0	0	0	0	2
<i>Rana</i> [<i>Lithobates</i>] <i>sphenocephala</i> (Southern Leopard Frog)	1	0	2	2	0	10
<i>Scaphiopus hurterii</i> (Hurter's Spadefoot Toad)	0	0	468 (85)	0	0	939 (5,370)
<u>Lizards</u>						
<i>Scincella lateralis</i> (Little Brown Skink)	3	1	3	2	1	5
<i>Cnemidophorus</i> [<i>Aspidozelis</i>] <i>sexlineatus</i> (Six-lined Race Runner)	18	12	14	27	18	25
<i>Sceloporus consobrinus</i> (Southern Prairie Lizard)	77	45	42	48	33	30
<u>Snakes</u>						
<i>Agkistrodon contortrix</i> (Broad-banded Copperhead)	1	0	0	0	0	0
<i>Heterodon platirhinos</i> (Eastern Hog-nose)	0	1	1	0	0	0

Table 3.1 continued.

<i>Leptotyphlops [Rena]</i>	0	0	0	0	0	1
<i>dulcis</i> (Texas Threadsnake)						
<i>Micrurus tener</i> (Texas Coralsnake)	0	0	0	1	0	1
<i>Nerodia erythrogaster</i> (Plain-bellied Watersnake)	0	0	0	0	0	2
<i>Storeria dekayi</i> (Texas Brownsnake)	0	0	0	0	0	1
<i>Tantilla gracilis</i> (Flat- headed Snake)	1	0	1	0	1	2
<i>Thamnophis proximus</i> (Ribbon Snake)	0	0	0	1	1	0

^aWildfires occurred on 4 September 2011 and 4 October 2011, burning 987 ha (50.7%), and 80.5 ha (4.1%) of the study area, respectively.

Lizard abundance and detection.—We captured 45 and 97 unique Six-lined Racerunners and Southern Prairie Lizards during the pre-burn sampling period, respectively, and 30 and 31 unique Six-lined Racerunners and Southern Prairie Lizards during the post-burn 1st sampling period, respectively. We recaptured 1 Six-lined Racerunner and 7 Southern Prairie Lizards that were marked during the pre-burn sampling period. Cumulative estimated abundance of Six-lined Racerunners was 25.5 individuals in the control habitat and 25.4 individuals in the wildfire habitat during the pre-burn sampling period, and 69.4 individuals in the control habitat and 68.3 individuals in the wildfire habitat during the post-burn 1st sampling period. For Six-lined Racerunners, we did not find a significant habitat-time interaction for abundance ($Z = 0.03$, $df = 44$, $P = 0.977$), and abundance did not differ by habitat ($Z = -0.10$, $df = 44$, $P =$

0.924). However, abundance differed by time ($Z = -4.32$, $df = 44$, $P < 0.001$).

Cumulative estimated abundance of Southern Prairie Lizards was 299.6 individuals in the control habitat and 306.1 individuals in the wildfire habitat during the pre-burn sampling period, and 33.8 individuals in the control habitat and 36.6 individuals in the wildfire habitat during the post-burn 1st sampling period. For Southern Prairie Lizards, we did not find a significant habitat-time interaction for abundance ($Z = -0.23$, $df = 44$, $P = 0.819$), and abundance did not differ by habitat ($Z = 0.33$, $df = 44$, $P = 0.741$). However, abundance differed by time ($Z = 12.02$, $df = 44$, $P < 0.001$).

For Six-lined Race Runners, we did not find a significant habitat-time interaction for P ($Z = -0.24$, $P = 0.808$), and P did not differ by habitat ($Z = 1.63$, $P = 0.103$), or time ($Z = -1.46$, $P = 0.145$). Estimated detection probabilities were higher post-burn in both habitats, and were higher in the wildfire habitat than the control habitat (Fig. 3.3a). For Southern Prairie Lizards, we did not find a significant habitat-time interaction for P ($Z = -1.20$, $P = 0.231$), and P did not differ by time ($Z = 1.19$, $P = 0.233$). However, P differed by habitat ($Z = -2.76$, $P = 0.006$). Estimated detection probabilities were higher pre-burn in both habitats, and were higher in the control habitat than the wildfire habitat (Fig. 3.3b).




Figure 3.3. Estimated detection probabilities for Six-lined Race Runners (A) and Southern Prairie Lizards (B), before (Pre) and after (Post) high severity wildfire in control (C) and burn (B) habitats on the Griffith League Ranch (GLR), Bastrop County, Texas, USA. Parameter estimates are shown with 95% confidence intervals. The wildfire did not appear to alter detection probabilities for these species in the short-term, which were sampled using drift fence arrays.

Hurter's Spadefoot Toad detection.—We captured 1349 unique adult Hurter's Spadefoot Toads, 123 of which were recaptured at least once. The highest ranked model included no covariate for apparent survival, and habitat as the covariate for P (AIC_C weight = 0.17). However, there was little support for this model over the second highest ranked model ($\Delta AIC_C = 0.09$), which included habitat as a covariate for apparent survival, and no covariate for P (AIC_C weight = 0.17), or additional models that included

sex as a covariate (Table 3.2). Despite the lack of strong support for a single model, results were consistent, with both apparent survival and P for both males and females being higher in the burned than in the unburned habitat (Fig. 3.4).

Table 3.2. Model selection results from a capture-recapture analysis used to determine if apparent survival (S) and detection probability (P) varied by sex and between non-burned and burned habitat for adult Hurter's Spadefoot Toads (*Scaphiopus hurterii*) on Griffith League Ranch (GLR), Bastrop County, Texas, USA, during spring 2012. We sampled the population using 24 drift fence arrays, 14 of which were located in burned habitat. We used a Cormack-Jolly-Seber model structure, which assumes the population was open, and did not include time effects on estimates of S and P . Model selection was based on Akaike Information Criterion, corrected for small sample size (AIC_C). We used 1,349 unique individuals (742 male and 607 female) in the capture-recapture analysis.

Model	AIC_C	ΔAIC_C	AIC_C weight	Parameters
$S(.) P(\text{treatment})$	1537.8	0.00	0.17	3
$S(\text{treatment}) P(.)$	1537.9	0.09	0.17	3
$S(\text{sex}) P(\text{treatment})$	1538.3	0.47	0.14	4
$S(\text{treatment}) P(\text{sex})$	1538.5	0.67	0.12	4
$S(\text{treatment}) P(\text{treatment})$	1539.0	1.22	0.09	4
$S(\text{sex} \times \text{treatment}) P(.)$	1540.0	2.23	0.06	5
$S(.) P(\text{sex} \times \text{treatment})$	1540.4	2.65	0.05	5
$S(\text{sex} \times \text{treatment}) P(\text{treatment})$	1541.1	3.31	0.03	6
$S(.) P(.)$	1541.2	3.42	0.03	2
$S(\text{sex}) P(.)$	1541.6	3.78	0.03	3
$S(\text{treatment}) P(\text{sex} \times \text{treatment})$	1541.6	3.85	0.03	6
$S(.) P(\text{sex})$	1541.7	3.88	0.02	3
$S(\text{sex} \times \text{treatment}) P(\text{sex})$	1541.7	3.94	0.02	6
$S(\text{sex}) P(\text{sex} \times \text{treatment})$	1542.1	4.30	0.02	6
$S(\text{sex}) P(\text{sex})$	1543.4	5.57	0.01	4
$S(\text{sex} \times \text{treatment}) P(\text{sex} \times \text{treatment})$	1543.8	6.02	0.01	8




Figure 3.4. Adult male (M) and female (F) Hurter's Spadefoot Toad (*Scaphiopus hurterii*) apparent survival (S ; A) and detection probability (P ; B) estimates in non-burned (C) and burned (B) habitat during spring 2012 on Griffith League Ranch (GLR), Bastrop County, Texas, USA. Parameter estimates represent model-averaged results from competing models (see Table 2), and are shown with 95% confidence intervals. Estimates of both S and P were higher for both males and females in burned compared to non-burned habitat. However, support for an optimal model was low.

Herpetofaunal species composition.—We found no difference in herpetofaunal species composition between unburned and burned habitat during the pre-burn ($R = -0.020$, $P = 0.56$), post-burn 1st ($R = 0.021$, $P = 0.32$), or post-burn 2nd sampling periods ($R = -0.073$, $P = 0.90$). During the pre-burn sampling period we detected 10 herpetofaunal species in unburned habitat and 10 species in burned habitat; 7 species were captured in both habitat types. During the post-burn 1st sampling period we detected 4 species in unburned habitat and 8 species in burned habitat; 4 species were captured in both habitat types. During the post-burn 2nd sampling period we detected 10 species in unburned habitat and 17 species in burned habitat; 9 species were captured in both habitat types. During the post-burn 1st sampling period, all species captured in unburned habitat were also captured in burned habitat. During the post-burn 2nd sampling period, the only species captured in unburned habitat that was not captured in burned habitat was the Eastern Hog-nosed Snake (*Heterodon platirhinos*), which was represented by only one individual.

Species richness of calling anurans.—We found that total species richness of calling anurans was equal between ponds in unburned and burned habitat (10 total species detected). However, the species accumulation curves indicated that species richness of calling anurans was higher at ponds in burned habitat when at least eight ponds were surveyed (i.e., SD did not overlap; Fig. 3.5).




Figure 3.5. Species accumulation curves for calling anurans at ponds located within non-burned (wildfire; $n = 9$) and burned (control; $n = 10$) habitat on the Griffith League Ranch (GLR), Bastrop County, Texas, USA. We completed 24 anuran call surveys between 31 January 2012 and 16 May 2012. We surveyed all ponds holding water on each survey night ($n = 15$ to 19).

Discussion

Our results agree with most previous studies, which concluded mortality from fire was minimal for amphibians and reptiles (e.g., Cunningham et al. 2002; Fenner and Bull 2007; Greenberg and Waldrop 2008; Radke et al. 2008; Brown et al. 2011; Milanovich et al. 2011). Observational studies during prescribed burns documented lizards escaping fire by burrowing under the soil and climbing trees (Bishop and Murrie 2004; Beane 2006), and Grafe et al. (2002) suggested that surface-aestivating anurans respond to auditory cues of approaching fire by seeking burn-resistant refugia. Most individuals that survived this wildfire were likely underground, given that crowning (i.e., fire in the tree canopy) was prevalent. For herpetofauna that were active at the time of the wildfire, with the possible exception of the terrestrial Three-toed Box Turtle (*Terrapene carolina triunguis*) and any dispersing semi-aquatic turtles, rapid burrowing under the soil was probably not difficult given the predominance of deep sandy soils in the Lost Pines ecoregion (Baker et al. 1979). While surveying and searching within the burn zone, we did find several Red-eared Slider (*Trachemys scripta elegans*) turtle shell fragments, and a partial Three-toed Box Turtle plastron, but cannot be certain those turtles died from the wildfire.

Our results indicated that detection probability (P) of two lizard species was not affected by the wildfire in the short-term. This gives some credence to results from previous studies using drift fence sampling that did not model P when assessing responses of lizards to fire, which as far as we are aware, includes all but one previous study (Driscoll et al. 2012). However, we note that Driscoll et al. (2012) found evidence that the detection probability model they used, which included a daily detection/non-

detection framework (as opposed to individual-based detection probabilities used here), did not capture what were likely differences in activity levels between burned and unburned areas. We also note that our P estimates lacked precision, as indicated by broad confidence intervals (see Fig. 3). Alternately, we obtained some evidence that P for Hurter's Spadefoot Toad was higher in the burned than in the unburned habitat. Chelgren et al. (2011) concluded that a wildfire positively influenced detection probabilities for five salamander species sampled using active searches. As far as we are aware, ours is the first study to indicate this phenomenon for amphibians using drift fences, which suggests that movement rates can increase after fires. However, we cannot conclude from these data what factors changed following fire that resulted in our observations, and thus we do not know if this was a general fire effect or a habitat-specific effect. An additional observation during the post-burn 2nd sampling period on adult Tiger Salamanders (*Ambystoma tigrinum*) is worth noting. Between spring 2008 and summer 2011 we trapped for 387 days, including 56 to 89 days each year between February and April, using 4 drift fence arrays located near a GLR pond within what would become the high-severity wildfire zone (Pond 9), and captured two Tiger Salamanders. During the 72 days of trapping in spring 2012, we captured 57 unique individuals near Pond 9. These data indicate that either movement rates or movement distances increased dramatically for Tiger Salamanders following the wildfire, or alternately some other factor led to a seemingly dramatic increase in detection for this species after the wildfire.

The evidence for higher anuran species richness at ponds in the wildfire habitat indicated that anurans were not only present in burned habitat, but used it for reproduction. Thus, despite the dramatic habitat changes, the post-fire landscape was not

perceived by anurans to be unsuitable habitat. This result agrees with previous studies that addressed impacts of a single wildfire on amphibian habitat use (Kirkland et al. 1996; Cummer and Painter 2007; Guscio et al. 2007; Hossack and Corn 2007). Our results indicated that pond occupancy within burned habitat may have increased after wildfire, and we intend to assess this possibility across the Lost Pines ecoregion in the coming years. As discussed earlier, the reduced vegetation and ground structural complexity could have increased movement rates, and thus potentially dispersal distances. In addition, we hypothesize that anuran calls traveled further in the burned habitat because of reduced vegetation, which could assist with pond colonization by breeding anurans. Hossack and Corn (2007) found that pond occupancy by Boreal Toads (*Bufo* [*Anaxyrus*] *boreas*) increased dramatically following a wildfire in Montana. If the endangered Houston Toad responds similarly in the Lost Pines, this could be viewed as a positive effect of wildfire. During spring 2012 anuran call surveys throughout the Lost Pines ecoregion, we detected Houston Toads at ponds in both unburned and burned habitat, including several ponds within the burn zone where we had not detected them during call surveys over the last decade. However, we note that because this species has critically low numbers (Duarte et al. 2011), and Houston Toad breeding success at a given pond increases exponentially with number of calling males (Gaston et al. 2010), increased occupancy in suboptimal habitat (e.g., ponds located within residential areas or similar small forest fragments) may actually expedite the extirpation of this species from the Lost Pines.

The limited mortality inferred from our analyses is a positive result for management of the endangered Houston Toad and other threatened herpetofauna (i.e., Texas Horned

Lizard *Phrynosoma cornutum* and Timber Rattlesnake) found in the Lost Pines (Brown et al., in press). Despite the positive immediate wildfire effects on herpetofauna found in this study, indirect effects (e.g., prey reduction, litter and coarse woody debris removal, overstory tree thinning, water quality impacts) are still a major concern. In our opinion, the most critical impact of the wildfire was overstory tree loss. Both the endangered Houston Toad and threatened Timber Rattlesnake prefer heavily canopied environments (U.S. Fish and Wildlife Service 1984; Brown 1993). The estimated loss of 63.8% of overstory trees on the GLR was similar to an overall projection from the Texas Forest Service for the entire burned area (78%; Lost Pines Recovery Team 2011), and it will take decades for these areas to return to mature forest.

Although most of the remaining wild Houston Toads are currently in the Lost Pines ecoregion (Brown 1971, 1975), it would be valuable to focus some future recovery efforts (e.g., headstarting, Vandewege et al. 2012) on other geographically disjunct populations to prevent against extinction in the wild, given the unknown trajectory of the Lost Pines population. This latter point is supported by previous studies emphasizing that probability of extinction in the Lost Pines will likely increase dramatically with a catastrophic wildfire (Seal 1994), or lack of multiple viable populations (Hatfield et al. 2004).

In conclusion, this study adds to the growing literature indicating high severity fires are not inherently detrimental to amphibians and reptiles, even in the immediate and short-term. Further, the wildfire could increase herpetofaunal diversity in the coming decades due to habitat diversification (e.g., the creation of large patches of early-succession forest habitat). We intend to continue monitoring herpetofauna in the Lost Pines to document

population and community trends in relation to natural forest recovery and active restoration initiatives.

Acknowledgements

This study was supported by the Texas Parks and Wildlife Department and U. S. Fish and Wildlife Service through a traditional Section 6 grant. Joseph Barnett, Devin Preston, Derek Wallace, Mindy Murray, Jennifer Oakley, Chris Tsay, and Amy Connolly assisted with rebuilding traps. Amber Harper, Devin Preston, Derek Wallace, Dionne Episcopo, Emrah Ozel, and Tyrone Davis assisted with trapping. Sara Weaver, Devin Preston, and Derek Wallace assisted with call surveys. The Capitol Area Council of the Boy Scouts of America provided access to the Griffith League Ranch, and we appreciate their continuing support of our research. Handling permits were provided by the Texas Parks and Wildlife Department (permit SPR-0102-191), and U.S. Fish and Wildlife Service (permit TE 039544-0). Trapping and handling methods were approved by the Texas State University-San Marcos Institutional Animal Care and Use Committee (Protocol No. 1012_0501_11).

Literature Cited

- Agee, J.K. 1996. Achieving conservation biology objectives with fire in the Pacific Northwest. *Weed Technology* 10:417–421.
- Al-Rabah'ah, M.A., and C.G. Williams. 2004. An ancient bottleneck in the Lost Pines of central Texas. *Molecular Ecology* 13:1075–1084.

- Allen, C.D., M. Savage, D.A. Falk, K.F. Suckling, T.W. Swetnam, T. Schulke, P.B. Stacey, P. Morgan, M. Hoffman, and J.T. Klingel. 2002. Ecological restoration of southwestern Ponderosa Pine ecosystems: a broad perspective. *Ecological Applications* 12:1418–1433.
- Baker, F.E. 1979. Soil survey of Bastrop County, Texas. U.S. Department of Agriculture, Soil Conservation Service. Washington, D.C., USA. 73 p.
- Beane, J.C. 2006. *Sceloporus undulatus hyacinthinus* (Northern Fence Lizard) fire avoidance behavior. *Herpetological Review* 37:92.
- Bess, E.C., R.R. Parmenter, S. McCoy, and M.C. Molles, Jr. 2002. Responses of a riparian forest-floor arthropod community to wildfire in the middle Rio Grande Valley, New Mexico. *Environmental Entomology* 31:774–784.
- Bishop, D.C., and T.W. Murrie. 2004. *Sceloporus undulatus* (Eastern Fence Lizard) fire avoidance behavior. *Herpetological Review* 35:397–398.
- Brown, D.J., J.T. Baccus, D.B. Means, and M.R.J. Forstner. 2011. Potential positive effects of fire on juvenile amphibians in a southern USA pine forest. *Journal of Fish and Wildlife Management* 2:135–145.
- Brown, D.J., T.M. Swannack, and M.R.J. Forstner. In press. Herpetofaunal survey of the Griffith League Ranch in the Lost Pines Ecoregion of Texas. *Texas Journal of Science*.
- Brown, L.E. 1971. Natural hybridization and trend toward extinction in some relict Texas Toad populations. *Southwestern Naturalist* 16:185–199.
- Brown, L.E. 1975. The status of the near-extinct Houston Toad (*Bufo houstonensis*) with recommendations for its conservation. *Herpetological Review* 6:37–40.

- Brown, W.S. 1993. Biology, status, and management of the Timber Rattlesnake (*Crotalus horridus*): A guide for conservation. Society for the Study of Amphibians and Reptiles, Oxford, Ohio, USA. 78 p.
- Bryant, V.M., Jr. 1977. A 16,000 year pollen record of vegetational change in central Texas. *Palynology* 1:143–156.
- Buddle, C.M., D.W. Langor, G.R. Pohl, and J.R. Spence. 2006. Arthropod responses to harvesting and wildfire: Implications for emulation of natural disturbance in forest management. *Biological Conservation* 128:346–357.
- Burnham, K.P., and D.R. Anderson. 1998. Model Selection and Multimodel Inference: A Practical Information-theoretic Approach. 2nd edition. Springer–Verlag, New York, New York, USA. 518 p.
- Chelgren, N.D., M.J. Adams, L.L. Bailey, and R.B. Bury. 2011. Using multilevel spatial models to understand salamander site occupancy patterns after wildfire. *Ecology* 92:408–421.
- Clarke, K.R. 1993. Non-parametric multivariate analysis of changes in community structure. *Australian Journal of Ecology* 18:117–143.
- Collins, B.M., S.L. Stephens, J.J. Moghaddas, and J. Battles. 2010. Challenges and approaches in planning fuel treatments across fire-excluded forested landscapes. *Journal of Forestry* 108:24–31.
- Cormack, R.M. 1964. Estimates of survival from the sighting of marked animals. *Biometrika* 51:429–438.

- Cummer, M.R., and C.W. Painter. 2007. Three case studies of the effect of wildfire on the Jemez Mountains Salamander (*Plethodon neomexicanus*): microhabitat temperatures, size distributions, and historical locality perspective. *Southwestern Naturalist* 52: 26–37.
- Cunningham, S.C., R.D. Babb, T.R. Jones, B.D. Taubert, and R. Vega. 2002. Reaction of lizard populations to a catastrophic wildfire in a central Arizona mountain range. *Biological Conservation* 107:193–201.
- Dail, D., and L. Madsen. 2011. Models for estimating abundance from repeated counts of an open metapopulation. *Biometrics* 67:577–587.
- Davis, C. 2001. The west in flames: the intergovernmental politics of wildfire suppression and prevention. *Publius* 31:97–110.
- Driscoll, D.A., A.L. Smith, S. Blight, and J. Maindonald. 2012. Reptile responses to fire and the risk of post-disturbance sampling bias. *Biodiversity and Conservation* 21:1607–1625.
- Duarte, A., D.J. Brown, and M.R.J. Forstner. 2011. Estimating abundance of the endangered Houston toad on a primary recovery site. *Journal of Fish and Wildlife Management* 2:207–215.
- Fenner, A.L., and C.M. Bull. 2007. Short-term impact of grassland fire on the endangered Pygmy Bluetongue Lizard. *Journal of Zoology* 272:444–450.
- Ferguson, A.W., and M.R.J. Forstner. 2006. A device for excluding predators from pitfall traps. *Herpetological Review* 37:316–317.

- Ferner, J.W. 2007. A Review of Marking and Individual Recognition Techniques for Amphibians and Reptiles. Society for the Study of Amphibians and Reptiles, Oxford, Ohio, USA. 72 p.
- Fiske, I. J., and R. B. Chandler. 2011. Unmarked: an R package for fitting hierarchical models of wildlife occurrence and abundance. *Journal of Statistical Software* 43:1–23.
- Gaston, M.A., A. Fuji, F.W. Weckerly, and M.R.J. Forstner. 2010. Potential component allee effects and their impact on wetland management in the conservation of endangered anurans. *PLoS One*:e10102.
- Gilliam, F.S., and W.J. Platt. 1999. Effects of long-term fire exclusion on tree species composition and stand structure in an old-growth *Pinus palustris* (Longleaf Pine) forest. *Plant Ecology* 140:15–26.
- Grafe, T.U., S. Dobler, and K.E. Linsenmair. 2002. Frogs flee from the sound of fire. *Proceedings of the Royal Society of London B: Biological Sciences* 269:999–1003.
- Greenberg, C.H., and T.A. Waldrop. 2008. Short-term response of reptiles and amphibians to prescribed fire and mechanical fuel reduction in a southern Appalachian upland hardwood forest. *Forest Ecology and Management* 255:2883–2893.
- Guscio, C.G., B.R. Hossack, L.A. Eby, and P.S. Corn. 2007. Post-breeding habitat use by adult Boreal Toads (*Bufo boreas*) after wildfire in Glacier National Park, USA. *Herpetological Conservation and Biology* 3:55–62.
- Hartnett, D.C., and D.M. Krofta. 1989. Fifty-five years of post-fire succession in a southern mixed hardwood forest. *Bulletin of the Torrey Botanical Club* 116:107–113.

- Hatfield, J.S., A.H. Price, D.D. Diamond, and C.D. True. 2004. Houston Toad (*Bufo houstonensis*) in Bastrop County, Texas: Need for Protecting Multiple Subpopulations. Pp. 292–298 *In* Species Conservation and Management: Case Studies. Akcakaya, H.R., M.A. Burgman, O. Kindvall, C.C. Wood, P. Sjogren-Gulve, J.S. Hatfield, and M.A. McCarthy (Eds.). Oxford University Press, New York, New York, USA.
- Hossack, B.R., and P.S. Corn. 2007. Responses of pond-breeding amphibians to wildfire: short-term patterns in occupancy and colonization. *Ecological Applications* 17:1403–1410.
- Houghton, R.A., J.L. Hackler, and K.T. Lawrence. 2000. Changes in terrestrial carbon storage in the United States 2: the role of fire and fire management. *Global Ecology and Biogeography* 9:145–170.
- Jackson, J.T., F.W. Weckerly, T.M. Swannack, and M.R.J. Forstner. 2006. Inferring absence of Houston toads given imperfect detection probabilities. *Journal of Wildlife Management* 70: 1461–1463.
- Jolly, G.M. 1965. Explicit estimates from capture-recapture data with both death and immigration-stochastic model. *Biometrika* 52:225–247.
- Kindt, R., and R. Coe. 2005. Tree diversity analysis: a manual and software for common statistical methods for ecological and biodiversity studies. World Agroforestry Centre, Nairobi, Kenya. 197 p.
- Kirkland, Jr., G.L., H.W. Snoddy, and T.L. Amsler. 1996. Impact of fire on small mammals and amphibians in a central Appalachian deciduous forest. *American Midland Naturalist* 135:253–260.

- Knebel, L., and T.R. Wentworth. 2007. Influence of fire and Southern Pine Beetle on pine-dominated forests in the Linville Gorge Wilderness, North Carolina. *Castanea* 72:214–225.
- Littell, J.S., D. McKenzie, D.L. Peterson, and A.L. Westerling. 2009. Climate and wildfire area burned in western U.S. ecoprovinces, 1916–2003. *Ecological Applications* 19:1003–1021.
- Lost Pines Recovery Team. 2011. Bastrop County complex fire, Lost Pines region, resources assessment and response report. Lost Pines Recovery Team, Bastrop, Texas, USA. 247 p.
- Milanovich, J.R., S.E. Trauth, and D.A. Saugey. 2011. Reproduction and age composition of a population of Woodland Salamanders (*Plethodon albagula*) after a prescribed burn in southwestern Arkansas. *Southwestern Naturalist* 56:172–179.
- Miller, J.D., H.D. Safford, M. Crimmins, and A.E. Thode. 2009. Quantitative evidence for increasing forest fire severity in the Sierra Nevada and Southern Cascade mountains, California and Nevada, USA. *Ecosystems* 12:16–32.
- Moore, B. 1977. Bastrop County 1691–1900. Revised Edition. Nortex Press, Wichita Falls, Texas, USA. 318 p.
- National Park Service. 2003. Fire monitoring handbook. Fire Management Program Center, Boise, Idaho, USA. 274 p.
- Neumann, F.G. 1991. Responses of litter arthropods to major natural or artificial ecological disturbances in Mountain Ash forest. *Australian Journal of Ecology* 16:19–32.

- Nowacki, G.J., and M.D. Abrams. 2008. The demise of fire and "mesophication" of forests in the eastern United States. *Bioscience* 58:123–138.
- Parmenter, R.R., M. Kreutzian, D.I. Moore, and D.C. Lightfoot. 2011. Short-term effects of a summer wildfire on a desert grassland arthropod community in New Mexico. *Environmental Entomology* 40:1051–1066.
- Pilliod, D.S., R.B. Bury, E.J. Hyde, C. A. Pearl, and P.S. Corn. 2003. Fire and amphibians in North America. *Forest Ecology and Management* 178:163–181.
- Radke, N.J., D.B. Wester, G. Perry, and S. Rideout-Hanzak. 2008. Short-term effects of prescribed fire on lizards in Mesquite-Ashe Juniper vegetation in central Texas. *Applied Herpetology* 5:281–292.
- Royle, J.A. 2004. *N*-mixture models for estimating population size from spatially replicated counts. *Biometrics* 60:108–115.
- Russell, K.R., D.H.V. Lear, and D.C. Guynn, Jr. 1999. Prescribed fire effects on herpetofauna: review and management implications. *Wildlife Society Bulletin* 27:374–384.
- Ruthven, D.C., III, R.T. Kzmaier, and M.W. Janis. 2008. Short-term response of herpetofauna to various burning regimes in the south Texas plains. *Southwestern Naturalist* 53:480–487.
- Schulte, L.A., and D.J. Mladenoff. 2005. Severe wind and fire regimes in northern forests: historical variability at the regional scale. *Ecology* 86:431–445.
- Seal, U.S. 1994. Population and habitat viability assessment: Houston Toad (*Bufo houstonensis*). IUCN/SSC Conservation Breeding Specialist Group, Apple Valley, Minnesota, USA. 127 p.

- Seber, G.A. 1965. A note on the multiple-recapture census. *Biometrika* 52:249–259.
- Shang, Z., H.S. He, D.E. Lytle, S.R. Shifley, and T.R. Crow. 2007. Modeling the long-term effects of fire suppression on central hardwood forests in Missouri Ozarks, using LANDIS. *Forest Ecology and Management* 242:776–790.
- Stephens, S.L., and L.W. Ruth. 2005. Federal forest-fire policy in the United States. *Ecological Applications* 15:532–542.
- Taylor, A.H. 2000. Fire regimes and forest changes in mid and upper montane forests of the southern Cascades, Lassen Volcanic National Park, California, U.S.A. *Journal of Biogeography* 27:87–104.
- U. S. Fish and Wildlife Service. 1984. Houston Toad recovery plan. U.S. Fish and Wildlife Service, Albuquerque, New Mexico, USA. 73 p.
- Vandewege, M.W., D.J. Brown, and M.R.J. Forstner. 2012. *Bufo* (= *Anaxyrus*) *houstonensis* (Houston Toad) head-start juvenile dispersal. *Herpetological Review* 43:117–118.
- Wells, K.D. 1977. The social behaviour of anuran amphibians. *Animal Behaviour* 25:666–693.
- White, G.C., and K.P. Burnham. 1999. Program MARK: survival estimation from populations of marked animals. *Bird Study* 46:120–139.
- Zuur, A.F., E.N. Leno, N.J. Walker, A.A. Saveliev, and G.M. Smith. 2009. *Mixed Effects Models and Extensions in Ecology with R*. Springer Science + Business Media, New York, New York, USA. 574 p.

CHAPTER IV

COMPARISON OF LOW, MODERATE, AND HIGH SEVERITY FIRE IMPACTS TO AQUATIC AND TERRESTRIAL ECOSYSTEM COMPONENTS OF A SOUTHERN USA MIXED PINE/HARDWOOD FOREST

Format follows the guidelines for Forest Ecology and Management

IV. Comparison of Low, Moderate, and High Severity Fire Impacts to Aquatic and Terrestrial Ecosystem Components of a Southern USA Mixed Pine/Hardwood Forest

ABSTRACT

Historically fire was an important natural disturbance shaping the structure and composition of pine-dominated forests in the southern United States. Longstanding fire suppression policies have resulted in structural and compositional changes, notably accumulation of heavy fuel loads and reduction in vegetation species diversity. Primary goals of forest management through prescribed burning include fuel load reduction and mimicking ecosystem impacts of historically natural wildfires. In addition to the influences of fire frequency and season, the influence of fire severity on ecosystem responses is currently of interest. In this study we assessed the impacts of low, moderate, and high severity fires to several aquatic and terrestrial ecosystem components of a southern U.S. mixed pine/hardwood forest using a before-after, control-impact (BACI) approach. The ecosystem components we assessed were water quality, community composition of aquatic arthropods (wildfire impacts only), forest structure characteristics, community composition of understory vegetation, and community composition of ground-dwelling arthropods. We found that fire severity was an important factor influencing the response of the ecosystem components we assessed. Fire at all severity levels increased aquatic nutrient levels and productivity, but the magnitude of effects increased with severity. Low and moderate severity fires had weak effects on forest structure characteristics, community composition of understory vegetation, and

community composition of ground-dwelling arthropods. In contrast, high severity fires dramatically reduced fine and large fuel loads, increased diversity of understory vegetation, and influenced community composition of ground-dwelling arthropods. However, we did not detect a fire response for aquatic arthropods. Our results contribute to the growing evidence that fire severity is a primary factor influencing responses of ecosystems to fire.

Keywords:

Arthropods

Forest

Prescribed fire

Vegetation

Water chemistry

Wildfire

Highlights:

1. The magnitude of fire effects on water quality increased with increasing fire severity.
2. Low and moderate severity fires had weak effects on forest structure characteristics and species composition of understory vegetation, whereas high severity fires dramatically reduced fine and large fuel loads and increased tree mortality and diversity and cover of understory vegetation.
3. Low and moderate severity fires had no detectable effects on species composition of ground-dwelling arthropods, whereas high severity fires influenced species composition.

1. Introduction

Climatic trends towards warmer and drier conditions, coupled with longstanding broad-scale fire suppression, have resulted in an increase in frequency of high severity wildfires in the southern and western United States (Davis, 2001; Miller *et al.*, 2009), with this trend projected to continue into the next century (Moritz *et al.*, 2012). The increase in wildfires is prevalent in pine-dominated forests (Miller *et al.*, 2009), which are naturally fire-maintained systems (Hartnett and Krofta, 1989; Schulte and Mladenoff, 2005). In the absence of fire, these forests typically progress towards a climax state dominated by hardwood trees (Gilliam and Platt, 1999; Knebel and Wentworth, 2007; Hanberry *et al.*, 2012). Further, suppression-induced increase in fuel loads often creates environments conducive to abnormally high severity wildfires (Davis, 2001; Allen *et al.*, 2002; Collins *et al.*, 2010). Thus, integration and maintenance of fire management is necessary for restoration and sustainability of healthy pine-dominated forests (Agee, 1996). Although the use of prescribed fire for reducing fuel loads and managing forest communities has increased dramatically over the last half century, much of the U.S. remains severely fire-suppressed (Houghton *et al.*, 2000; Shang *et al.*, 2007; Gebert and Black, 2012).

In addition to reducing fuel loads, a common goal of prescribed burning is to mimic ecosystem impacts of historically natural wildfires within a controlled setting (Vose, 2000). Thus, increasing our knowledge of similarities and differences between prescribed fire and wildfire, with respect to ecosystem impacts, is of interest to both ecologists and land managers (Schwilk *et al.*, 2006; Glasgow and Matlack, 2007; Arkle and Pilliod, 2010). This question has resulted in much research being devoted to effects

of burn timing and burn frequency, particularly on vegetation (e.g., Cain *et al.*, 1998; Sparks *et al.*, 1998; Taylor, 2000; Webster and Halpern, 2010). However, much less is known about effects of burn severity, given that prescribed burns are typically low or moderate severity fires (Knapp *et al.*, 2009). In contrast, high severity wildfires are unplanned, and thus wildfire research is by necessity a response to non-designed treatments, and opportunities for comparative research are more limited.

Clearly, fire impacts many terrestrial ecosystem components, ranging from soil properties (Certini, 2005), to habitat selection of large mammals (Long *et al.*, 2009). Fire has also been found to impact aquatic ecosystems in pine-dominated forests (e.g., Spencer and Hauer, 1991; Battle and Golladay, 2003). These aquatic ecosystems provide habitat for a variety of vertebrates (e.g., amphibians and fish) and invertebrates, with many threatened and endangered species dependent upon the presence and quality of aquatic environments in forested landscapes, such as the endangered Houston toad (*Bufo* [*Anaxyrus*] *houstonensis*), and threatened gila trout (*Oncorhynchus gilae*). With the exception of direct effects through increased water temperatures (Hitt, 2003) and burning when aquatic areas are dry (Sacerdote and King, 2009), fire primarily impacts aquatic components indirectly through a variety of impacts to the surrounding and adjacent terrestrial environment (Spencer and Hauer, 1991; Gresswell, 1999). Due to the interconnected nature of aquatic and terrestrial systems, fire research that incorporates effects on both ecosystem types is needed to improve our understanding of fire impacts to ecosystems (Bisson *et al.*, 2003; Rieman *et al.*, 2003).

Previous research suggests there are consistencies with regards to fire impacts on nutrients in both terrestrial and aquatic environments. In the short-term, fire often

increases soil nitrogen (N) and phosphorus (P) availability through conversion of organic to inorganic forms, with fire severity positively related to the magnitude of inorganic nutrient increases (Wan *et al.*, 2001; Certini, 2005). Similarly, N and P in both still and flowing waters typically increases after fire (Battle and Golladay, 2003; Earl and Blinn, 2003). In both terrestrial and aquatic environments, nutrient availability tends to decrease to background levels over a period of weeks to a few years (Wan *et al.*, 2001; Earl and Blinn, 2003; Spencer *et al.*, 2003). However, in some cases fire can impact terrestrial and aquatic nutrient levels for decades (McEachern *et al.*, 2000; Duran *et al.*, 2010).

Fire can directly impact biotic communities through both heat-induced mortality (animals and plants) and heat-induced reproduction (plants; Gauthier *et al.*, 1996; Schwilk *et al.*, 2006; Engstrom, 2010). Fire can indirectly impact biotic communities through alteration of nutrient availability (Lewis, 1974; Gilliam, 1988; Battle and Golladay, 2003), structural habitat modification (e.g., removal or addition of debris; Sweeney and Biswell, 1961; Tinker and Knight, 2000; Hall *et al.*, 2006), and alteration of inter- and intraspecific interactions. Thus, fire effects on biotic ecosystem components are inherently complex and difficult to extrapolate from one ecosystem to another. Further, the impacts of high severity fires are typically not short-lived, but rather a fire can influence a given ecosystem from decades to centuries (Hall *et al.*, 2006; Lecomte *et al.*, 2006; Webster and Halpern, 2010).

Fire can influence plant community composition primarily through direct impacts on mortality and reproduction (Gauthier *et al.*, 1996; Simmons *et al.*, 2007), changes in nutrient availability (Wan *et al.*, 2001), and alteration of ground structure (e.g., soil organic matter), which influences spacing of individual plants and light availability

(Sweeney and Biswell, 1961; Tinker and Knight, 2000; Hall *et al.*, 2006). Because all of these factors are affected by fire severity, we would expect higher severity fires to have a greater impact on plant communities than low severity fires in fire-suppressed forest ecosystems, where probability of tree survival is heavily influenced by fire intensity (Oosting, 1944; Safford *et al.*, 2012; Thies and Westlind, 2012), and understory plant growth is largely limited by the density of litter and duff layers (Hodgkins, 1958; Glasgow and Matlack, 2007; Wayman and North, 2007).

Compared to nutrient and vegetation impacts, effects of fire on animal taxa are much more equivocal and unpredictable, likely due to more complex trophic interactions and the ability of some taxa to adapt to habitat changes through both movement and behavioral responses (Geluso and Bragg, 1986; Jones *et al.*, 2004; Engstrom, 2010). Animal-based studies often detect minimal or no effects of prescribed fire (Ford *et al.*, 1999; Greenberg and Waldrop, 2008; Dickson *et al.*, 2009; Greenberg *et al.*, 2010). However, studies assessing impacts of fire severity on animals have found that severity is an important factor affecting population and community dynamics. Smucker *et al.* (2005) reported that some bird species (i.e., hermit thrush [*Catharus guttatus*] and western tanager [*Piranga ludoviciana*]) responded positively to low severity fire and negatively to high severity fire in the same study area, and Roberts *et al.* (2008) found equivalent responses in small mammals. Alternatively, by meta-analysis on bird and small mammal responses to fire severity, Fontaine and Kennedy (2012) concluded that fire severity did not consistently impact species response direction (e.g., a negative or positive impact), but response magnitude of animals increased with fire severity.

The above ensemble of previous fire research suggests that fire severity is an important, and potentially driving, factor in determining fire impacts to essentially all fire-affected ecosystem components (Knapp *et al.*, 2009). Thus, to improve our understanding of the potential of fire for ecosystem management, and to better understand the role of fire severity, there is a need to quantify the impacts of fire severity within ecosystems. Further, because fire impacts to terrestrial components can influence aquatic components, we examined both terrestrial and aquatic ecosystem and community responses in this study. Specifically, we used water quality, aquatic arthropod (data for wildfire impacts only), vegetation (live and dead), and terrestrial ground-dwelling arthropod data collected prior to and following low severity winter prescribed burns, moderate severity summer prescribed burns, moderate severity summer wildfires, and high severity summer wildfires to compare and contrast effects of fire severity on several ecosystem components within a mixed pine/hardwood forest in the south-central USA. Based on previous research, we hypothesized that impacts to terrestrial components would increase with increasing fire severity, and subsequent effects on aquatic components would also be more pronounced as fire severity increased.

In addition to our basic interest concerning fire severity effects, we also have an applied interest specific to the impacts of fire on our study area, the Lost Pines ecoregion of Texas. This ecoregion is the last remaining stronghold for the federally endangered Houston toad, and populations have been declining within the ecoregion for decades, to the point now where the species is at high risk of extinction in the wild (Gaston *et al.*, 2010; Duarte *et al.*, 2011; Brown *et al.*, 2013). Thus, we are interested in fire as a habitat restoration tool in this ecoregion, with particular interest in potential and realized effects

on this endangered species (Brown *et al.*, 2011; Brown *et al.*, 2012). To this end, we included a discussion of results of this study with respect to potential impacts on the Houston toad.

2. Methods

2.1. Study area

This study was conducted in the Lost Pines ecoregion in Bastrop County, Texas, USA. The Lost Pines is a 34,400-ha remnant patch of pine-dominated forest that is thought to have been isolated from the East Texas Piney Woods ecoregion between 10,000 and 14,000 years ago (Bryant, 1977), with the pines of the area beginning to diverge up to 30,000 years ago (Al-Rabah'ah and Williams, 2004). The Lost Pines was extensively logged in the 1800s and early 1900s (Moore, 1977). Since the early to mid-1900s broad-scale fire suppression has been implemented throughout the ecoregion, resulting in the accumulation of heavy fuel loads.

The study area for this project was the 1,948-ha Griffith League Ranch (GLR). The GLR is primarily a forested ranch with an overstory dominated by loblolly pine (*Pinus taeda*), post oak (*Quercus stellata*), and eastern red cedar (*Juniperus virginiana*), and a pre-burn understory dominated by yaupon holly (*Ilex vomitoria*), American beautyberry (*Callicarpa Americana*), and farkleberry (*Vaccinium arboreum*). The property is underlain by deep sandy soils of the Patilo-Demona-Silstid Association (Baker *et al.*, 1979). The GLR contains 3 permanent ponds (i.e., ponds have not dried in at least 12 years), 10 semi-permanent ponds (i.e., ponds typically dry several times per

decade), and 10 or more ephemeral pools that hold water for days to months annually depending on rainfall.

2.2. Fires

2.2.1. Prescribed burns

We conducted prescribed burns on the GLR on 13 November 2009, 10 January 2010, and 7 August 2010, with the prescribed burn areas encompassing ca. 21 ha, 95 ha, and 262 ha, respectively. The habitat management goal of the prescribed burns was fuel load reduction, and burn intensities were low to reduce the potential for the fires to spread beyond the designated burn units, and to reduce the probability of crowning (i.e., aerial fire in the forest canopy).

2.2.2. Wildfires

Two moderate severity wildfires occurred on the GLR on 21 August 2010, which burned 36 ha and 153 ha, respectively. These fires were started from embers in the 7 August 2010 prescribed burn unit that were wind-thrown. Burn breaks were installed during the fires to restrict their spread. On 4 September 2011 a high severity wildfire began from multiple initial fire outbreaks across the Lost Pines. The fire was unstoppable due to wind gusts in excess of 58 kph resulting from the passage of tropical storm Lee, coupled with extreme drought conditions in central Texas (Lost Pines Recovery Team, 2011). After 18 days the fire was 95% contained, with the total burn area encompassing 13,406 ha. A fire break was installed on the GLR during the burn, restricting the fire on

the property to 987 ha. On 11 October 2011 the wildfire breached a fire break on the GLR, burning an additional 125 ha (Figure 4.1).




Figure 4.1. Aerial image of the Griffith League Ranch (GLR), Bastrop County, Texas, USA, and its location with respect to a 13,406 ha wildfire that occurred in the Lost Pines ecoregion in September 2011, with a breach of the fire break resulting in an additional 125 ha burned in October 2011. Overlain on the image are the locations of all fires included in this study, and the locations of the ponds, terrestrial habitat plots, and drift fence arrays used for this study assessing the impacts of fire severity on aquatic and terrestrial ecosystem components.

2.3. Data collection

2.3.1. Water quality

We assessed water quality opportunistically at 16 ponds on the GLR between 17 April 2009 and 4 February 2012, with sampling frequency highest shortly before and following fires to capture immediate impacts. We sampled the study area a total of 32 times, with each pond sampled between 3 and 25 times (median = 15). Days between sampling ranged from 2 to 150 (mean = 33). Whether or not a given pond was sampled usually depended on whether or not the pond held water at the time of sampling. However, a subset of 5 ponds were sampled monthly in 2009 for an independent study (Gaertner et al. 2012), and ponds located in burned units were sampled more frequently immediately following all burn events (i.e., 1 to 3 times within the first 2 weeks). We sampled water within 1 m of pond edges using 1 L Nalgene[®] collection bottles. Within 24 hours of collection we estimated pH using a SympHony 5B70P pH meter, filtered pond water through Gelman A/E glass-fiber filters (1- μ m pore size), and preserved water samples with sulfuric acid. We extracted chlorophyll *a* (Chl-*a*) from filters with acetone, and analyzed Chl-*a* using a Turner Designs Trilogy fluorometer. We quantified total suspended solids (TSS) in the water column by filtering water through pre-combusted and pre-weighed A/E filters and then drying filters at 60 °C for 48 to 72 hours, and re-weighing the filters. Quantification of non-volatile suspended solids (NVSS) was determined by subsequently combusting filters at 550 °C for four hours and then taking weights (Heiri *et al.*, 2001). We obtained filter mass values using a Mettler Toledo MX5 microbalance.

We used a Varian Cary 50 Ultraviolet-Visible light spectrophotometer for the remaining water quality analyses. We measured soluble reactive phosphorus (SRP) and total phosphorus (TP) using the molybdenum blue method (Wetzel and Likens, 2000). To estimate TP we digested unfiltered samples with potassium persulfate then quantified SRP. We measured nitrate (NO_3^-) and total nitrogen (TN) using second-derivative UV spectroscopy (Crompton *et al.*, 1992). To estimate TN we digested unfiltered samples with alkaline potassium persulfate then quantified NO_3^- . We analyzed ammonium (NH_4^+) using the phenol-hypochlorite method (Wetzel and Likens, 2000). For all water quality analyses we collected two water samples per pond on each sampling date, and used the average of the replicates for this study.

In addition to measuring water quality variables, we also estimated pond depth and percent canopy cover around each pond for use as covariables in water quality analyses. We estimated pond depth using permanent staff gauges located at the deepest point of each pond. We estimated percent canopy cover around each pond using a spherical densiometer at two to six randomly selected points at the pond edge, with higher numbers of estimation points corresponding to ponds of larger size. We averaged the estimates at each pond per sampling date.

The terrestrial habitat surrounding 4 ponds was burned during the November 2009 and January 2010 prescribed fires, 2 ponds during the August 2010 prescribed fire, 2 ponds during the August 2010 wildfires (including 1 previously burned pond from the 13 November 2009 prescribed fire), and 7 ponds during the September 2011 wildfire (including 6 previously burned ponds). The remaining 8 ponds served as unburned ‘controls’ in this study. However, we note that these ponds were not necessarily

completely free of fire impacts, as volatilized nitrogen and carbon could have been deposited in control ponds through smoke and ash deposition (Battle and Golladay, 2003; Earl and Blinn, 2003).

2.3.2. Aquatic arthropods

Unfortunately, sampling of aquatic arthropods in ponds was initiated after the low and moderate severity fires and most of the sampled ponds were not part of those burn treatments. Thus, we were only able to assess impacts of the high severity fire on the composition of pond arthropods. The data set included 4 control (unburned) ponds and 4 wildfire ponds, with 3 pre-burn sampling events and 4 post-burn sampling events. We sampled aquatic arthropods seasonally, collecting 7 samples at 8 ponds between August 2010 and January 2013 using a dip net (900 μm aperture netting). For each sampling event at each pond we sampled 3 points ca. 1 m from the perimeter, maintaining approximately even spacing between points, and performed 3 successive dip net sweeps per point. We combined the samples obtained at each pond and stored them in 95% ethanol. We identified insects to family, with the exception of Ephemeroptera (mayflies) and Odonata (dragonflies and damselflies), which we identified to order, and we identified other arthropods to class or order. Further, we removed larval forms from the data set with the exception of Chironomidae (non-biting midges), Culicidae (mosquitos), Ephemeroptera, Odonata, and Trichoptera (caddisflies), for which all captures were larvae. The resulting data set contained 26 aquatic arthropod groups.

2.3.3. Terrestrial vegetation

We used National Park Service (2003) fire monitoring guidelines to assess fire severity and impacts of the fires on forest structure characteristics and community composition of understory vegetation. We randomly placed thirty-one 20 m x 50 m plots in forested habitat, 29 of which were used in this study. The remaining 2 plots were abandoned after fire breaks were accidentally installed within them during wildfires. Only 1 plot was burned in the 2010 moderate severity wildfires, and this plot was used only in fire severity analyses. In addition, because no plots were burned in the November 2009 low severity prescribed fire, we installed 1 temporary plot adjacent to a permanent plot to assess fire severity. Two plots were burned during the January 2010 low severity prescribed fire, 4 plots during the August 2010 moderate severity prescribed fire, 1 plot during the August 2010 moderate severity wildfire, and 17 plots during the September and October 2011 high severity wildfires (including 4 plots that had been burned previously).

We surveyed vegetation plots throughout summer and fall of 2008, 2009, 2010, and 2012, and assessed fire severity at all plots within 66 days of a fire. In addition, we surveyed burned plots in 2011 to assess tree mortality from the low and moderate severity fires. We quantified the following variables in all plots: percent canopy cover, overstory ($\text{DBH} \geq 15 \text{ cm}$) and pole-sized ($\text{DBH} \geq 2.5 \text{ cm}$ and $< 15 \text{ cm}$) tree mortality by species, seedling tree ($\text{DBH} < 2.5 \text{ cm}$) abundance, understory vegetation species cover (i.e., number of transect points intersecting vegetation), shrub and herbaceous vegetation abundance (i.e., number of stems detected in plots), litter and duff depth (mm), and fuel abundance (1 hour [0 to 0.62 cm diameter], 10 hour [0.63 to 2.54 cm diameter], 100 hour

[2.55 to 7.62 cm diameter], and 1000 hour [>7.62 cm diameter]). We identified shrubs, vines, and forbs to genus or, typically, to species. We identified the 2 most common grasses in the study area to genus (i.e., *Dicanthelium* and *Eragrostis*), with the remaining grasses grouped into the family Poaceae for analyses. To assess fire severity to substrate within each burned plot we used four 15 m transect lines, each consisting of 4 points spaced 5 m apart. We assigned points a burn severity ranking from 0 (unburned) to 4 (heavily burned), with the same researcher performing all assessments to ensure consistency in ranks. Overstory canopy cover is not included in the National Park Service (2003) fire monitoring guidelines, but we estimated percent canopy cover using a spherical densiometer at the 4 corners and center of each plot, then computed the mean of those 5 estimates for our plot estimate each survey year.

2.3.4. Ground-dwelling arthropods

We sampled ground-dwelling arthropods 24 times between 8 March 2009 and 21 April 2012 using 18 Y-shaped and 7 linear drift fence arrays. Y-shaped arrays consisted of three 15 m arms with a 19 L center bucket and a 19 L bucket at each arm terminus. Linear arrays consisted of a 15 m arm with a 19 L bucket at each end. We sampled at least once in spring, summer, and fall annually during the sampling period, with additional sampling events occurring during the spring months and following the burns. For each sampling event we allowed pitfall traps to collect arthropods for 7 days prior to collection. We euthanized arthropods through freezing, sorted arthropod captures by drift fence array, and identified insects to family and other arthropods to order. We did not include larvae with the exception of Myrmeleontidae (antlions), and we removed captures

of primarily flying arthropods (i.e., Diptera [flies], Cicadidae [cicadas], Aculeata [wasps], Lepidoptera [moths], and Odonata [dragonflies]) because our sampling design was likely inadequate for estimating relative abundance differences for flying taxa. We also did not include Formicidae (ant) captures due to their ability to easily crawl into and out of pitfall traps. The resulting data set contained 49 ground-dwelling arthropod groups.

The terrestrial habitat surrounding 4 drift fence arrays was burned during the November 2009 low severity prescribed fire, 4 drift fence arrays during the August 2010 moderate severity prescribed fire, and 14 drift fence arrays during the September and October 2011 high severity wildfires (including all 8 drift fence arrays that had burned previously). The remaining 10 drift fence arrays served as controls in this study.

2.4. Data Analyses

2.4.1. Water quality

We assessed impacts of the fires to pond water quality using Principal Response Curves (PRC), an extension of Redundancy Analysis (RDA; Van den Brink and Ter Braak, 1999). This multivariate analysis method is designed to assess treatment effects over time, and allows both an overall community response and specific response variables to be assessed through comparisons to control sites (Lepš and Šmilauer, 2003). The significance of treatment x time interactions is assessed using Monte Carlo permutation tests, where sites are randomized within, but not across sampling periods (p -values in the results represent permutation tests for the first canonical axis).

Of the 16 ponds monitored throughout the study period, 8 were subjected to fire, with 6 subjected to multiple fire events. Because the fires occurred at different times over the study period, we conducted 3 separate analyses, assessing impacts of low (i.e., winter 2010 and 2011 prescribed burns), moderate (i.e., summer 2010 prescribed burn and wildfires), and high (i.e., summer 2011 wildfires) severity fire. To minimize the influence of fires not included in a given analysis, we removed ponds from the data set (i.e., sampling events) after they were burned in another fire. Ponds that were burned multiple times re-entered the data set after the fire of interest. Thus, the pre-burn samples were truly pre-burn in all analyses. In addition, for the low and moderate severity fire analyses, we did not include any sampling events after the 2011 wildfires, given that all but 1 of the ponds re-burned in those wildfires.

For all 3 analyses we included sampling period, pond depth, and canopy cover at the pond edge as covariates. Including sampling event allowed us to test for a treatment x time interaction. The data set included several missing values (i.e., 20 of 2710 observations), and we estimated those values using the mean for a given variable in a pond across all sampling events. We centered and standardized the response data because variables were measured on different scales (i.e., response variables had a zero average and unit variance). In addition, we $\log_{10}$ transformed the response data so that percentage rather than absolute changes in captures were analyzed. We performed these analyses using the program CANOCO (version 4.5).

2.4.2. Aquatic arthropods

We assessed wildfire impacts on both total number of captured individuals and community composition of aquatic arthropods. To assess wildfire impacts on total captures we used generalized least squares analyses, with the capture data transformed using the arcsinh (i.e., inverse hyperbolic sine) transformation to satisfy assumptions of normality and homoscedasticity (Fowler *et al.*, 1998). We accounted for non-independence in our repeated measures data using a continuous autoregressive term (corCAR1) nested within each pond (i.e., time series; Zuur *et al.*, 2009). We then tested for an interaction between treatment (control or fire) and burn status (pre-burn or post-burn). We performed this analyses using the program R (version 2.14).

To assess impacts of the high severity wildfire on community composition we used redundancy analysis (RDA), which is an extension of principal components analysis (PCA), but includes explanatory variables. We chose RDA over canonical correspondence analysis (CCA) because our gradient lengths were short (< 4) and our predictors were categorical (Lepš and Šmilauer, 2003). To assess whether or not the wildfire impacted community composition we tested for a treatment x burn status interaction using a Monte Carlo permutation test (p -value in the results represents a permutation test for the first canonical axis). We included pond identity as a covariate. By including this covariate we subtracted the average number of captures and assessed only capture changes within each pond (Lepš and Šmilauer, 2003). We $\log_{10}$ transformed the response data so that percentage rather than absolute changes in captures were analyzed. We performed this analysis using the program CANOCO (version 4.5).

2.4.3. Terrestrial vegetation

We summarized differences in burn severity with respect to substrate and tree mortality using a visual descriptive statistics approach. We calculated the mean, median, and range of fire severity ranks in the low, moderate, and high severity fire vegetation plots. Further, we categorized the high severity plots by whether they were control plots prior to the 2011 wildfire, or had been burned in the prior winter or summer fires. This allowed us to assess whether or not prior burning reduced severity of the 2011 wildfire. To compare impacts on tree mortality, we separated trees into 3 genus-level groups: pine (loblolly pine), oak (post oak, blackjack oak [*Quercus marilandica*], and water oak [*Quercus nigra*]), and cedar (eastern red cedar), and plotted the proportion of overstory and pole-sized trees that died within 1 year following the fires for each group.

To assess impacts of the fires to forest structure characteristics and community composition of understory vegetation, we used RDA. For forest structure characteristics, we included 12 response variables: 1 hour fuel, 10 hour fuel, 100 hour fuel, 1000 hour fuel, total above-ground litter depth, total above-ground duff depth, overstory canopy cover, understory species cover, vegetation species richness, cedar seedling abundance, oak seedling abundance, and pine seedling abundance. Our understory species cover metric was the number of points along a transect line (i.e., 166 total points, with points located every 0.3 m along a 50 m transect) intersected by vegetation. Our species richness metric included all vegetation groups (see section 2.3.2) encountered from the tree, shrub and vine, and herbaceous vegetation sub-plots, and the understory species cover transect.

To assess whether or not the fires impacted forest structure characteristics and understory community composition, we tested for treatment x burn status interactions

using a Monte Carlo permutation test (p -values in the results represent permutation tests for the first canonical axis). We included vegetation plot identity and cumulative precipitation between January and May (i.e., prior to initiation of sampling) for each sample year as covariates. By including plot identity as a covariate, we subtracted the average response variable values and assessed only changes in response variable values within each plot (Lepš and Šmilauer, 2003).

For the forest structure characteristics analyses, we centered and standardized the response data because variables were measured on different scales. In addition, we $\log_{10}$ transformed the response data so that percentage rather than absolute changes in captures were analyzed. For the understory community composition analyses, we centered and $\log_{10}$ transformed the response data. We removed 10 shrub and 41 herbaceous vegetation records because we were unable to identify the plants. In addition, we removed dead individuals and species with less than 5 total observations (shrubs: $n = 5$; forbs: $n = 23$), resulting in 28 vegetation groups.

We performed additional univariate analyses for forest structure characteristics and species that appeared to display a strong response in the high severity wildfire analyses, including fuel (all classes), litter depth, duff depth, overstory canopy cover, understory species cover, and vegetation species richness, and the species yaupon holly, flowering spurge (*Euphorbia corollata*), pokeweed (*Phytolacca americana*), horseweed (*Conyza canadensis*), sedges (*Cyperus* spp.), and panic grasses (*Dicanthelium* spp.). For these analyses we used a generalized least squares approach, with the data transformed using the arcsinh (i.e., inverse hyperbolic sine) transformation to satisfy assumptions of normality and homoscedasticity (Fowler *et al.*, 1998). We included cumulative

precipitation between January and May (i.e., prior to initiation of sampling) for each sample year as a covariate in the community composition analyses, and accounted for non-independence in our repeated measures data using a continuous autoregressive term (corCAR1) nested within each plot (i.e., time series; Zuur *et al.*, 2009). We then tested for an interaction between treatment (control or fire) and burn status (pre-burn or post-burn). We performed these analyses using the program R (version 2.14).

The winter prescribed burn data sets included two pre-burn sampling years (2008 and 2009) and one post-burn sampling year (2010), with 2 treatment plots and 26 control plots. The summer prescribed burn data sets included 3 pre-burn sampling years (2008 to 2010) and 1 post-burn sampling year (2012). In addition, plots burned in the remaining fires were removed, as they did not represent true controls, resulting in 3 treatment plots and 9 control plots. The high severity wildfire data set included 3 pre-burn sampling years (2008 to 2010) and 1 post-burn sampling year (2012). For the wildfire analyses, we separated plots into 3 categories, those that were controls prior to the wildfire ($n = 13$), those that were within the low severity winter prescribed burn zone ($n = 2$), and those that were within the moderate severity summer prescribed burn zone ($n = 1$). Three summer prescribed burn plots were not re-burned in the wildfire and were removed from this data set, and the remaining 9 plots were used as controls. We performed these analyses using the program CANOCO (version 4.5).

2.4.4. Ground-dwelling arthropods

We assessed impacts of the fires on both total number of captured individuals and community composition of ground-dwelling arthropods. In addition, we performed

univariate analyses for species groups that appeared to show a strong response in the high severity wildfire analysis. These groups included Araneae (spiders), Carabidae (ground beetles), Curculionidae (snout beetles), Diplopoda (millipedes), Gryllacrididae (raspy crickets), Scarabaeidae (scarab beetles), and Scorpiones (scorpions). To assess fire impacts on total captures and individual taxonomic groups we used generalized least squares analyses, with the capture data transformed using the arcsinh (i.e., inverse hyperbolic sine) transformation to satisfy assumptions of normality and homoscedasticity (Fowler *et al.*, 1998). For all analyses, we included sampling season and cumulative precipitation during the 7-day sampling period as covariates, and accounted for non-independence in our repeated measures data using a continuous autoregressive term (corCAR1) nested within each array (i.e., time series; Zuur *et al.*, 2009). We then tested for an interaction between treatment (control or fire) and burn status (pre-burn or post-burn).

The winter and summer prescribed burn analyses included all sampling periods prior to the September 2011 wildfire, and each analysis excluded the arrays within the burn zone of the other fire, as those arrays were not true controls. For the wildfire analysis, we separated arrays into 3 categories: (1) those that were controls prior to the wildfire, (2) those that were within the low severity winter prescribed burn zone, and (3) those that were within the moderate severity summer prescribed burn zone. Because the same response data were used in multiple analyses for the total capture analyses, we adjusted the level considered significant from the standard $\alpha = 0.05$ using the Bonferroni correction (i.e., $\alpha = 0.017$). We performed these analyses using the program R (version 2.14).

To assess impacts of the fires to community composition, we used RDA. To assess whether or not the fires impacted community composition, we tested for treatment x burn status interactions using a Monte Carlo permutation test (p -values in the results represent permutation tests for the first canonical axis). We included cumulative precipitation during each sampling period, sampling season, and arrays as covariates. By including arrays as covariates, we subtracted the average number of captures and assessed only capture changes within each array (Lepš and Šmilauer, 2003). We $\log_{10}$ transformed the response data so that percentage rather than absolute changes in captures were analyzed. As with the total abundance analyses, the winter and summer prescribed burn data sets included all sampling periods prior to the September 2011 wildfire, and each analysis excluded the arrays within the burn zone of the other fire; whereas, for the wildfire analysis we separated arrays into 3 categories, those that were controls prior to the wildfire, those that were within the low severity winter prescribed burn zone, and those that were within the moderate severity summer prescribed burn zone. We performed these analyses using the program CANOCO (version 4.5).

3. Results

3.1. Water quality

The PRC analyses indicated a significant treatment x time interaction for the low-severity, winter-fire ponds ($P = 0.032$) and the high-severity, summer-fire ponds ($P = 0.024$), but not for the moderate-severity, summer-fire ponds ($P = 0.60$), with the first axis explaining 5%, 5%, and 4% of the variance, respectively. The PRC diagrams

indicated increased pond nutrient levels in all burns, with corresponding increases in chl-*a* concentrations (Figure 4.2). Impacts on pH and DOC were equivocal, with pH decreasing following summer fires, but increasing slightly after winter fires, and DOC increasing following the low and moderate severity fires, but decreasing after the high severity fire.




Figure 4.2. Results from Principal Response Curve (PRC) analyses used to assess the impacts of low (A), moderate (B), and high (C) severity fires on pond water quality on the Griffith League Ranch (GLR), Bastrop County, Texas, USA. Arrows indicate the first post-fire sampling event following significant precipitation (cm). Number of days before and after fires shown on the X-axes, and PRC scores shown on the left Y-axes. The right Y-axes allow the estimation of percentage differences between treatments for individual variables at any point in time. This is accomplished through the following equation:

$$\text{Difference (\%)} = 100 \times e^{(\text{PRC1 score at time point [left Y-axis]} \times \text{variable score [right Y-axis]})}$$

The magnitude of impacts increased with fire severity, but water quality began to return to baseline levels within 200 days for all burn severities.

3.2. Aquatic arthropods

We did not detect a high severity wildfire impact on total number of captured individuals of aquatic arthropods ($F_{1,52} = 0.71$, $P = 0.404$). Likewise, the RDA analysis indicated the high severity wildfire had no impact on community composition of aquatic arthropods ($P = 0.949$), with 3.2% of the variation explained by the model. In addition, the RDA biplot confirmed the statistical test, with all aquatic arthropod groups located either near the origin or at nearly orthogonal to the treatment x burn status predictor.

3.3. Terrestrial vegetation

3.3.1. Fire severity

The distribution of fire severity ranks to substrate ranged from 0.5 to 1.44 for the low severity winter fires, 0.5 and 2.5 for the moderate severity summer fires, and 1.68 to 4 for the high severity summer fires when plots were controls prior to the fires (Figure 4.3). Our plots indicated that burn severity for the 2011 wildfire was not affected by the prior low severity winter burn, but was dramatically reduced in the prior moderate severity summer burn zone (Figure 4.4).



Figure 4.3. Examples of vegetation plots following low (A), moderate (B), and high (C) severity fires (left panels), and ca. 1 year following the fires (right panel), on the Griffith League Ranch (GLR), Bastrop County, Texas, USA. Low and moderate severity fires did not significantly impact forest structure characteristics or species composition of understory vegetation, whereas high severity fires reduced overstory canopy cover, litter and duff depth, and the amount of fuel in all fuel classes, and increased diversity and cover of understory vegetation.




Figure 4.4. Visual descriptive statistics displaying fire severity to substrate on the Griffith League Ranch (GLR), Bastrop County, Texas, USA for low, moderate, and high severity fires. We assessed fire severity using four 15 m transect lines, each consisting of 4 points spaced 5 m apart, and assigned points a burn severity ranking from 0 (unburned) to 4 (heavily burned). We used the mean severity ranking of the 4 transects in this analysis. The high severity summer fires included 3 categories: (1) plots that were controls prior to the wildfire [High (control)], (2) plots that were part of the prior winter low severity fires [High (winter)], and (3) plots that were part of the prior summer moderate severity fires [High (summer)]. Boxes enclose the range, diamonds show the mean, and horizontal bars delineate the median fire severity rank for each category. Stars in the low and moderate severity categories denote the mean severity rank when unburned points were removed. Fire severity rank increased with fire severity, and prior moderate severity fires reduced substrate burn severity from the high severity fires.

Both overstory and pole-sized tree mortality was substantially greater in the high severity fire plots compared to the low and moderate severity fire plots (Figure 4.5).

Similar to substrate impacts, our plots indicated that tree mortality following the high severity summer fires was not reduced in the low severity winter burn zone, but was dramatically reduced in the moderate severity summer burn zone, with the exception of overstory cedar trees. For all of the fires, pine tree mortality was greater than either oak or cedar tree mortality.




Figure 4.5. Overstory (A) and pole-sized (B) tree mortality on the Griffith League Ranch (GLR), Bastrop County, Texas, USA following low, moderate, and high severity fires. The high severity summer fires included 3 categories: (1) plots that were controls prior to the wildfire [High (control)], (2) plots that were part of the prior winter low intensity fires [High (winter)], and (3) plots that were part of the prior summer moderate intensity fires [High (summer)]. We separated trees into 3 genus-level groups: pine (loblolly pine), oak (post oak, blackjack oak [*Quercus marilandica*], and water oak [*Quercus nigra*]), and cedar (eastern red cedar), and plotted the proportion of overstory and pole-sized trees that died within 1 year following the fires for each group. Tree mortality increased with fire severity, and prior moderate severity fires reduced tree mortality caused by the high severity fires. Note there were no overstory cedar trees in the low severity winter plots.

3.3.2. Forest structure characteristics and understory community composition

The RDA analyses indicated the low severity winter fires had no significant impact on forest structure characteristics ($P = 0.133$) or understory community composition ($P = 0.145$), the moderate severity summer fires had no significant impact on forest structure characteristics ($P = 0.102$) or understory community composition ($P = 0.082$), and the high severity summer fires did impact forest structure characteristics ($P = 0.002$) and understory community composition ($P = 0.002$). The amount of total variation explained by the high severity fire models was 18.1% and 9.6% for forest structure characteristics and community composition of understory vegetation, respectively.

The RDA biplot was congruent with significance tests (Figure 4.6). No response variables were strongly associated with the low and moderate severity fires, as indicated by the direction and length of response arrows. For plots that were controls prior to the high severity wildfires, overstory canopy cover ($F_{1,96} = 32.49$, $P < 0.001$), litter depth ($F_{1,96} = 66.63$, $P < 0.001$), duff depth ($F_{1,96} = 62.18$, $P < 0.001$), and amount of fuel in all fuel classes ($F_{1,96} = 24.55$, $P < 0.001$ [1 hour fuel]; $F_{1,96} = 7.00$, $P = 0.010$ [10 hour fuel]; $F_{1,96} = 11.49$, $P = 0.001$ [100 hour fuel]; $F_{1,96} = 14.70$, $P < 0.001$ [1000 hour fuel]) was reduced ca. 1 year after the fire, whereas species richness ($F_{1,96} = 4.08$, $P = 0.046$) was higher ca. one year after the fire. Understory vegetation cover was not significantly different ($F_{1,96} = 0.61$, $P = 0.44$).

For the vegetation taxa with a clear response to the wildfire, all responses were positive with the exception of yaupon holly. The univariate analyses supported the positive responses of pokeweed ($F_{1,95} = 87.61$, $P < 0.001$), panic grasses ($F_{1,95} = 5.51$, $P = 0.021$), and sedges ($F_{1,95} = 33.59$, $P < 0.001$), and the negative response of yaupon

holly ($F_{1,95} = 45.31$, $P < 0.001$), to high severity wildfire. The wildfire analyses indicated that prior fires influenced the outcome for understory community composition.

Abundance of flowering spurge ($F_{1,95} = 3.81$, $P = 0.054$) appeared to be positively associated with plots previously burned with low severity winter fires, whereas abundance of horseweed ($F_{1,95} = 3.23$, $P = 0.075$) appeared to be negatively associated with these plots. Plots previously burned with moderate severity summer fires indicated the opposite associations ($F_{1,95} = 3.42$, $P = 0.068$ and $F_{1,95} = 27.97$, $P < 0.001$, respectively).




Figure 4.6. Results from Redundancy Analyses (RDA) used to assess the impacts of low severity winter fires (A), moderate severity summer fires (B), and high severity summer fires (C) on forest structure characteristics (left biplot) and species composition of understory vegetation (right biplot) on the Griffith League Ranch (GLR), Bastrop County, Texas, USA. The high severity summer fires included 3 categories: (1) plots that were controls prior to the wildfire, (2) plots that were part of the prior winter low intensity fires, and (3) plots that were part of the prior summer moderate intensity fires. Dashed lines represent treatment-burn status interactions and solid lines represent response variable associations with those interactions. Of the 28 vegetation groups included in the analyses, only the 8 to 11 showing the greatest interaction response were included for ease of interpretation. The analyses indicated that low and moderate severity fires had weak effects on forest structure characteristics and species composition of understory vegetation. In contrast, high severity fires dramatically reduced fine and large fuel loads and increased diversity and cover of understory vegetation.

3.4. Ground-dwelling arthropods

We found no significant effect of fire on total captures of ground-dwelling arthropods in any of the fire severity analyses, as indicated by no significant treatment x burn status interaction effects (Table 4.1). The RDA analyses indicated the low-severity winter prescribed burn had no impact on community composition ($P = 0.110$); whereas, the moderate-severity summer prescribed burn ($P = 0.024$) and high-severity wildfires ($P = 0.002$) did affect community composition. However, the amount of total variation explained by the low, moderate, and high severity models was low; 0.8%, 1.0%, and 2.9%, respectively. The RDA biplot indicated little or no response to fires for most arthropod groups (Figure 4.7). However, the univariate analyses indicated several groups responded to fire. For areas that were controls prior to the high-severity wildfire, Curculionidae ($F_{1,594} = 44.29$, $P < 0.001$) and Gryllacrididae ($F_{1,594} = 6.10$, $P = 0.014$) responded positively, and Araneae responded negatively ($F_{1,594} = 6.56$, $P = 0.011$). The positive response of Scarabaeidae to areas burned in both the summer prescribed fire and the high-severity wildfire was trending towards significance ($F_{1,594} = 3.17$, $P = 0.076$). Although the RDA biplot indicated that Diplopoda responded negatively to fire in general, that response was not strongly associated with a particular category and was not found to be significant for areas that were controls prior to the wildfire ($F_{1,594} = 0.54$, $P = 0.464$), or areas that were part of the prior summer prescribed burn ($F_{1,594} = 1.32$, $P = 0.252$). Although Carabidae and Scorpiones appeared to show a moderate positive and negative response in areas that were controls prior to the wildfire, respectively, neither relationship was supported from univariate analyses ($F_{1,594} = 0.01$, $P = 0.943$ and $F_{1,594} = 1.42$, $P = 0.233$, respectively).

Table 4.1. Results of generalized least squares analyses used to assess if a low severity winter prescribed burn, a moderate severity summer prescribed burn, and a high severity summer wildfire impacted total captures of ground-dwelling arthropods on the Griffith League Ranch (GLR), Bastrop County, Texas, USA, with sampling season and cumulative precipitation during the sampling period included as covariates. We sampled arthropods using 24 drift fence arrays with pitfall traps 24 times between 8 March 2009 and 21 April 2012. The low severity winter prescribed burn, moderate severity summer prescribed burn, and high severity wildfire occurred on 13 November 2009, 7 August 2010, and 4 September 2011, respectively. The high severity summer fires included 3 categories: (1) plots that were controls prior to the wildfire (control), (2) plots that were part of the prior winter low intensity fires (winter), and (3) plots that were part of the prior summer moderate intensity fires (summer). The lack of significant treatment x burn status interaction effects indicated the fires did not significantly impact total abundance of ground-dwelling arthropods.

Fire	Coefficient ($\pm$ SE)	F-value	df	P
<i>Prescribed (winter)</i>				
Season	0.192 (0.07)	6.26	1,393	0.013
Precipitation	0.003 (0.002)	1.85	1,393	0.175
Treatment	0.426 (0.44)	2.28	1,393	0.132
Status	-0.081 (0.21)	1.60	1,393	0.207
Treatment*status	-0.807 (0.46)	3.06	1,393	0.081
<i>Prescribed (summer)</i>				
Season	0.114 (0.07)	3.07	1,393	0.080
Precipitation	0.004 (0.00)	5.52	1,393	0.019
Treatment	0.447 (0.28)	2.44	1,393	0.119
Status	0.228 (0.19)	1.14	1,393	0.286
Treatment*status	-0.206 (0.41)	0.26	1,393	0.613
<i>Wildfire</i>				
Season	0.101 (0.05)	2.87	1,590	0.091
Precipitation	0.007 (0.00)	16.15	1,590	< 0.001
Treatment (winter)	-0.154 (0.23)	2.45	1,590	0.118
Treatment (summer)	0.529 (0.23)	2.01	1,590	0.156
Treatment (control)	0.368 (0.19)	2.55	1,590	0.111
Status	-0.109 (0.23)	7.65	1,590	0.006
Treatment*status (winter)	-0.148 (0.42)	0.00	1,590	0.944
Treatment*status (summer)	-0.914 (0.42)	3.66	1,590	0.056
Treatment*status (control)	-0.427 (0.35)	1.48	1,590	0.224




Figure 4.7. Results from Redundancy Analyses (RDA) used to assess the impacts of a low severity winter fire (A), moderate severity summer fires (B), and high severity summer fires (C) on ground-dwelling arthropods in the Lost Pines ecoregion of Texas, USA. We sampled arthropods using drift fence arrays with pitfall traps. The high severity summer fires included 3 categories: (1) traps that were controls prior to the wildfire, (2) traps that were part of the prior winter prescribed burn, and (3) traps that were part of the prior summer prescribed burn. Dashed lines represent treatment-burn status interactions and solid lines represent arthropod group associations with those interactions. Of the 49 arthropod groups included in the analyses, only the 9 to 15 showing the greatest interaction response were included for ease of interpretation. The analyses indicated low and moderate severity fires had no detectable effects on species composition of ground-dwelling arthropods, whereas several arthropod groups responded to the high severity fires.

4. Discussion

The results of our study indicated fire severity was an important factor influencing responses of the ecosystem components we assessed, and directly contributes to the growing literature supporting severity as a critical determinant of fire effects on ecosystems (Schwilk *et al.*, 2006; Wayman and North, 2007; Knapp *et al.*, 2009). Further, based on the data used in this study and our qualitative observations in the study area, moderate severity prescribed burning was effective for reducing wildfire severity, and consequently tree mortality was much lower, litter and duff was reduced but not eliminated, and understory taxonomic diversity increased slightly. In contrast, the low severity fires did little to mitigate tree mortality or substrate burn severity. Overall, our study supports conclusions of the majority of studies, which concluded fuel reduction helps reduce fire severity in pine-dominated forests (Schoennagel *et al.*, 2004; Mitchell *et al.*, 2009; Safford *et al.*, 2012).

Interestingly, we found pine mortality was greater than oak or cedar at all fire severities. This result would appear to contradict the general consensus that pines in the southern United States are not only fire-adapted (Oosting, 1944; Moore *et al.*, 1999; Stambaugh *et al.*, 2011), but fire is necessary for long-term persistence of pine forests (Hartnett and Krofta, 1989; Waldrop *et al.*, 1992; Schulte and Mladenoff, 2005). We believe there are likely several interacting factors in our study responsible for this finding. The first is geographical; the Lost Pines loblolly pine community represents the westernmost edge of the loblolly pine distribution in the U.S. (Al-Rabah'ah and Williams, 2004). The climate in this region is drier than in the expansive East Texas Pineywoods ecoregion (Owen 1989), and this disjunct population has managed to persist under what

are likely suboptimal environmental conditions. As a result, the Lost Pines population may be more vulnerable to climate change and related disturbance (e.g., increased prevalence of wildfires) impacts. Higher vulnerability to climate change and related disturbance impacts at distribution edges, particularly dry edges, appears to be common for conifers (Galiano *et al.*, 2010; Littell *et al.*, 2010; Coops and Waring, 2011; Benavides *et al.*, 2013). Secondly, and related to this, was the timing of the fires. The 2009 and 2010 low and moderate severity fires occurred following exceptional drought conditions (based on the Palmer Drought Severity Index) throughout most of 2009. This drought killed 5.7% of the overstory loblolly pine trees, and 12.0% of the post oak trees, in our vegetation plots. In contrast, eastern red cedar appeared to be very drought resistant, with 0% overstory tree mortality. Thus, the remaining loblolly pines were likely stressed, and thus more vulnerable to fire mortality, than if the burns occurred following several average rainfall or wetter than average years. Third, the Lost Pines loblolly pine population currently has a pine engraver beetle (*Ips* spp.) infestation (KES Consulting *et al.*, 2007). Pine engraver and bark beetles are known to select drought-stressed trees, and further increase vulnerability to mortality (Grosman and Upton, 2006; Schwilk *et al.*, 2006). Finally, fire intensities were potentially higher at the base of loblolly pines compared to the hardwoods due to higher pine needle fuel load densities, which are typically more flammable than hardwood leaves due to lower moisture and higher resin content (Hély *et al.*, 2000; Nowacki and Abrams, 2008). Based on haphazard sampling (3 independent samples per tree species) conducted in December 2008 to gauge differences in litter moisture content, it appeared that loblolly pine had the lowest litter moisture

content (mean = 8.7%, SD = 1.6%), followed by post oak (mean = 8.9%, SD = 0.6%) and eastern red cedar (mean = 14.2%, SD = 4.9%).

We found all fire severities resulted in increased nutrient levels in ponds, and the magnitude of those increases were greatest following high severity fire, which agreed with our hypothesis that aquatic impacts would increase as terrestrial impacts increased. However, for all the fire severity analyses, the magnitude of effects was not substantially greater than the highest natural variability observed prior to burning, and further our results indicated that nutrient levels began returning to baseline levels within 200 days following the fires. The observed increases were clearly influenced by precipitation-induced runoff following the fires, which is both intuitive and congruent with other studies (Gresswell, 1999; Battle and Golladay, 2003). Thus, if fire were to be used as a tool to stimulate aquatic productivity in oligotrophic ponds in similar ecosystems, both fire severity and timing are important considerations. Conversely, ponds and associated drainage zones should be excluded from prescribed burns when temporary nutrient increases are viewed as harmful to aquatic ecosystem integrity.

Despite the impacts of high severity fire on water quality, our analyses indicated those effects did not translate to changes in total captures or community composition of aquatic arthropods. Potential direct mortality of aquatic arthropods would likely have been caused primarily by significant increases in water temperature, which has been documented during high severity fires (Gresswell, 1999; Hitt, 2003; Pilliod *et al.*, 2003). Unfortunately those data were not collected for our study, or as far as we are aware, any water bodies within this wildfire zone, and thus we do not know if the wildfires impacted water temperature. Malison and Baxter (2010) reported a fire severity effect for benthic

stream insects 5 to 10 years post-burn, with greater captures of emergent insects and greater larval biomass in higher severity stream reaches, and several other studies have reported impacts years after fire events (reviewed in Gresswell, 1999). Thus, it is possible the wildfires may affect the aquatic arthropod communities in our study area at a longer time-scale (i.e., lagged responses in population trends). However, given that water quality appeared to be returning to background levels by the end of the study, and all but 3 ponds on our study area dry periodically, we believe long-term impacts are unlikely for aquatic arthropods.

We found that in contrast to high severity fire, low and moderate severity fire in this severely fire-suppressed ecosystem was largely ineffective for enhancing vegetation diversity. This is probably directly related to less fine fuel consumption, and thus less open ground following the fires, as well as higher percent canopy cover due to the majority of trees surviving the fires. We found that even low severity fire was sufficient to top-kill yaupon holly, the pre-burn dominant shrub in our study area. However, this was followed by significant basal sprouting after the low and moderate severity fires (D.J. Brown, personal observation), which agrees with other findings in central Texas (Mitchell *et al.*, 2005). In contrast, the high severity fires not only top-killed, but consumed entirely the majority of yaupon holly individuals, and this species did not reestablish within the hottest areas of the wildfire zones within the time-frame of our study. Rather, pokeweed, a species rarely observed on the property prior to the wildfires, replaced yaupon holly as the dominant shrub in the wildfire zones. This positive fire severity response for pokeweed has been noted elsewhere (Glasgow and Matlack, 2007). In addition, the dominant herbaceous vegetation shifted from panic grasses

(*Dicanthelium* spp.) and flowering spurge (*Euphorbia corollata*), to panic grasses, sedges (*Cyperus* spp.), and horseweed (*Conyza canadensis*), another species rarely observed prior to the wildfires, but one that is known to respond positively to high severity fire (Barclay *et al.*, 2004).

Given the moderate severity fires affected subsequent fire severity, whereas the low severity fires did not, we expected that impacts to forest structure characteristics and understory vegetation composition would be different in the moderate severity plots re-burned during the high severity wildfires versus the prior control plots that burned during the high severity wildfires, and our analyses indicated that this was the case. However, we expected the opposite result for low severity plots, and our analyses indicated the low severity burn plots also differed from the prior control plots. The reasons for this are unclear, but could be a low sample size ($n = 2$) effect for twice-burned low severity plots.

As with the vegetation results, we found that in contrast to high severity fire, low and moderate severity fire had no detectable effect on community composition of ground-dwelling arthropods. Further, the arthropod wildfire analysis agreed with our expectation for re-burned areas, traps in moderate fire severity zones would differ from those in high severity zones that were controls prior to the high severity wildfires; whereas, traps in low fire severity zones would be similar. However, there were few arthropod groups that seemed to have a strong response to fire within the time-frame of our study, and there were no clear correlations related to feeding guilds. Qualitatively, the predators Scorpiones (scorpions [*Centruroides vittatus*]; Taber *et al.*, 2003), Opiliones (harvestmen) and Araneae (spiders) were negatively impacted, whereas the predatory beetle family Carabidae showed a positive response. Similarly, the herbivorous Gryllidae

(crickets) showed a negative response, whereas the herbivorous Gryllacrididae (camel crickets) and Curculionidae (snout beetles) showed a positive response. However, our univariate analyses indicated most of these responses were not strong. Taber *et al.* (2008) found that low severity prescribed burning at another site in the Lost Pines positively impacted Diplopoda (millipedes), whereas we did not detect a strong response for diplopods, and the response direction was negative. Thus, as is typical with studies assessing fire impacts to terrestrial arthropods, our results were ambivalent and patterns were unclear.

This study has several implications with respect to management of the Houston toad using prescribed fire and in the Lost Pines post-wildfire landscape. First, one of the questions in relation to prescribed burning is whether winter or summer burns should be conducted (KES Consulting *et al.*, 2007). We believe there are benefits to conducting burns in both seasons. Summer burns were more effective for reducing fuel loads, and appeared to negatively impact some of the arthropod groups that are known predators of juvenile amphibians (Toledo, 2005), which are active during the summer in our study area (Brown *et al.*, 2011). In contrast, winter burns may positively impact tadpole growth and survivorship through increased aquatic productivity during the spring when the Houston toad breeds (Hillis *et al.*, 1984; Brown *et al.*, 2013). However, we note the relationships between pond productivity and tadpole growth and time-to-metamorphosis have not been examined for the Houston toad, and we recommend future research on this topic.

The wildfire dramatically impacted the terrestrial landscape. In our opinion, the most concerning of the impacts was complete overstory tree loss and elimination of fine

fuel throughout much of the high severity wildfire zone. The Houston toad exhibits strong preference for heavily canopied environments (U.S. Fish and Wildlife Service, 1984), and the suitability of the post-2011 wildfire landscape for this species is currently unclear. However, we note that although overstory canopy cover has been dramatically reduced, understory cover has increased since the completion of this study (D.J. Brown, personal observation). Further, although the majority of large fuel was consumed, which is the preferred refugia for adult Houston toads (Swannack, 2007), it is currently being rapidly replaced by fallen trees and large limbs. Thus, although in general the soil is likely warmer and drier within the wildfire zone (i.e., increased probability of desiccation), Houston toads could potentially mitigate this problem in the short-term by seeking out suitable refugia. However, we note that natural pine regeneration following the 2011 wildfire was low and patchy across most of the burn zone. Thus, assisted pine restoration through seedling tree planting is currently a major recovery initiative in the ecoregion (Lost Pines Recovery Team 2012).

With the increase in understory vegetation abundance and diversity an increase in overall arthropod abundance and diversity could occur in the coming years, which would be a positive impact on terrestrial food resources for the Houston toad (Neumann, 1991; Swengel, 2001; Moretti *et al.*, 2004; Buddle *et al.*, 2006). However, a concern with respect to arthropod responses is a potential increase in distribution and density of the invasive red imported fire ant (*Solenopsis invicta*), a known predator of juvenile Houston toads (Freed and Neitman, 1988), and possible predator of adult Houston toads (M.C. Jones, personal observation), given that abundance of this species appears to be strongly inversely correlated with overstory canopy cover in our study area (Brown *et al.*, 2012,

2013). Thus, although the high severity wildfire could certainly have both short-term and long-term negative impacts on the Houston toad, it could also have positive impacts, and we believe it is inappropriate for managers to consider the burn zone as unsuitable habitat without the population trend data to support that assumption.

In conclusion, this study represented a unique opportunity to compare the influence of fire severity on several ecosystem components within a BACI framework, and our results indicated this factor affected the ecosystem components we assessed. Given our results, future work should seek to address if multiple low to moderate severity fires produce the desirable outcomes of high severity fires (e.g., increased diversity of understory vegetation), while minimizing the unwanted outcomes (e.g., significant loblolly pine mortality). We intend to continue monitoring the response of this ecoregion to the high severity wildfires to assess the longer-term impacts to biotic and abiotic ecosystem components, and to provide data that can assist with recovery and management initiatives.

5. Acknowledgments

We thank J. Becker and P. Diaz for assistance with aquatic habitat sampling and analyses, and D. Lemke and J. Angermeier for assistance with plant identification. We thank B. DeVolld, A. Harper, M. Jones, A. Bosworth, A. Gomez, A. Schultz, M. Ray, C. Wiggins, K. Jamerson, E. Feichtinger, S. Dolino, J. Lattanzio, J. Shaw, J. Vrla, R. Bogan, J. Barnett, A. Duarte, K. Wooten, M. Heather, S. Uthuppuru, and D. Wallace for assistance with arthropod identification, sampling, and sorting. We thank Jay and Joshua Kane, T. Schumann, M. Sanders, J. Chenoweth, B. Armstrong, and L. Kothmann for

assistance with prescribed burns. We thank the Texas Forest Service and local fire departments for assistance with controlling the wildfires. This study was supported by the Texas Parks and Wildlife Department and U. S. Fish and Wildlife Service through a traditional Section 6 grant. The Capitol Area Council of the Boy Scouts of America provided access to the Griffith League Ranch, and we appreciate their continuing support of our research.

6. References

- Agee, J.K., 1996. Achieving conservation biology objectives with fire in the Pacific Northwest. *Weed Technol.* 10, 417–421.
- Al-Rabah'ah, M.A., Williams, C.G., 2004. An ancient bottleneck in the Lost Pines of central Texas. *Mol. Ecol.* 13, 1075–1084.
- Allen, C.D., Savage, M., Falk, D.A., Suckling, K.F., Swetnam, T.W., Schulke, T., Stacey, P.B., Morgan, P., Hoffman, M., Klingel, J.T., 2002. Ecological restoration of southwestern ponderosa pine ecosystems: a broad perspective. *Ecol. Appl.* 12, 1418–1433.
- Arkle, R.S., Pilliod, D.S., 2010. Prescribed fires as ecological surrogates for wildfires: a stream and riparian perspective. *For. Ecol. Manag.* 259, 893–903.
- Baker, F.E., 1979. Soil survey of Bastrop County, Texas. U.S. Department of Agriculture, Soil Conservation Service, Washington, D.C.
- Barclay, A.D., Betancourt, J.L., Allen, C.D., 2004. Effects of seeding ryegrass (*Lolium multiflorum*) on vegetation recovery following fire in a ponderosa pine (*Pinus ponderosa*) forest. *Int. J. Wildland Fire* 13, 183–194.

- Battle, J., Golladay, S.W., 2003. Prescribed fire's impact on water quality of depressional wetlands in southwestern Georgia. *Am. Midl. Nat.* 150, 15–25.
- Benavides, R., Rabasa, S.G., Granda, E., Escudero, A., Hodar, J.A., Martinez-Vilalta, J., Rincon, A.M., Zamora, R., Valladares, F., 2013. Direct and indirect effects of climate on demography and early growth of *Pinus sylvestris* at the rear edge: changing roles of biotic and abiotic Factors. *Plos One* 8, e59824.
- Bisson, P.A., Rieman, B.E., Luce, C., Hessburg, P.F., Lee, D.C., Kershner, J.L., Reeves, G.H., Gresswell, R.E., 2003. Fire and aquatic ecosystems of the western USA: current knowledge and key questions. *Forest Ecol. Manag.* 178, 213–229.
- Brown, D.J., Baccus, J.T., Means, D.B., Forstner, M.R.J., 2011. Potential positive effects of fire on juvenile amphibians in a southern USA pine forest. *J. Fish Wildlife Manage.* 2, 135–145.
- Brown, D.J., DeVolld, B., Nowlin, W.H., Forstner, M.R.J., 2012. Fire ants in Houston toad habitat: annual activity and responses to canopy cover and fire. *J. Fish Wildlife Manage.* 3, 142–149.
- Brown, D.J., Preston, D.B., Ozel, E., Forstner, M.R.J., 2013. Wildfire impacts on red imported fire ant captures in the Lost Pines ecoregion of Texas. *J. Fish Wildlife Manage.* 4, 129–133.
- Brown, D.J., Swannack, T.M., Forstner, M.R.J., 2013. Predictive models for calling and movement activity of the endangered Houston toad (*Bufo houstonensis*). *Am. Midl. Nat.* 169, 303–321.
- Bryant, V.M., Jr., 1977. A 16,000 year pollen record of vegetational change in central Texas. *Palynology* 1, 143–156.

- Buddle, C.M., Langor, D.W., Pohl, G.R., Spence, J.R., 2006. Arthropod responses to harvesting and wildfire: implications for emulation of natural disturbance in forest management. *Biol. Conserv.* 128, 346–357.
- Cain, M.D., Wigley, T.B., Reed, D.J., 1998. Prescribed fire effects on structure in uneven-aged stands of loblolly and shortleaf pines. *Wildl. Soc. Bull.* 26, 209–218.
- Certini, G., 2005. Effects of fire on properties of forest soils: a review. *Oecologia* 143, 1–10.
- Collins, B.M., Stephens, S.L., Moghaddas, J.J., Battles, J., 2010. Challenges and approaches in planning fuel treatments across fire-excluded forested landscapes. *J. Forest.* 108, 24–31.
- Coops, N.C., Waring, R.H., 2011. Estimating the vulnerability of fifteen tree species under changing climate in Northwest North America. *Ecol. Model.* 222, 2119–2129.
- Crumpton, W.G., Isenhardt, T.M., Mitchell, P.D., 1992. Nitrate and organic N analyses with second-derivative spectroscopy. *Limnol. Oceanogr.* 37, 907–913.
- Davis, C., 2001. The west in flames: the intergovernmental politics of wildfire suppression and prevention. *Publius* 31, 97–110.
- Dickson, B.G., Noon, B.R., Flather, C.H., Jentsch, S., Block, W.M., 2009. Quantifying the multi-scale response of avifauna to prescribed fire experiments in the southwest United States. *Ecol. Appl.* 19, 608–621.
- Duarte, A., Brown, D.J., Forstner, M.R.J., 2011. Estimating abundance of the endangered Houston toad on a primary recovery site. *J. Fish Wildlife Manage.* 2, 207–215.

- Duran, J., Rodriguez, A., Fernandez-Palacios, J.M., Gallardo, A., 2010. Long-term decrease of organic and inorganic nitrogen concentrations due to pine forest wildfire. *Ann. For. Sci.* 67, 207.
- Earl, S.R., Blinn, D.W., 2003. Effects of wildfire ash on water chemistry and biota in south-western USA streams. *Freshw. Biol.* 48, 1015–1030.
- Engstrom, R.T., 2010. First-order fire effects on animals: review and recommendations. *Fire Ecol.* 6, 115–130.
- Fontaine, J.B., Kennedy, P.L., 2012. Meta-analysis of avian and small-mammal response to fire severity and fire surrogate treatments in U.S. fire-prone forests. *Ecol. Appl.* 22, 1547–1561.
- Ford, W.M., Menzel, M.A., McGill, D.W., Laerm, J., McCay, T.S., 1999. Effects of a community restoration fire on small mammals and herpetofauna in the southern Appalachians. *For. Ecol. Manag.* 114, 233–243.
- Fowler, J., Cohen, L., Jarvis, P., 1998. *Practical Statistics for Field Biology*, second ed. Wiley, New York.
- Freed, P.S., Neitman, K., 1988. Notes on predation of the endangered Houston toad, *Bufo houstonensis*. *Tex. J. Sci.* 40, 454–456.
- Gaertner, J.P., Brown, D.J., Mendoza, J.A., Forstner, M.R.J., Bonner, T., Hahn, D., 2012. Geographic variation in *Batrachochytrium dendrobatidis* occurrence among populations of *Acris crepitans blanchardi* in Texas, USA. *Herpetol. Rev.* 43, 274–278.

- Galiano, L., Martinez-Vilalta, J., Lloret, F., 2010. Drought-induced multifactor decline of scots pine in the Pyrenees and potential vegetation change by the expansion of co-occurring oak species. *Ecosystems* 13, 978–991.
- Gaston, M.A., Fuji, A., Weckerly, F.W., Forstner, M.R.J., 2010. Potential component allee effects and their impact on wetland management in the conservation of endangered anurans. *Plos One* 5, e10102.
- Gauthier, S., Bergeron, Y., Simon, J.-P., 1996. Effects of fire regime on the serotiny level of jack pine. *J. Ecol.* 84, 539–548.
- Gebert, K.M., Black, A.E., 2012. Effect of suppression strategies on federal wildland fire expenditures. *J. Forest.* 110, 65–73.
- Geluso, K.N., Bragg, T.B., 1986. Fire-avoidance behavior of meadow voles (*Microtus pennsylvanicus*). *Am. Midl. Nat.* 116, 202–205.
- Gilliam, F.S., 1988. Interactions of fire with nutrients in the herbaceous layer of a nutrient- poor coastal plain forest. *B. Torrey Bot. Club* 115, 265–271.
- Gilliam, F.S., Platt, W.J., 1999. Effects of long-term fire exclusion on tree species composition and stand structure in an old-growth *Pinus palustris* (longleaf pine) forest. *Plant Ecol.* 140, 15–26.
- Glasgow, L.S., Matlack, G.R., 2007. Prescribed burning and understory composition in a temperate deciduous forest, Ohio, USA. *For. Ecol. Manag.* 238, 54–64.
- Greenberg, C.H., Forrest, T.G., Waldrop, T., 2010. Short-term response of ground-dwelling arthropods to prescribed fire and mechanical fuel reduction in a southern Appalachian upland hardwood forest. *For. Sci.* 56, 112–121.

- Greenberg, C.H., Waldrop, T.A., 2008. Short-term response of reptiles and amphibians to prescribed fire and mechanical fuel reduction in a southern Appalachian upland hardwood forest. *For. Ecol. Manag.* 255, 2883–2893.
- Gresswell, R.E., 1999. Fire and aquatic ecosystems in forested biomes of North America. *Trans. Am. Fish. Soc.* 128, 193–221.
- Grosman, D.M., Upton, W.W., 2006. Efficacy of systemic insecticides for protection of loblolly pine against southern fine engraver beetles (Coleoptera: Curculionidae: Scolytinae) and wood borers (Coleoptera: Cerambycidae). *J. Econ. Entomol.* 99, 94–101.
- Hall, S.A., Burke, I.C., Hobbs, N.T., 2006. Litter and dead wood dynamics in ponderosa pine forests along a 160-year chronosequence. *Ecol. Appl.* 16, 2344–2355.
- Hanberry, B.B., Dey, D.C., He, H.S., 2012. Regime shifts and weakened environmental gradients in open oak and pine ecosystems. *Plos One* 7, e41337.
- Hartnett, D.C., Krofta, D.M., 1989. Fifty-five years of post-fire succession in a southern mixed hardwood forest. *B. Torrey Bot. Club* 116, 107–113.
- Heiri, O., Lotter, A.F., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *J. Paleolimnol.* 25, 101–110.
- Hély, C., Bergeron, Y., Flannigan, M.D., 2000. Effects of stand composition on fire hazard in mixed-wood Canadian Boreal Forest. *J. Veg. Sci.* 11, 813–824.
- Hillis, D.M., Hillis, A.M., Martin, R.F., 1984. Reproductive ecology and hybridization of the endangered Houston toad (*Bufo houstonensis*). *Journal of Herpetology* 18, 56–72.

- Hitt, N.P., 2003. Immediate effects of wildfire on stream temperature. *J. Freshwater Ecol.* 18, 171–173.
- Hodgkins, E.J., 1958. Effects of fire on undergrowth vegetation in upland southern pine forests. *Ecology* 39, 36–46.
- Houghton, R.A., Hackler, J.L., Lawrence, K.T., 2000. Changes in terrestrial carbon storage in the United States. 2: the role of fire and fire management. *Glob. Ecol. Biogeogr.* 9, 145–170.
- Jones, D.D., Conner, L.M., Storey, T.H., Warren, R.J., 2004. Prescribed fire and raccoon use of longleaf pine forests: implications for managing nest predation? *Wildl. Soc. Bull.* 32, 1255–1259.
- KES Consulting, Loomis Austin, Forstner, M.R.J., 2007. Lost Pines habitat conservation plan for Bastrop County, Texas. http://www.co.bastrop.tx.us/site/content/lostpines_habitat (accessed 6.17.13).
- Knapp, E.E., Estes, B.L., Skinner, C.N., 2009. Ecological effects of prescribed fire season: a literature review and synthesis for managers. U.S. Department of Agriculture, General Technical Report PSW–GTR–224, Redding, CA.
- Knebel, L., Wentworth, T.R., 2007. Influence of fire and southern pine beetle on pine-dominated forests in the Linville Gorge Wilderness, North Carolina. *Castanea* 72, 214–225.
- Lecomte, N., Simard, M., Fenton, N., Bergeron, Y., 2006. Fire severity and long-term ecosystem biomass dynamics in coniferous Boreal Forests of eastern Canada. *Ecosystems* 9, 1215–1230.

- Lepš, J., Šmilauer, P., 2003. *Multivariate Analysis of Ecological Data using CANOCO*. Cambridge University Press, New York.
- Lewis, W.M., Jr., 1974. Effects of fire on nutrient movement in a South Carolina pine forest. *Ecology* 55, 1120–1127.
- Littell, J.S., Oneil, E.E., McKenzie, D., Hicke, J.A., Lutz, J.A., Norheim, R.A., Elsner, M.M., 2010. Forest ecosystems, disturbance, and climatic change in Washington State, USA. *Clim. Change* 102, 129–158.
- Long, R.A., Rachlow, J.L., Kie, J.G., 2009. Sex-specific responses of North American elk to habitat manipulation. *J. Mammal.* 90, 423–432.
- Lost Pines Recovery Team, 2011. Bastrop County complex fire Lost Pines region: resource assessment & response report. Lost Pines Recovery Team, Bastrop County, Texas, USA. <http://bastroprecovery.org> (accessed 6.17.13).
- Lost Pines Recovery Team, 2012. Bastrop Lost Pines habitat recovery project master plan. Lost Pines Recovery Team, Bastrop County, Texas, USA. <http://bastroprecovery.org> (accessed 6.17.13).
- Malison, R.L., Baxter, C.V., 2010. Effects of wildfire of varying severity on benthic stream insect assemblages and emergence. *J. N. Am. Benthol. Soc.* 29, 1324–1338.
- McEachern, P., Prepas, E.E., Gibson, J.J., Dinsmore, W.P., 2000. Forest fire induced impacts on phosphorus, nitrogen, and chlorophyll a concentrations in Boreal subarctic lakes of northern Alberta. *Can. J. Fish. Aquat. Sci.* 57, 73–81.

- Miller, J.D., Safford, H.D., Crimmins, M., Thode, A.E., 2009. Quantitative evidence for increasing forest fire severity in the Sierra Nevada and Southern Cascade Mountains, California and Nevada, USA. *Ecosystems* 12, 16–32.
- Mitchell, R., Cathey, J.C., Dabbert, B., Prochaska, D.F., DuPree, S., Ron, S., 2005. Managing yaupon with fire and herbicides in the Texas Post Oak Savannah. *Rangelands* 27, 17–19.
- Mitchell, S.R., Harmon, M.E., O'Connell, K.E.B., 2009. Forest fuel reduction alters fire severity and long-term carbon storage in three Pacific Northwest ecosystems. *Ecol. Appl.* 19, 643–655.
- Moore, B., 1977. Bastrop County 1691–1900, revised ed. Nortex Press, Wichita Falls, TX.
- Moore, M.M., Covington, W.W., Fule, P.Z., 1999. Reference conditions and ecological restoration: a southwestern ponderosa pine perspective. *Ecol. Appl.* 9, 1266–1277.
- Moretti, M., Obrist, M.K., Duelli, P., 2004. Arthropod biodiversity after forest fires: winners and losers in the winter fire regime of the southern Alps. *Ecography* 27, 173–186.
- Moritz, M.A., Parisien, M.-A., Batllori, E., Krawchuk, M.A., Van Dorn, J., Ganz, D.J., Hayhoe, K., 2012. Climate change and disruptions to global fire activity. *Ecosphere* 3, article 49.
- National Park Service. 2003. Fire monitoring handbook. Fire Management Program Center, Boise, ID.

- Neumann, F.G., 1991. Responses of litter arthropods to major natural or artificial ecological disturbances in mountain ash forest. *Aust. J. Ecol.* 16, 19–32.
- Nowacki, G.J., Abrams, M.D., 2008. The demise of fire and "mesophication" of forests in the eastern United States. *Bioscience* 58, 123–138.
- Oosting, H.J., 1944. The comparative effect of surface and crown fire on the composition of a loblolly pine community. *Ecology* 25, 61–69.
- Owen, J.G., 1989. Patterns of herpetofaunal species richness: relation to temperature, precipitation, and variance in elevation. *J. Biogeogr.* 16, 141–150.
- Rieman, B., Lee, D., Burns, D., Gresswell, R., Young, M., Stowell, R., Rinne, J., Howell, P., 2003. Status of native fishes in the western United States and issues for fire and fuels management. *Forest Ecol. Manag.* 178, 197–211.
- Roberts, S.L., van Wagtendonk, J.W., Miles, A.K., Kelt, D.A., Lutz, J.A., 2008. Modeling the effects of fire severity and spatial complexity on small mammals in Yosemite National Park, California. *Fire Ecol.* 4, 83–104.
- Sacerdote, A.B., King, R.B., 2009. Dissolved oxygen requirements for hatching success of two ambystomatid salamanders in restored ephemeral ponds. *Wetlands* 29, 1202–1213.
- Safford, H.D., Stevens, J.T., Merriam, K., Meyer, M.D., Latimer, A.M., 2012. Fuel treatment effectiveness in California yellow pine and mixed conifer forests. *For. Ecol. Manag.* 274, 17–28.
- Schoennagel, T., Veblen, T.T., Romme, W.H., 2004. The interaction of fire, fuels, and climate across Rocky Mountain forests. *Bioscience* 54, 661–676.

- Schulte, L.A., Mladenoff, D.J., 2005. Severe wind and fire regimes in northern forests: historical variability at the regional scale. *Ecology* 86, 431–445.
- Schwilk, D.W., Knapp, E.E., Ferrenberg, S.M., Keeley, J.E., Caprio, A.C., 2006. Tree mortality from fire and bark beetles following early and late season prescribed fires in a Sierra Nevada mixed-conifer forest. *For. Ecol. Manag.* 232, 36–45.
- Shang, Z., He, H.S., Lytle, D.E., Shifley, S.R., Crow, T.R., 2007. Modeling the long-term effects of fire suppression on central hardwood forests in Missouri Ozarks, using LANDIS. *For. Ecol. Manag.* 242, 776–790.
- Simmons, M.T., Windhager, S., Power, P., Lott, J., Lyons, R.K., Schwope, C., 2007. The response of old world bluestem to mid- and late-summer fire in two grassland ecoregions in central Texas, in: Masters, R.E., Galley, K.E.M. (Eds.), *Proceedings, 23rd, Tall Timbers Fire Ecology Conference: Fire in Grassland and Shrubland Ecosystems*. Tall Timbers Research Station, Tallahassee, FL, pp. 246–250.
- Smucker, K.M., Hutto, R.L., Steele, B.M., 2005. Changes in bird abundance after wildfire: importance of fire severity and time since fire. *Ecol. Appl.* 15, 1535–1549.
- Sparks, J.C., Masters, R.E., Engle, D.M., Palmer, M.W., Bukenhofer, G.A., 1998. Effects of late growing-season and late dormant-season prescribed fire on herbaceous vegetation in restored pine-grassland communities. *J. Veg. Sci.* 9, 133–142.
- Spencer, C.N., Hauer, F.R., 1991. Phosphorus and nitrogen dynamics in streams during a wildfire. *J. N. Am. Benthol. Soc.* 10, 24–30.

- Spencer, C.N., Gabel, K.O., Hauer, F.R., 2003. Wildfire effects on stream food webs and nutrient dynamics in Glacier National Park, USA. *For. Ecol. Manag.* 178, 141–153.
- Stambaugh, M.C., Guyette, R.P., Marschall, J.M., 2011. Longleaf pine (*Pinus palustris* Mill.) fire scars reveal new details of a frequent fire regime. *J. Veg. Sci.* 22, 1094–1104.
- Swannack, T.M., 2007. Modeling aspects of the ecological and evolutionary dynamics of the endangered Houston toad. PhD dissertation. Texas A&M University, College Station, Texas, 155 p.
- Sweeney, J.R., Biswell, H.H., 1961. Quantitative studies of the removal of litter and duff by fire under controlled conditions. *Ecology* 42, 572–575.
- Swengel, A.B., 2001. A literature review of insect responses to fire, compared to other conservation managements of open habitat. *Biodivers. Conserv.* 10, 1141–1169.
- Taber, S.W., Fleenor, S.B., 2003. *Insects of the Texas Lost Pines*. Texas A&M University Press, College Station.
- Taber, S.W., Fleenor, S.B., Privette, J.L., 2008. Arthropod abundance following a prescribed burn in the Lost Pines forest of central Texas. *Southwest. Entomol.* 33, 65–77.
- Taylor, A.H., 2000. Fire regimes and forest changes in mid and upper montane forests of the southern Cascades, Lassen Volcanic National Park, California, U.S.A. *J. Biogeogr.* 27, 87–104.

- Thies, W.G., Westlind, D.J., 2012. Validating the Malheur model for predicting ponderosa pine post-fire mortality using 24 fires in the Pacific Northwest, USA. *Int. J. Wildland Fire* 21, 572–582.
- Tinker, D.B., Knight, D.H., 2000. Coarse woody debris following fire and logging in Wyoming lodgepole pine forests. *Ecosystems* 3, 472–483.
- Toledo, L.F., 2005. Predation of juvenile and adult amphibians by invertebrates: current knowledge and perspectives. *Herpetol. Rev.* 36, 395–400.
- U.S. Fish and Wildlife Service., 1984. Houston toad recovery plan. U.S. Fish and Wildlife Service, Albuquerque, New Mexico, USA. <http://ecos.fws.gov/speciesProfile/profile/speciesProfile.action?sPCODE=D004>.
- Van den Brink, P.J., Ter Braak, C.J.F., 1999. Principal response curves: analysis of time-dependent multivariate responses of biological community to stress. *Environ. Toxicol. Chem.* 18, 138–148.
- Vose, J.M., 2000. Perspectives on using prescribed fire to achieve desired ecosystem conditions, in: Moser, W.K., Moser, C. E. (Eds.), *Proceedings, 21st, Tall Timbers Fire Ecology Conference: Fire and Forest Ecology: Innovative Silviculture and Vegetation Management*. Tall Timbers Research Station, Tallahassee, FL, pp. 12–17.
- Waldrop, T.A., White, D.L., Jones, S.M., 1992. Fire regimes for pine-grassland communities in the southeastern United States. *For. Ecol. Manag.* 47, 195–210.
- Wan, S., Hui, D., Luo, Y., 2001. Fire effects on nitrogen pools and dynamics in terrestrial ecosystems: a meta-analysis. *Ecol. Appl.* 11, 1349–1365.

- Wayman, R.B., North, M., 2007. Initial response of a mixed-conifer understory plant community to burning and thinning restoration treatments. *For. Ecol. Manag.* 239, 32–44.
- Webster, K.M., Halpern, C.B., 2010. Long-term vegetation responses to reintroduction and repeated use of fire in mixed-conifer forests of the Sierra Nevada. *Ecosphere* 1, article 9.
- Wetzel, R.G., Likens, G.E., 2000. *Limnological Analyses*, third ed. Springer, New York.
- Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M., 2009. *Mixed Effects Models and Extensions in Ecology with R*. Springer Science+Business Media, New York.

CHAPTER V

FIRE ANTS IN HOUSTON TOAD HABITAT: ANNUAL ACTIVITY AND RESPONSES TO CANOPY COVER AND FIRE

Format follows the guidelines for the Journal of Fish and Wildlife Management

V. Fire Ants in Houston Toad Habitat: Annual Activity and Responses to Canopy Cover and Fire

Abstract

The red imported fire ant (RIFA; *Solenopsis invicta*) is an invasive species found throughout the south and southeastern U.S. Since its introduction, RIFA has been shown to negatively impact a wide range of native vertebrate and invertebrate species. The purposes of this study were to delineate the annual RIFA activity pattern, investigate the association between overstory canopy cover and RIFA captures, and evaluate the effects of low and moderate severity prescribed fire around pond edges on RIFA in the Lost Pines ecoregion of Texas, which provides habitat for most of the remaining endangered Houston toads (*Bufo* [*Anaxyrus*] *houstonensis*). We found that annual RIFA activity followed a quadratic curve, with above average activity between May and October. We found an inverse relationship between mean percent canopy cover near pond edges and mean number of RIFA captured. We found low and moderate severity prescribed burning had no significant influence on RIFA captures during our study period. However, strong spatial and temporal capture variability was apparent, and thus a strong impact would have been necessary to detect an effect. While this study provides evidence that should decrease concerns that wildlife managers, conservation biologists, and landowners in the Lost Pines ecoregion may have about exacerbating RIFA abundance when utilizing fire as an ecosystem management tool, we recommend additional work be conducted using a greater sample size, greater sampling effort, and longer study duration. We found that RIFA activity was highest during the time frame in which juvenile Houston toads emerge

from ponds. Thus, RIFA control may be a useful Houston toad recovery tool where breeding ponds are not within dense canopy habitats.

Keywords: Houston toad; Lost Pines; prescribed fire; red imported fire ant; Texas.

Introduction

The red imported fire ant (RIFA; *Solenopsis invicta*) is an invasive species in the U.S., predominantly in the south and southeast. Since its introduction to Alabama in the 1930s, the distribution of RIFA has expanded east to North Carolina and west to central Texas (Callcott and Collins 1996), and the species has been introduced to California (Ward 2005). When introduced, RIFA has been shown to negatively impact a wide range of native vertebrate and invertebrate species (Porter and Savignano 1990; Allen et al. 1994; Allen et al. 2004; Stuble et al. 2009; Diffie et al. 2010; Epperson and Allen 2010). Vertebrates are particularly vulnerable to RIFA predation during early life-stages (Landers et al. 1980; Freed and Neitman 1988; Pedersen et al. 1996). Abundance and diversity of native arthropods can also be negatively affected by RIFA competition and predation (Morrison 2002; Plowes et al. 2007; Epperson and Allen 2010).

In the U.S. the RIFA prefers environments characterized by disturbance, and can use roads and powerline cuts as expansion corridors (Tschinkel 1988; Stiles and Jones 1998; Todd et al. 2008). In primarily forested habitats, RIFA tend to be more abundant in forest gaps (Colby and Prowell 2006). The RIFA is adapted to periodically flooded habitats (Allen et al. 1974), and has been found to prefer edges of ponds and other water bodies (Lyle and Fortune 1948; Tschinkel 1988; Stuble et al. 2009; Vogt et al. 2009).

Because RIFA can be more flood tolerant than some native ant species, it may have a competitive advantage over these taxa in flood-prone environments (Vogt et al. 2009). The potential for high RIFA density around ponds is of concern in amphibian management, given the potential vulnerability of juvenile amphibians to RIFA predation when they enter the terrestrial landscape (Freed and Neitman 1988). Red imported fire ants prefer canopy gaps, thus ponds with less canopy cover or near the edges of forested fragments may have higher RIFA abundances and may subsequently have higher RIFA predation pressure on juvenile amphibians.

The RIFA arrived in Bastrop County, central Texas between 1973 and 1977, and are now well-established in the Lost Pines ecoregion (Cokendolpher and Phillips 1989; Taber and Fleenor 2003). This ecoregion is a 34,400 ha remnant of a pine-dominated forest that occurred in east and east-central Texas approximately 14,000 to 10,000 years ago (Bryant 1977). The Lost Pines houses the majority of the remaining breeding aggregations of the endangered Houston toad (*Bufo* [*Anaxyrus*] *houstonensis*; Gottschalk 1970), a species that has declined rapidly in the last half century due primarily to habitat loss and degradation (U.S. Fish and Wildlife Service 1984; Brown and Mesrobian 2005). The RIFA is known to prey upon Houston toads after their emergence from ponds as terrestrial juveniles (Freed and Neitman 1988).

We conducted this investigation as part of a larger study on the use of prescribed fire for Houston toad habitat recovery and conservation initiatives. The majority of the Lost Pines has been fire suppressed for the past century, resulting in heavy fuel loads and dense thickets of fire-intolerant shrubs, primarily yaupon holly (*Ilex vomitoria*). Currently there is limited published information on RIFA response to prescribed burning, and RIFA

population responses to fire are unclear. Forbes (1999) documented short-term (i.e., five months post-burn) negative effects on RIFA due to prescribed burning in a Texas coastal prairie, presumably due to reduced soil moisture and food availability. Conversely, Norton (2003) found no effect of burning on RIFA abundance in habitats similar to those of Forbes (1999). In contrast, Hanula and Wade (2003) found that RIFA abundances increased with burn frequency in a Florida longleaf pine (*Pinus palustris*) forest. Thus, the results of the aforementioned studies present an unclear picture of whether the use of prescribed fire as a recovery tool in pine-dominated ecosystems may increase invasive potential for RIFA or whether fire may be an effective method to reduce RIFA in invaded pine forests.

The purposes of this study were to delineate the annual RIFA activity pattern in the Lost Pines ecoregion of Texas, and to investigate RIFA responses to overstory canopy cover and low and moderate severity prescribed fire. We were interested in temporal activity in order to better understand RIFA activity patterns in relation to Houston toad activity patterns, and to delineate optimum time intervals for RIFA control measures. Percent canopy cover is an easily quantifiable variable that could be used to determine which Houston toad ponds likely contain large RIFA populations, and thus where emergent Houston toads are most vulnerable to RIFA predation. Prescribed fire is a common habitat management tool used across the south and southeastern U.S., and thus it is important to know the impacts of fire on this invasive species.

Study Site

We conducted this study on the 1,900 ha Griffith League Ranch (GLR) in Bastrop County, Texas. The GLR is primarily forested, with vegetation typical of the Lost Pines ecoregion. The overstory is dominated by loblolly pine (*Pinus taeda*), post oak (*Quercus stellata*), blackjack oak (*Quercus marilandica*), and eastern red cedar (*Juniperus virginiana*), and the understory is dominated by yaupon holly (*Ilex vomitoria*), American beautyberry (*Callicarpa americana*), and farkleberry (*Vaccinium arboreum*).

The GLR contains three permanent ponds (i.e., ponds have not dried in at least 12 years), 10 semi-permanent ponds (i.e., ponds typically dry several times per decade), and dozens of ephemeral pools that hold water for days to months annually depending on rainfall. Ten of these ponds were used for this study. Three of the ponds were ephemeral (i.e. typically dry several times per year), six were semi-permanent (i.e., typically dry several times per decade), and one was permanent (i.e., has not dried in at least 11 years). Four of the pond edges were primarily covered by pine and oak leaf litter (>80% ground cover), with the remaining ponds having edges dominated by grasses and forbs (>80% ground cover). We chose these ponds because they spanned the distribution of size, hydroperiod, and canopy cover variability on the GLR, they were known Houston toad breeding ponds, and, when applicable, they were located in units projected to be burned. Houston toad calling activity occurred at all ponds used in this study within the previous two years (Duarte et al. 2011).

Methods

We conducted this study over a one-year period between 26 September 2009 and 10 October 2010, sampling the area around each of the 10 ponds a total of 27 times. Days between sampling ranged from 0 (i.e., consecutive sampling days) to 73, with a mean of 15 days between samples. We used a standard bait cup method for sampling ants (Porter and Tschinkel 1987; Mueller et al. 1999), which consisted of half a Vienna sausage placed in a 9-cm diameter round plastic dish. We placed between 3 and 15 dishes around the perimeter of each pond, depending on pond size (i.e., perimeter length). During each sampling event, we placed dishes 1-2 m from the pond edge at 3-4 m intervals, and allowed them to attract ants for 30 minutes. After 30 minutes we collected ants from each dish and euthanized them by freezing. We removed native ants from samples and RIFA were dried at $\sim 50^{\circ}\text{C}$ for at least 48 hours. We then counted RIFA collected in each bait cup at each pond on each sampling date. Alternately, we weighed the cumulative number of ants in each bait cup at each pond on each sampling date, and sample weight was estimated to the nearest 1×10^{-4} g. We estimated RIFA abundances in these samples using an empirically-derived relationship between a known number of RIFA and the weight of the sample. This relationship was derived from samples collected from around study ponds on three randomly selected sampling dates ($n = 109$ samples, $y = 2140.8 \times \text{weight (g)}$, $r^2 = 0.92$). We used the estimated RIFA abundances as count data in statistical analyses.

Four of the ponds were subjected to prescribed burning, with all ponds sampled before and after burning. However, because ponds were spread over multiple burn units, areas around individual ponds were burned at different times of the year. The area around

one pond was burned on 13 November 2009 (Pond 10), one pond on 10 January 2010 (Pond 13), and the remaining two ponds were burned on 21 August 2010 (ponds 14 and 15). This burn scenario created the ability to examine the effect of low severity winter fires (November and January) versus moderate severity summer fires (August). The goal for all fires was fuel reduction; burns removed the upper portion (i.e., 1 to 2 cm) of the leaf litter layer and charred but typically did not consume live woody vegetation.

In order to assess if RIFA counts were higher in open canopy areas, between 20 July 2008 and 16 April 2010, we estimated percent canopy cover around each pond between 4 and 24 times, depending on hydroperiod status at the time of sampling (i.e., estimates were not taken when ponds did not contain water). For each sample, we estimated percent canopy cover using a spherical densiometer (Forestry Suppliers Inc., Jackson, MS) at two to six randomly selected points at the pond edge, with higher numbers of estimation points corresponding to larger ponds. We then averaged the estimates at each pond per sampling date. For this study, we included measurements taken during the leaf-off period (December to March) because ponds with consistently high canopy cover at our study site were dominated by loblolly pine, which retains needles throughout the year.

Statistical analyses

We used a Generalized Additive Model (GAM; Hastie and Tibshirani 1990) to summarize the annual activity pattern based on our sampling results. We used the CPUE at each pond on each day sampled (i.e., 27 samples with 10 observations per sample), included Day of Year (DOY) as a predictor, and fit a smoothing curve to the data (cubic

regression spline), with the optimal amount of smoothing determined using a cross-validation algorithm (Zuur et al. 2009). Thus, the smoothing curve represented the model-fitted relationship between RIFA CPUE and DOY. Generalized Additive Models assume normality and homoscedasticity, and we investigated these assumptions using graphical diagnostics plots (Zuur et al. 2009). Because data appeared to violate the assumption of homoscedasticity, we transformed CPUE using the arcsinh (i.e., inverse hyperbolic sine) transformation (Fowler et al. 1998), which was effective. We performed this analysis using the program R (R Version 2.10.1, www.r-project.org), with the mgcv package (Wood 2004).

To examine the influence of canopy cover on RIFA CPUE, we utilized the RIFA capture and canopy cover data from all of the study ponds ($n = 10$), and calculated the mean percent canopy cover of each pond over the study period. We regressed mean RIFA CPUE on mean percent canopy cover for all ponds using ordinary least squares (OLS) linear regression, and gauged model fit using the coefficient of determination (r^2 ; Sokal and Rohlf 1995). We performed this analysis using program R (R Version 2.10.1, www.r-project.org).

To assess if prescribed fire influenced RIFA CPUE around ponds, we used the Before-After Control-Impact (BACI) approach described by Stewart-Oaten et al. (1986, 1992). One burn pond (Pond 13) was not included in the statistical analyses because we captured RIFA on very few sampling events, including none following the burn. We paired treatment (i.e., burned) ponds with control ponds that were similar in pre-burn temporal CPUE dynamics. For each pair, we computed the difference in CPUE between the control and treatment pond on each sampling occasion, and used a standard t -test to

determine if the mean difference between the control and treatment ponds changed after the prescribed fire. An important assumption with this analysis is that effects are additive (Stewart-Oaten et al. 1986). We tested this assumption using Tukey's test of additivity (Tukey 1949), and two of the three pond pairs indicated non-additive effects ($P < 0.001$). We transformed CPUE for all ponds using the arcsinh transformation, which was effective for all pairs ($P = 0.904$ [ponds 5 and 10]; $P = 0.392$ [ponds 9 and 14]; $P = 0.407$ [ponds 16 and 15]), and used the transformed data for the t -tests. Levene's test for equality of variances indicated that variances were equal for all comparisons, and thus unequal variance t -tests were not used. However, results from unequal variance t -tests were equivalent. We performed this analysis using program SPSS (SPSS Statistics Version 20).

Results

We captured an estimated 18,050 RIFA during this study, with estimated number of captures among ponds ranging from 4 (Pond 13) to 3,642 (Pond 11; Fig. 5.1). The GAM model fit a quadratic curve to the CPUE-DOY relationship, with above average activity between May and October (Fig. 5.2). We found a strong negative relationship between percent canopy cover and mean RIFA CPUE (Fig. 5.3; $t_8 = 11.08$, $P < 0.0001$, $r^2 = 0.82$). We did not detect a prescribed fire effect on RIFA CPUE for ponds burned in the winter (Pond 10: $t_{25} = 1.338$, $P = 0.193$; Pond 13: number of pre-burn captures = 4, number of post-burn captures = 0). We also did not detect a prescribed fire effect on RIFA CPUE for ponds burned in the summer (Pond 14: $t_{25} = -0.316$, $P = 0.755$; Pond 15: $t_{25} = -0.630$, $P = 0.534$).




Figure 5.1. Red imported fire ant (RIFA; *Solenopsis invicta*) captures at pond edges in the Lost Pines ecoregion of Texas. We sampled each pond 27 times between 26 September 2009 and 10 October 2010. The solid vertical bars delineate pre-burn and post-burn samples at burn ponds. Captures decreased immediately following a summer prescribed burn at ponds 14 and 15, but returned to background levels within weeks.




Figure 5.2. Annual activity pattern of red imported fire ants (RIFA) in the Lost Pines ecoregion of Texas based on captures-per-unit-effort (CPUE). We sampled RIFA 27 times at 10 ponds between 26 September 2009 and 10 October 2010. Sampling dates are shown with hash marks inside the x-axis. For this analysis we used CPUE at each pond on each day sampled (i.e., 27 samples with 10 observations per sample). We modeled CPUE as a function of Day of Year (DOY) using a Generalized Additive Model (GAM). The Y-axis shows the predicted activity level (with point-wise 95% confidence bands) based on results from days we sampled (inner tick marks) and relative to the observed mean CPUE for all observations (0). The vertical lines enclose the time period in which terrestrial juvenile Houston toads (*Bufo houstonensis*) may be present near ponds.




Figure 5.3. Relationship between mean percent canopy cover ($\pm$ SE) and mean red imported fire ant (RIFA) captures ($\pm$ SE) at 10 ponds in the Lost Pines ecoregion of Texas. We sampled RIFA at ponds 27 times between 26 September 2009 and 10 October 2010 and used the mean of the samples for this analysis. For percent canopy cover we used the mean of 4 to 24 samples taken between 20 July 2008 and 16 April 2010. We found a strong inverse relationship between mean RIFA captures and percent canopy cover.

Discussion

The results of this study indicate that disturbance from low and moderate severity prescribed burning did not affect RIFA abundance around ponds in our study area. This finding is in contrast to other studies reporting that disturbance from livestock use or vegetation-clearing can lead to increases in RIFA abundance (Lofgren et al. 1975; Stiles and Jones 1998; Todd et al. 2008; Vogt et al. 2009). In the present study, the number of RIFA captured at the two summer burn ponds decreased immediately after burning, but

returned to background levels within weeks. The winter burns were conducted during a period of low RIFA activity at all ponds, and no short-term changes were apparent. These results indicate that, at least in the short-term, prescribed burning did not benefit RIFA. However, each experimental pond was burned only once, and burn frequency may increase RIFA abundance (Hanula and Wade 2003). Additional research is needed concerning effects of burn frequency and severity on RIFA in both forest and non-forest systems. Further, due to strong spatial and temporal capture variability a strong impact would have been necessary to detect an effect. Future work on this topic would benefit from greater sample sizes, greater sampling effort, and longer study durations. We also note that RIFA were present at our study ponds prior to burning, and thus we did not address the potential for habitat invasion following disturbance through fire.

Percent canopy cover was a viable predictor of RIFA CPUE in our study area. This result was not surprising, as this species is known to select for open and edge habitats (Stiles and Jones 1998; Colby and Prowell 2006). We recommend that future studies seek to determine whether substrate-type (i.e., litter-dominated or vegetation-dominated) is an important RIFA CPUE predictor in forested environments, as fire can increase understory herbaceous vegetation density by removing litter and sunlight-capturing woody vegetation (Hodgkins 1958; Sparks et al. 1998; Donato et al. 2009). Because of constantly shifting pond borders due to deposition of precipitation and evapotranspiration, we were unable to test substrate-type in this study, as substrate-type shifted at several ponds. Quantifying substrate relationships would potentially increase predictive power for estimating RIFA densities around ponds, as well as increase our understanding of the causal mechanisms behind fire effects on RIFA.

Monitoring RIFA over the course of a year showed that activity varied seasonally, with highest activity between May and October. This includes the period where juvenile Houston toads are vulnerable to predation near ponds. Houston toad emergence from ponds in the Lost Pines typically occurs between March and June, and juveniles remain near the pond for the first one to two months after emergence (Greuter and Forstner 2003; Greuter 2004). Given that they are active, RIFA abundance differences could potentially significantly affect survivorship rates of terrestrial juvenile Houston toads. Thus, future research on juvenile Houston toad predation by the RIFA is warranted.

Based on the results of this study, low and moderate severity prescribed burning did not decrease RIFA CPUE around pond margins, suggesting that this kind of fire does not appear to be a useful management tool for eliminating RIFA. Chemical control has been shown to be locally effective, but insecticides can have severe impacts non-target wildlife, particularly amphibians (Lofgren et al. 1975; Boone and James 2003; Relyea 2003). If chemical control is necessary for managing RIFA in the Lost Pines ecoregion, we hypothesize that fall or winter application may have the lowest amphibian impact because amphibian activity is low relative to spring and summer (Brown et al. in press). Further, although broadcast application is the most effective way to control RIFA, strategic bait placement (e.g., on mounds) can also be effective (Williams et al. 2001; Allen et al. 2004). Because amphibians do not appear to actively avoid soils contaminated with harmful chemicals (Hatch et al. 2001; Storrs Méndez et al. 2009), and these chemicals can be readily absorbed through the skin, strategic bait placement is a more attractive management option. Biological controls, such as the fire ant decapitating fly (*Pseudacteon curvatus*) and the disease-causing protozoan *Thelohania solenopsae*,

may eventually prove to be effective broad-scale control agents (Brinkman and Gardner 2001; Porter 2010).

Acknowledgements

We thank L. Eddins for assistance with laboratory work and ant sampling, and W. DeVolld for assistance with ant sampling. J. Veech and B. Weckerly provided helpful statistical suggestions. This study was supported by the Texas Parks and Wildlife Department and U.S. Fish and Wildlife Service through a traditional Section 6 grant. This study was also funded in part by a Francis Rose undergraduate research scholarship awarded to B. DeVolld.

References

- Allen CR, Demarais S, Lutz RS. 1994. Red imported fire ant impact on wildlife- an overview. *Texas Journal of Science* 46:51–59.
- Allen CR, Epperson DM, Garmestani AS. 2004. Red imported fire ant impacts on wildlife: a decade of research. *American Midland Naturalist* 152:88–103.
- Allen GE, Buren WF, Williams RN, Demeneze M, Whitcomb WH. 1974. Red imported fire ant, *Solenopsis invicta*, distribution and habitat in Mato Grosso, Brazil. *Annals of the Entomological Society of America* 67:43–46.
- Boone MD, James SM. 2003. Interactions of an insecticide, herbicide, and natural stressors in amphibian community mesocosms. *Ecological Applications* 13:829–841.

- Brinkman MA, Gardner WA. 2001. Use of diatomaceous earth and entomopathogen combinations against the red imported fire ant (Hymenoptera: Formicidae). *Florida Entomologist* 84:740–741.
- Brown DJ, Swannack TM, Forstner MRJ. In press. Herpetofaunal survey of the Griffith League Ranch in the Lost Pines ecoregion of Texas. *Texas Journal of Science*.
- Brown LE, Mesrobian A. 2005. Houston toads and Texas politics. Pages 150–160 in Lannoo M, editor. *Amphibian declines: the conservation status of United States species*. Ewing, New Jersey: University of California Press.
- Bryant VM Jr. 1977. A 16,000 year pollen record of vegetational change in central Texas. *Palynology* 1:143–156.
- Callcott A-MA, Collins HL. 1996. Invasion and range expansion of imported fire ants (Hymenoptera: Formicidae) in North America from 1918-1995. *Florida Entomologist* 79:240–251.
- Cokendolpher JC, Phillips SA Jr. 1989. Rate of spread of the red imported fire ant, *Solenopsis invicta* (Hymenoptera: Formicidae), in Texas. *Southwestern Naturalist* 34:443–449.
- Colby D, Prowell D. 2006. Ants (Hymenoptera: Formicidae) in wet longleaf pine savannas in Louisiana. *Florida Entomologist* 89:266–269.
- Crawley MJ. 2007. *The R book*. England: John Wiley & Sons Ltd.
- Diffie S, Miller J, Murray K. 2010. Laboratory observations of red imported fire ant (Hymenoptera: Formicidae) predation on reptilian and avian eggs. *Journal of Herpetology* 44:294–296.

- Donato DC, Fontaine JB, Robinson WD, Kauffman JB, Law BE. 2009. Vegetation response to a short interval between high-severity wildfires in a mixed-evergreen forest. *Journal of Ecology* 97:142–154.
- Duarte A, Brown DJ, Forstner MRJ. 2011. Estimating abundance of the endangered Houston toad on a primary recovery site. *Journal of Fish and Wildlife Management* 2:207–215.
- Epperson DM, Allen CR. 2010. Red imported fire ant impacts on upland arthropods in southern Mississippi. *American Midland Naturalist* 163:54–63.
- Forbes A. 1999. Effect of bobwhite habitat management on red imported fire ant populations. Master's thesis. Lubbock: Texas Tech University.
- Fowler JF, Cohen L, Jarvis P. 1998. Practical statistics for field biology. 2nd edition. England: John Wiley & Sons Ltd.
- Freed PS, Neitman K. 1988. Notes on predation on the endangered Houston toad, *Bufo houstonensis*. *Texas Journal of Science* 40:454–456.
- Gottschalk JS. 1970. United States list of endangered native fish and wildlife. Federal Register 35:16047–16048. Available: http://ecos.fws.gov/docs/federal_register/federal_register_35_16047-16048.pdf (August 2011).
- Greuter KL. 2004. Early juvenile ecology of the endangered Houston toad, *Bufo houstonensis* (Anura: Bufonidae). Master's thesis. San Marcos: Texas State University.
- Greuter KL, Forstner MRJ. 2003. *Bufo houstonensis* (Houston toad) growth. *Herpetological Review* 34:355–356.

- Hanula JL, Wade DD. 2003. Influence of long-term dormant season burning and fire exclusion on ground-dwelling arthropod populations in a longleaf pine flatwoods ecosystem. *Forest Ecology and Management* 175:163–184.
- Hastie T, Tibshirani R. 1990. *Generalized additive models*. England: Chapman and Hall.
- Hatch AC, Belden LK, Scheessele E, Blaustein AR. 2001. Juvenile amphibians do not avoid potentially lethal levels of urea on soil substrate. *Environmental Toxicology and Chemistry* 20:2328–2335.
- Hodgkins EJ. 1958. Effects of fire on undergrowth vegetation in upland southern pine forests. *Ecology* 39:36–46.
- Landers JL, Garner JA, McRae WA. 1980. Reproduction of gopher tortoises (*Gopherus polyphemus*) in southwestern Georgia. *Herpetologica* 36:353–361.
- Lofgren CS, Banks WA, Glancey BM. 1975. Biology and control of imported fire ants. *Annual Review of Entomology* 20:1–30.
- Lyle C, Fortune I. 1948. Notes on an imported fire ant. *Journal of Economic Entomology* 41:833–834.
- Morrison LW. 2002. Long-term impacts of an arthropod-community invasion by the imported fire ant, *Solenopsis invicta*. *Ecology* 83:2337–2345.
- Mueller JM, Dabbert CB, Demarais S, Forbes AR. 1999. Northern bobwhite chick mortality caused by red imported fire ants. *Journal of Wildlife Management* 63:1291–1298.
- Norton AA. 2003. Integrating prescribed burning and insecticide to reduce fire ant impacts on bobwhite chicks. Master's thesis. Lubbock: Texas Tech University.

- Pedersen EK, Grant WE, Longnecker MT. 1996. Effects of red imported fire ants on newly-hatched northern bobwhite. *Journal of Wildlife Management* 60:164–169.
- Plowes RM, Dunn JG, Gilbert LE. 2007. The urban fire ant paradox: native fire ants persist in an urban refuge while invasive fire ants dominate natural habitats. *Biological Invasions* 9:825–836.
- Porter SD. 2010. Distribution of the formosa strain of the fire ant decapitating fly *Pseudacteon curvatus* (Diptera: Phoridae) three and a half years after releases in north Florida. *Florida Entomologist* 93:107–112.
- Porter SD, Savignano DA. 1990. Invasion of polygyne fire ants decimates native ants and disrupts arthropod community. *Ecology* 71:2095–2106.
- Porter SD, Tschinkel WR. 1987. Foraging in *Solenopsis invicta* (Hymenoptera: Formicidae): effects of weather and season. *Environmental Entomology* 16:802–808.
- Relyea RA. 2003. Predator cues and pesticides: a double dose of danger for amphibians. *Ecological Applications* 13:1515–1521.
- Sokal RR, Rohlf FJ. 1995. *Biometry: the principles and practice of statistics in biological research*. 3rd edition. New York: W. H. Freeman and Company.
- Sparks JC, Masters RE, Engle DM, Palmer MW, Bukenhofer GA. 1998. Effects of late growing-season and late dormant-season prescribed fire on herbaceous vegetation in restored pine-grassland communities. *Journal of Vegetation Science* 9:133–142.
- Stewart-Oaten A, Bence JR, Osenberg CW. 1992. Assessing effects of unreplicated perturbations: no simple solutions. *Ecology* 73:1396–1404.

- Stewart-Oaten A, Murdoch WW, Parker KR. 1986. Environmental impact assessment: “pseudoreplication” in time? *Ecology* 67:929–940.
- Stiles, JH, Jones RH. 1998. Distribution of the red imported fire ant, *Solenopsis invicta*, in road and powerline habitats. *Landscape Ecology* 13:335–346.
- Storrs Méndez SI, Tillitt DE, Rittenhouse TAG, Semlitsch RD. 2009. Behavioral response and kinetics of terrestrial atrazine exposure in American toads (*Bufo americanus*). *Archives of Environmental Contamination and Toxicology* 57:590–597.
- Stuble KL, Kirkman LK, Carroll CR. 2009. Patterns of abundance of fire ants and native ants in a native ecosystem. *Ecological Entomology* 34:520–526.
- Taber SW, Fleenor SB. 2003. *Insects of the Texas Lost Pines*. College Station: Texas A&M University Press.
- Todd, BD, Rothermel BB, Reed RN, Luhring TM, Schlatter K, Trenkamp L, Gibbons JW. 2008. Habitat alteration increases invasive fire ant abundance to the detriment of amphibians and reptiles. *Biological Invasions* 10:539–546.
- Tschinkel WR. 1988. Distribution of the fire ants *Solenopsis invicta* and *S. geminata* (Hymenoptera: Formicidae) in northern Florida in relation to habitat and disturbance. *Annals of the Entomological Society of America* 81:76–81.
- Tukey, JW. 1949. One degree of freedom for non-additivity. *Biometrics* 5:232–242.
- U.S. Fish and Wildlife Service. 1984. Houston toad recovery plan. Albuquerque, New Mexico: U.S. Fish and Wildlife Service. Available: <http://ecos.fws.gov/speciesProfile/profile/speciesProfile.action?sPCODE=D004>.

- Vogt JT, Allen ML, Wallet B, Boykin D, Smith WA. 2009. Distribution patterns of imported fire ants (Hymenoptera: Formicidae) on a sheep and goat farm in Oklahoma. *Environmental Entomology* 38:551–560.
- Ward PS. 2005. A synoptic review of the ants of California (Hymenoptera:Formicidae). *Zootaxa* 936:1–68.
- Williams DF, Collins HL, Oi DH. 2001. The red imported fire ant (Hymenoptera: Formicidae): an historical perspective of treatment programs and the development of chemical baits for control. *American Entomologist* 47:146–159.
- Wood, SN. 2004. Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association* 99:673–686.
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. 2009. *Mixed effects models and extensions in ecology with R*. New York: Springer Science+Business Media.

CHAPTER VI

WILDFIRE IMPACTS ON RED IMPORTED FIRE ANT CAPTURES IN THE LOST PINES ECOREGION OF TEXAS

Format follows the guidelines for the Journal of Fish and Wildlife Management

VI. Wildfire Impacts on Red Imported Fire Ant Captures in the Lost Pines Ecoregion of Texas

Abstract

The impacts of high intensity wildfires on red imported fire ant (*Solenopsis invicta*) abundances in forest systems are currently unknown. The purpose of this study was to evaluate the effects of a high intensity wildfire around pond edges on red imported fire ant captures in the Lost Pines ecoregion of Texas, which provides habitat for most of the remaining endangered Houston toads (*Bufo houstonensis*). The red imported fire ant is a known predator of Houston toads, and thus there is interest in understanding the potential and realized impacts of this species on Houston toad survivorship. Our results suggested the wildfire did not directly impact captures-per-unit-effort, but it had indirect positive impacts through reductions in canopy cover due to overstory tree mortality. The results of this study indicate that both area occupied by red imported fire ants and site-specific red imported fire ant densities could increase dramatically in the Lost Pines as a result of the wildfire and subsequent human-based recovery efforts.

Keywords: Canopy cover; Houston toad; Lost Pines; red imported fire ant; Texas; wildfire.

Introduction

The red imported fire ant (RIFA; *Solenopsis invicta*) is an invasive species in the southern and southeastern U.S. It was introduced to Alabama in the 1930s, and has since expanded east to North Carolina and west to central Texas (Callcott and Collins 1996), and has been documented in California (Ward 2005). Negative ecological impacts of RIFA invasion have been demonstrated for a wide range of vertebrates and invertebrates (Porter and Savignano 1990; Allen et al. 1994; Allen et al. 2004; Stuble et al. 2009; Diffie et al. 2010; Epperson and Allen 2010). Among vertebrates, juveniles tend to be particularly vulnerable to RIFA predation (Landers et al. 1980; Freed and Neitman 1988; Pedersen et al. 1996).

The red imported fire ant appeared in Bastrop County, central Texas between 1973 and 1977, and has since become well established throughout the Lost Pines ecoregion (Cokendolpher and Phillips 1989; Taber and Fleenor 2003), which has historically been the last remaining stronghold for the federally endangered Houston toad (*Bufo [Anaxyrus] houstonensis*; U.S. Endangered Species Act [ESA] 1973, as amended). This species is a known predator of juvenile Houston toads (Freed and Neitman 1988), and thus there is interest in understanding the potential and realized impacts of RIFA on Houston toad survivorship.

Brown et al. (2012) recently conducted a study in the Lost Pines aimed at delineating annual RIFA activity patterns, elucidating the relationship between overstory canopy cover and RIFA captures, and evaluating the effects of low intensity prescribed fire around pond edges (where juvenile Houston toads are vulnerable to predation after emergence) on RIFA captures. Brown et al. (2012) found that annual RIFA activity was

highest between May and October, which includes the period in which juvenile Houston toads disperse from ponds. They also found a strong inverse relationship between RIFA captures and canopy cover near pond edges. It is possible that greater sunlight exposure, and thus greater heat, in open-canopied environments results in more suitable habitat for RIFA (Vogt et al. 2003). Lastly, Brown et al. (2012) detected no effects of low intensity prescribed fire on RIFA captures, indicating no sustained direct impact of fire on the species. However, the authors noted that research on high intensity fire impacts was warranted, given the greater potential for habitat alteration consequent of hotter fires.

On 4 September 2011 a high intensity wildfire began in the Lost Pines. The fire was unstoppable due to wind gusts in excess of 58 kph caused by tropical storm Lee, coupled with extreme drought conditions in central Texas. After 18 days the fire was 95% contained, with the total burn area encompassing 13,406 ha (ca. 39% of the Lost Pines ecoregion). Seven of the 10 ponds used in the Brown et al. (2012) study burned in the wildfire, allowing us the opportunity to assess the impacts of a high intensity wildfire on RIFA in the Lost Pines ecoregion using 7 burned and 3 unburned ponds. Given the inverse relationship between canopy cover and RIFA captures found in the previous study, we hypothesized that RIFA captures would increase at ponds with substantial tree mortality caused by the wildfire.

Study Site

We conducted this study on the 1,948 ha Griffith League Ranch (GLR). Vegetation on the GLR is typical of the Lost Pines ecoregion, with an overstory dominated by loblolly pine (*Pinus taeda*), post oak (*Quercus stellata*), blackjack oak

(*Quercus marilandica*), and eastern red cedar (*Juniperus virginiana*), and an understory dominated by yaupon holly (*Ilex vomitoria*), American beautyberry (*Callicarpa americana*), and farkleberry (*Vaccinium arboreum*). The GLR contains three permanent ponds (i.e., ponds have not dried in at least 12 years), 10 semi-permanent ponds (i.e., ponds typically dry several times per decade), and dozens of ephemeral ponds that hold water for days to months annually depending on rainfall. For this study we used the 7 burned and 3 unburned ponds with RIFA sampling data prior to the wildfire, all of which are known Houston toad breeding ponds. Three of the ponds were ephemeral, six were semi-permanent, and one was permanent.

Methods

The pre-wildfire RIFA capture data used in this study was collected between 26 September 2009 and 10 October 2010, with the area around each of the 10 ponds sampled a total of 27 times (Brown et al. 2012). We collected post-wildfire RIFA capture data for this study between 25 September 2011 and 4 August 2012, sampling the area around each of the 10 ponds a total of 11 times. To maximize comparability we used an identical sampling design to that of Brown et al. (2012). Days between sampling ranged from 15 to 78, with a mean of 31 days between samples. We used a standard bait cup method for sampling ants (Porter and Tschinkel 1987; Mueller et al. 1999), which consisted of half a Vienna sausage placed in a 9-cm-diameter round plastic dish. We placed between 4 and 14 dishes around the perimeter of each pond, depending on pond size (i.e., perimeter length). During each sampling event, we placed dishes 1–2 m from the pond edge at 3–4 m intervals, and allowed them to attract ants for 30 minutes. After 30 minutes we

collected ants from each dish and euthanized them by freezing. We removed native ants from samples and counted the number of RIFA individuals collected in each bait cup at each pond on each sampling date. Because the number of bait cups varied by pond and sampling date, we calculated captures-per-unit-effort (CPUE; defined as the number of captures / number of bait cups), and used CPUE as our response variable.

The pre-wildfire canopy cover data used in this study was collected between 20 July 2008 and 16 April 2010, with percent canopy cover around each pond estimated between 4 and 24 times, depending on hydroperiod status at the time of sampling (i.e., estimates were not taken when ponds did not contain water; Brown et al. 2012). We estimated percent canopy cover around each pond four times between 18 April 2012 and 27 June 2012. For both studies, we estimated percent canopy cover using a spherical densiometer (Forestry Suppliers Inc., Jackson, MS) at two to six randomly selected points at the pond edge, with higher numbers of estimation points corresponding to larger ponds. We then averaged the estimates at each pond per sampling date. For this study we used the difference between the mean pre-wildfire and post-wildfire canopy cover estimates as a predictor of post-wildfire RIFA CPUE.

Statistical analyses

Our data included both spatial (multiple samples were taken from around the same pond) and temporal (the same ponds were sampled repeatedly) non-independence. To account for this we used a mixed effects modeling approach for our analyses, treating within-day sample nested within pond as a random effect. Preliminary analyses indicated a strong positive relationship between air temperature and RIFA captures, and thus we

included mean daily temperature as a covariate in our final analyses. We obtained temperature data from the Austin-Bergstrom International Airport weather station, located approximately 36 km from the GLR (station number: 410429).

We initially tested interactions among and between treatment (control/wildfire), time (pre-burn/post-burn), and canopy cover change (change in mean % canopy cover between the pre-wildfire and post-wildfire sampling periods). This analysis indicated an interaction between canopy cover change and time, and changes in canopy cover were not consistent among ponds due to variability in tree mortality caused by the wildfire, as well as pond size (canopy cover at the edge of ponds typically increases as pond size increases). Thus, we analyzed each pond individually using mixed effects models to determine if RIFA CPUE differed by burn status. We investigated assumptions of normality and homoscedasticity using graphical diagnostics plots (Zuur et al. 2009). Because the data appeared to violate the assumption of normality, we transformed RIFA CPUE using the arcsinh (i.e., inverse hyperbolic sine) transformation (Fowler et al. 1998). We performed these analyses with the program R (R Version 2.10.1, www.r-project.org) using the nlme (Pinheiro et al. 2009) package. We considered effects to be significant at $\alpha = 0.05$. Finally, we tested the relationship between changes in RIFA CPUE between sampling periods and changes in mean % canopy cover between sampling periods using ordinary least squares (OLS) regression (Sokal and Rohlf 1995). For this analysis we used RIFA captures between May and early August (i.e., the peak RIFA activity months common to both the pre-wildfire and post-wildfire sampling periods). We considered effects to be significant at $\alpha = 0.05$.

Results

The initial analysis testing interactions among and between treatment, time, and canopy cover change indicated no treatment x time x canopy cover change interaction effect ($F_{1,365} = 0.03$, $P = 0.853$), no treatment x canopy cover change interaction effect ($F_{1,6} = 1.72$, $P = 0.237$), and no treatment x time interaction effect ($F_{1,365} = 0.02$, $P = 0.902$). However, we detected a time x canopy cover change interaction effect ($F_{1,365} = 27.52$; $P < 0.001$). These results indicate changes in CPUE between sampling periods (i.e., time) were not directly associated with the wildfire. Rather, CPUE differed between sampling periods, and the effect was influenced by changes in % canopy cover. The pond-specific tests indicated a time effect on RIFA CPUE at 5 of the 10 ponds, including 1 of 3 control ponds, and 4 of 7 wildfire ponds (Table 6.1). The time coefficients indicated that post-burn CPUE decreased at the control pond (Pond 6), decreased at two of the wildfire ponds (ponds 10 and 11), and increased at two of the wildfire ponds (ponds 13 and 14). The directional change in RIFA captures for all of these ponds was inversely related to canopy cover change (Pond 7 [+5.14%]; Pond 10 [+5.01%]; Pond 11 [+17.52%]; Pond 13 [-11.30%]; Pond 14 [-41.86%]). The OLS regression indicated that RIFA CPUE decreased with increased mean % canopy cover ($F_{1,7} = 13.17$; $P = 0.008$; $r^2 = 0.65$; Figure 6.1).

Table 6.1. Results from pond-specific tests of a burn status (pre-wildfire/post-wildfire) effect on red imported fire ant (RIFA; *Solenopsis invicta*) captures-per-unit-effort (CPUE) around pond edges in the Lost Pines ecoregion, Bastrop County, Texas, USA. We used mixed effects models to test for burn status effects, treating within-day sample nested within pond as a random effect, and we included mean daily temperature as a covariate. Pre-burn status coefficients show the direction of change in RIFA captures between sampling periods (pre-wildfire sampling period: September 2009 to October 2010; post-wildfire sampling period: September 2011 to August 2012), with positive coefficients indicating higher pre-wildfire RIFA captures.

Pond	Treatment	Pre-burn status coefficient ($\pm$ SE)	$F_{1,35}$	P
5	Control	0.53 (0.61)	0.76	0.388
6	Control	0.86 (0.34)	6.33	0.017
7	Control	0.81 (0.56)	2.07	0.159
9	Wildfire	0.85 (0.53)	2.57	0.118
10	Wildfire	1.14 (0.45)	6.55	0.015
11	Wildfire	1.27 (0.53)	5.79	0.022
13	Wildfire	-0.50 (0.17)	8.17	0.007
14	Wildfire	-1.46 (0.45)	10.68	0.002
15	Wildfire	-0.67 (0.50)	1.80	0.188
16	Wildfire	0.27 (0.41)	0.42	0.523




Figure 6.1. Ordinary least squares (OLS) regression assessing the relationship between differences in mean % canopy cover between the pre-wildfire (September 2009 to October 2010) and post-wildfire (September 2011 to August 2012) sampling periods, and red imported fire ant (RIFA; *Solenopsis invicta*) captures-per-unit-effort (CPUE) between May and early August (i.e., the peak RIFA activity months common to both the pre-wildfire and post-wildfire sampling periods) around pond edges in the Lost Pines ecoregion, Bastrop County, Texas, USA. The analysis indicated post-wildfire RIFA CPUE was typically higher around ponds where % canopy cover was lower during the post-wildfire sampling period, and typically lower around ponds where % canopy cover was higher during the post-wildfire sampling period ($F_{1,7} = 13.17$; $P = 0.008$). One pond (Pond 13) was removed from this analysis because we obtained no captures during the pre-burn sampling period and thus could not calculate the proportional change in CPUE. For this pond the mean % canopy cover change was -11.3% and the mean post-wildfire CPUE during the peak RIFA activity period was 21.4.

Discussion

We found that a high intensity wildfire did not have a direct impact on RIFA CPUE, which was the same conclusion found for low intensity prescribed fires in this system (Brown et al. 2012). Rather, CPUE differences between the two sampling periods were driven by changes in canopy cover. Our results indicated the wildfire indirectly benefited RIFA through overstory tree mortality (and thus canopy cover reduction), supporting our hypothesis based on previous research (Brown et al. 2012). We speculate that the two wildfire ponds with significant post-wildfire decreases in RIFA CPUE

actually positively impacted RIFA, but the effect was masked by an increase in canopy cover at these ponds due to heavy rainfall in fall 2011 that resulted in the pond edges reaching into the adjacent upland forest canopy.

Our primary motivation for conducting research on RIFA is to assist with management activities that benefit recovery of the Houston toad. The results of this study are cause for concern in that the wildfire burned a significant portion of the Lost Pines ecoregion (ca. 39%), with an estimated overstory tree loss of 78% across the burned area (Lost Pines Recovery Team 2011). Based on 13 randomly placed 20 m x 50 m vegetation plots on the GLR within the wildfire zone, the mean overstory tree loss was 81.9%, with a corresponding 30.2% decrease in canopy cover (D.J. Brown, unpublished data). Further, the Federal Emergency Management Agency (FEMA), Bastrop County, and local landowners are currently removing a substantial amount of dead trees due to conflicts with human-based recovery needs. Red imported fire ants are known to prefer forest gaps (Colby and Prowell 2006), and to use openings such as roads and powerline cuts as expansion corridors (Tschinkel 1988; Stiles and Jones 1998; Todd et al. 2008). Thus, the mortality and subsequent removal of trees in the Lost Pines has the potential to dramatically increase both area occupied by RIFA and site-specific RIFA densities in the Lost Pines.

The research conducted in this study and Brown et al. (2012) has increased our understanding of the spatial and temporal habitat use of RIFA in the Lost Pines ecoregion. It is clear that these patterns overlap with those of the Houston toad (Brown et al. 2012), and thus RIFA has the potential to be a serious threat to the persistence of the Houston toad. However, although we know that RIFA prey upon juvenile Houston toads

(Freed and Neitman 1988), and we have evidence that they also prey upon adult Houston toads (M. C. Jones, Texas State University, personal communication), we do not have a good understanding of their impacts on Houston toad survivorship rates. Thus, it is possible that RIFA predation is partially or entirely compensatory (i.e., some or all of the mortality caused by RIFA could be substituting for mortality that would have occurred due to other factors). However, we believe the increased prevalence of RIFA in the Lost Pines as a result of post-wildfire canopy cover loss will likely add additional stress to Houston toad persistence in the ecoregion. Fortunately, RIFA control through insecticides, if warranted, is a viable option for Houston toad recovery efforts.

Acknowledgements

This study was supported by the Texas Parks and Wildlife Department and U.S. Fish and Wildlife Service through a traditional Section 6 grant, and relied upon methods developed with funding from the Texas Department of Transportation. The Capitol Area Council of the Boy Scouts of America provided access to the Griffith League Ranch, and we appreciate their continuing support of our research. We thank B. DeVolld, W. DeVolld, and L. Eddins for their assistance with data collection for the pre-wildfire data set.

References

Allen CR, Demarais S, Lutz RS. 1994. Red imported fire ant impact on wildlife- an overview. *Texas Journal of Science* 46:51–59.

- Allen CR, Epperson DM, Garmestani AS. 2004. Red imported fire ant impacts on wildlife: a decade of research. *American Midland Naturalist* 152:88–103.
- Brown DJ, DeVold B, Forstner MRJ. 2012. Fire ants in Houston toad habitat: annual activity and responses to canopy cover and fire. *Journal of Fish and Wildlife Management* 3:142–149.
- Callcott A-MA, Collins HL. 1996. Invasion and range expansion of imported fire ants (Hymenoptera: Formicidae) in North America from 1918-1995. *Florida Entomologist* 79:240–251.
- Cokendolpher JC, Phillips SA Jr. 1989. Rate of spread of the red imported fire ant, *Solenopsis invicta* (Hymenoptera: Formicidae), in Texas. *Southwestern Naturalist* 34:443–449.
- Colby D, Prowell D. 2006. Ants (Hymenoptera: Formicidae) in wet longleaf pine savannas in Louisiana. *Florida Entomologist* 89:266–269.
- Diffie S, Miller J, Murray K. 2010. Laboratory observations of red imported fire ant (Hymenoptera: Formicidae) predation on reptilian and avian eggs. *Journal of Herpetology* 44:294–296.
- Epperson DM, Allen CR. 2010. Red imported fire ant impacts on upland arthropods in southern Mississippi. *American Midland Naturalist* 163:54–63.
- [ESA] US Endangered Species Act of 1973, as amended, Pub. L. No. 93-205, 87 Stat. 884 (Dec. 28, 1973). Available: <http://www.fws.gov/endangered/esa-library/pdf/ESAall.pdf>.
- Fowler JF, Cohen L, Jarvis P. 1998. *Practical statistics for field biology*. 2nd edition. England: John Wiley & Sons Ltd.

- Freed PS, Neitman K. 1988. Notes on predation on the endangered Houston toad, *Bufo houstonensis*. Texas Journal of Science 40:454–456.
- Landers JL, Garner JA, McRae WA. 1980. Reproduction of gopher tortoises (*Gopherus polyphemus*) in southwestern Georgia. Herpetologica 36:353–361.
- Lost Pines Recovery Team. 2011. Bastrop County complex fire, Lost Pines region, resources assessment and response report. Lost Pines Recovery Team. Available: http://co.bastrop.tx.us/bcdisaster/files/PDF_Files/LPRT%20Recovery%20Report%20FINAL%201-10-11%20reduced.pdf (November 2012).
- Mueller JM, Dabbert CB, Demarais S, Forbes AR. 1999. Northern bobwhite chick mortality caused by red imported fire ants. Journal of Wildlife Management 63:1291–1298.
- Pedersen EK, Grant WE, Longnecker MT. 1996. Effects of red imported fire ants on newly-hatched northern bobwhite. Journal of Wildlife Management 60:164–169.
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. 2009. nlme: linear and non-linear mixed effects models. R package version 3.1–96. Available: <http://cran.r-project.org/web/packages/nlme/nlme.pdf> (November 2012).
- Porter SD, Savignano DA. 1990. Invasion of polygyne fire ants decimates native ants and disrupts arthropod community. Ecology 71:2095–2106.
- Porter SD, Tschinkel WR. 1987. Foraging in *Solenopsis invicta* (Hymenoptera: Formicidae): effects of weather and season. Environmental Entomology 16:802–808.

- Sokal RR, Rohlf FJ. 1995. Biometry: the principles and practice of statistics in biological research. 3rd edition. New York: W. H. Freeman and Company.
- Stiles, JH, Jones RH. 1998. Distribution of the red imported fire ant, *Solenopsis invicta*, in road and powerline habitats. *Landscape Ecology* 13:335–346.
- Stuble KL, Kirkman LK, Carroll CR. 2009. Patterns of abundance of fire ants and native ants in a native ecosystem. *Ecological Entomology* 34:520–526.
- Taber SW, Fleenor SB. 2003. Insects of the Texas Lost Pines. College Station: Texas A&M University Press.
- Todd, BD, Rothermel BB, Reed RN, Luhring TM, Schlatter K, Trenkamp L, Gibbons JW. 2008. Habitat alteration increases invasive fire ant abundance to the detriment of amphibians and reptiles. *Biological Invasions* 10:539–546.
- Tschinkel WR. 1988. Distribution of the fire ants *Solenopsis invicta* and *S. geminata* (Hymenoptera: Formicidae) in northern Florida in relation to habitat and disturbance. *Annals of the Entomological Society of America* 81:76–81.
- Vogt JT, Smith WA, Grantham RA, Wright RE. 2003. Effect of temperature and season on foraging activity of red imported fire ants (Hymenoptera: Formicidae) in Oklahoma. *Environmental Entomology* 32:447–451.
- Ward PS. 2005. A synoptic review of the ants of California (Hymenoptera:Formicidae). *Zootaxa* 936:1–68.
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. 2009. Mixed effects models and extensions in ecology with R. New York: Springer Science+Business Media.

CHAPTER VII

SUMMARY AND CONCLUSIONS

Format follows the guidelines for the Journal of Wildlife Management

VII. Summary and Conclusions

This dissertation provided useful information for land managers in the Lost Pines ecoregion of Texas concerning the use of fire for habitat management, as well as post-fire management in the high severity wildfire zone. These studies served to increase our knowledge of fire impacts on amphibians and reptiles, the least studied major terrestrial vertebrate groups in relation to fire research (Russell et al. 1999, Pilliod et al. 2003), and red imported fire ants (*Solenopsis invicta*), a species for which almost no information existed concerning fire impacts. Further, this dissertation provided important information regarding effects of fire severity on several ecosystem components within a single ecosystem, which is currently a topic of interest given that fire severity is possibly the most important single factor influencing ecosystem responses to fire (Knapp et al. 2009). Finally, through the completion of these studies several relevant unanswered questions became apparent. In the following paragraphs I provide a brief summary of major findings of this dissertation research, address implications of those findings, and discuss future research opportunities that would improve our knowledge concerning the impacts of fire on ecosystems.

Chapters II and III investigated immediate and short-term impacts of moderate and high severity fire on herpetofauna. Neither study indicated that direct mortality was significant. This is the typical conclusion of previous prescribed fire research (e.g., Ford et al. 1999, Ruthven et al. 2008), but direct mortality impacts from high severity wildfires were virtually unknown, and this research indicated herpetofauna can survive even high

severity fires. However, it is important to note that the Lost Pines ecoregion is characterized by loose sandy soils (Baker et al. 1979), a substrate conducive to rapid burrowing. Thus, it would not be surprising if direct mortality of herpetofauna is typically greater in regions containing harder soils.

The results of Chapter II not only indicated that direct mortality was low, but also that survivorship of juvenile amphibians may have increased following a moderate severity summer prescribed fire, and the potential increase in survivorship could have been related to a reduction in arthropod predation. However, my study was not designed to explicitly test that relationship, and thus the evidence is both correlative and weak. Effects of terrestrial predation on amphibian population dynamics are largely unknown, and thus this is an area of future research with great potential.

The results of Chapter III indicated the post-wildfire landscape provided suitable habitat for herpetofauna ca. 6 months after the wildfire. This is directly opposed to the general assumption of wildlife managers in the area that the fire killed the majority of herpetofauna and destroyed their habitat (Lost Pines Recovery Team 2011). Thus, management and restoration initiatives in the wildfire zone should operate under the assumption that the wildfire zone is currently providing suitable habitat for wildlife. For example, when planning significant habitat alterations (e.g., clearcutting), an approach that preserves adjacent non-manipulated patches could provide temporary refuge zones during the early stages of forest restoration, when the altered habitat would be unsuitable for some species, such as the endangered Houston toad (*Bufo* [*Anaxyrus*] *houstonensis*).

An additional interesting result of Chapter III was that movement-rates of Hurter's spadefoot toads (*Scaphiopus hurterii*) appeared to increase following the wildfire. To my knowledge this is the first study that investigated the impacts of fire on amphibian movement-rates. However, movement was not assessed directly, but rather indirectly through detection probabilities for captures. Thus, validation of this phenomenon would require direct assessments of movement-rates (e.g., using radio telemetry). This is a topic worth investigating, as there could be multiple reasons why movement-rates would increase following high severity fires. For example, significant reduction of fine and coarse fuel loads could simply make it easier to traverse the landscape, which could benefit amphibians migrating to breeding ponds by reducing energy expenditure. Conversely, increases in movement during foraging events could indicate a reduction in habitat quality, as greater movement is required to obtain sufficient food resources (e.g., arthropods). A detailed study that incorporated movement purposes would provide extremely valuable, and more causative-based, information concerning fire impacts on amphibians.

Chapter IV investigated the influence of fire severity on responses of ecosystem components to fire. The study indicated that fire severity was an important factor, and the influence was related to magnitude, but not direction, of effects. For some components (e.g., pond nutrient levels) the magnitude effect was clear, whereas for others (e.g., species composition of understory vegetation) it was dichotomous in that no effect was apparent for low severity fire and a strong effect was apparent for high severity fire. It is likely that most terrestrial responses were associated with reduction of both living and

dead organic matter, which was substantially greater for high severity fire. Thus, it is possible that implementing multiple prescribed burns could result in habitat effects similar to those induced by high severity wildfire, particularly for forest structure characteristics and diversity of vegetation.

An important finding of Chapter IV for managers in the Lost Pines was that mortality of loblolly pine (*Pinus taeda*) trees was nearly 100% in the high severity wildfire zone, and subsequently loblolly pine tree recruitment was low. In contrast, mortality of eastern red cedar (*Juniperus virginiana*) trees was lower, and although mortality of post oak (*Quercus stellata*) trees was high, the species was re-establishing itself ca. 1 year after the wildfire through basal sprouting following top-kill. Thus, restoration of the Lost Pines will require significant active management through reseedling of loblolly pines, with the alternative being a shift to hardwood-dominated forest patches in the high severity wildfire zone. Because the former loblolly pine-dominated patches are both ecologically and culturally significant, a major reseedling initiative throughout the high severity wildfire zone is advised.

Chapters V and VI investigated temporal and spatial characteristics of the invasive red imported fire ant (RIFA; *Solenopsis invicta*) around forested ponds in the Lost Pines ecoregion, and the impacts of fire on this species. The results indicated that peak annual RIFA activity coincides with the period when juvenile Houston toads are found at high densities around breeding ponds, and thus are particularly vulnerable to population-level impacts of predation. Further, there was a strong inverse relationship between RIFA captures (an assumed proxy for abundance) and overstory canopy cover.

High severity fire appeared to positively affect RIFA through reduction in canopy cover. This is a concerning result given the substantial tree mortality across the high severity wildfire zone, and thus the potential for increases in RIFA abundance and distribution in the Lost Pines ecoregion. Fortunately, RIFA control through chemical means is a viable management option, and its use is worth considering at active Houston toad breeding ponds. However, we currently do not have a clear understanding of the population-level impacts of RIFA predation, and future research on this topic would not only benefit managers in the Lost Pines ecoregion, but also other regions where this invasive species could be negatively impacting amphibian populations.

LITERATURE CITED

- Baker, F. E. 1979. Soil survey of Bastrop County, Texas. U.S. Department of Agriculture, Soil Conservation Service, Washington, D.C., USA.
- Ford, W.M., M. A. Menzel, D. W. McGill, J. Laerm, and T. S. McCay. 1999. Effects of a community restoration fire on small mammals and herpetofauna in the southern Appalachians. *Forest Ecology and Management* 114:233–243.
- Gresswell, R. E. 1999. Fire and aquatic ecosystems in forested biomes of North America. *Transactions of the American Fisheries Society* 128:193–221.
- Knapp, E. E., B. L. Estes, and C. N. Skinner. 2009. Ecological effects of prescribed fire season: a literature review and synthesis for managers. U. S. Department of Agriculture General Technical Report PSW–GTR–224, Redding, California, USA.

- Lost Pines Recovery Team. 2011. Bastrop County complex fire, Lost Pines region, resources assessment and response report. Lost Pines Recovery Team, Bastrop, Texas, USA.
- Russell, K. R., D. H. Van Lear, and D. C. Guynn, Jr. 1999. Prescribed fire effects on herpetofauna: review and management implications. *Wildlife Society Bulletin* 27:374–384.
- Ruthven, D. C., III, R. T. Kzmaier, and M. W. Janis. 2008. Short-term response of herpetofauna to various burning regimes in the south Texas plains. *Southwestern Naturalist* 53:480–487.

VITA

Donald J. Brown was born on 8 June 1981 in Alameda, California to James and Debra Brown. Donald received a B.Sc. in Fisheries & Wildlife-Wildlife Specialization from the University of Minnesota-Twin Cities in May 2007, and a M.Sc. in Wildlife Ecology from Texas State University-San Marcos in December 2008. He entered the Aquatic Resources Ph.D. program at Texas State University-San Marcos in January 2009. While at Texas State Donald worked as a National Science Foundation- Project Flowing Waters fellow, a lab instructor for Wildlife Management, Genetics, and Functional Biology, a research assistant for Michael Forstner, and as a field-based consultant for Houston toad population monitoring and impact mitigation activities. Donald has currently written 13 peer-reviewed publications, is a co-author of 9 others, and has presented his research at 13 regional, national, and international conferences, including World Congress of Herpetology and Ecological Society of America.

Permanent Email Address: djb.ecology@gmail.com

This dissertation was typed by Donald J. Brown